



wwPDB X-ray Structure Validation Summary Report ⓘ

Jan 27, 2024 – 03:06 PM EST

PDB ID : 1AYR
Title : ARRESTIN FROM BOVINE ROD OUTER SEGMENTS
Authors : Granzin, J.; Wilden, U.; Choe, H.-W.; Labahn, J.; Krafft, B.; Bueldt, G.
Deposited on : 1997-11-10
Resolution : 3.30 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	1205 (3.34-3.26)
Ramachandran outliers	138981	1183 (3.34-3.26)
Sidechain outliers	138945	1182 (3.34-3.26)
RSRZ outliers	127900	1115 (3.34-3.26)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	368	5% (Poor fit) 40% (0 outliers) 45% (1 outlier) 14% (2 outliers) 0% (3+ outliers)
1	B	368	6% (Poor fit) 43% (0 outliers) 40% (1 outlier) 15% (2 outliers) 0% (3+ outliers)
1	C	368	5% (Poor fit) 41% (0 outliers) 45% (1 outlier) 13% (2 outliers) 0% (3+ outliers)
1	D	368	7% (Poor fit) 42% (0 outliers) 42% (1 outlier) 15% (2 outliers) 0% (3+ outliers)

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 11743 atoms, of which 0 are hydrogens and 0 are deuteriums.


In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ARRESTIN.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	368	2882	1841	488	543	10	0	0	0
1	B	363	2847	1821	482	534	10	0	0	0
1	C	368	2882	1841	488	543	10	0	0	0
1	D	363	2847	1821	482	534	10	0	0	0

- Molecule 2 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	99	Total	O	0	0
			99	99		
2	B	46	Total	O	0	0
			46	46		
2	C	78	Total	O	0	0
			78	78		
2	D	62	Total	O	0	0
			62	62		



• Molecule 1: ARRESTIN



• Molecule 1: ARRESTIN



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	169.17Å 185.53Å 90.93Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	130.00 – 3.30 26.98 – 3.30	Depositor EDS
% Data completeness (in resolution range)	95.1 (130.00-3.30) 41.9 (26.98-3.30)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	0.17	Depositor
$\langle I/\sigma(I) \rangle$ ¹	11.37 (at 3.31Å)	Xtrriage
Refinement program	X-PLOR 3.851	Depositor
R, R_{free}	0.254 , 0.316 0.254 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å ²)	64.0	Xtrriage
Anisotropy	0.742	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 38.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.88	EDS
Total number of atoms	11743	wwPDB-VP
Average B, all atoms (Å ²)	27.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.62% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.31	0/2940	0.57	3/3987 (0.1%)
1	B	0.30	0/2905	0.57	1/3941 (0.0%)
1	C	0.30	0/2940	0.57	3/3987 (0.1%)
1	D	0.30	0/2905	0.57	2/3941 (0.1%)
All	All	0.30	0/11690	0.57	9/15856 (0.1%)

There are no bond length outliers.

The worst 5 of 9 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	9	ASN	N-CA-C	-7.17	91.65	111.00
1	A	9	ASN	N-CA-C	-7.05	91.97	111.00
1	C	9	ASN	N-CA-C	-6.86	92.47	111.00
1	B	9	ASN	N-CA-C	-6.86	92.48	111.00
1	A	7	ALA	N-CA-C	-5.22	96.90	111.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2882	0	2971	190	0
1	B	2847	0	2937	164	0
1	C	2882	0	2971	174	0
1	D	2847	0	2937	166	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	A	99	0	0	17	0
2	B	46	0	0	8	0
2	C	78	0	0	14	0
2	D	62	0	0	11	0
All	All	11743	0	11816	680	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 29.

The worst 5 of 680 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:10:HIS:O	1:B:11:VAL:HG23	1.68	0.94
1:A:2:LYS:HG2	1:A:29:ARG:HH11	1.36	0.90
1:D:2:LYS:HG2	1:D:29:ARG:HH11	1.37	0.89
1:B:2:LYS:HG2	1:B:29:ARG:HH11	1.35	0.88
1:B:91:PHE:HB3	1:B:92:PRO:HD3	1.56	0.88

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	366/368 (100%)	314 (86%)	40 (11%)	12 (3%)	4	22
1	B	361/368 (98%)	310 (86%)	40 (11%)	11 (3%)	4	24
1	C	366/368 (100%)	313 (86%)	41 (11%)	12 (3%)	4	22
1	D	361/368 (98%)	309 (86%)	41 (11%)	11 (3%)	4	24
All	All	1454/1472 (99%)	1246 (86%)	162 (11%)	46 (3%)	4	22

5 of 46 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	4	ASN
1	A	11	VAL
1	A	35	VAL
1	A	341	GLU
1	B	4	ASN

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	327/327 (100%)	247 (76%)	80 (24%)	0	2
1	B	324/327 (99%)	244 (75%)	80 (25%)	0	2
1	C	327/327 (100%)	245 (75%)	82 (25%)	0	2
1	D	324/327 (99%)	246 (76%)	78 (24%)	0	2
All	All	1302/1308 (100%)	982 (75%)	320 (25%)	0	2

5 of 320 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	318	LYS
1	D	182	ARG
1	C	344	SER
1	D	78	SER
1	D	246	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 23 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	155	HIS
1	D	34	HIS
1	C	301	HIS
1	D	89	GLN
1	A	357	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	368/368 (100%)	-0.12	17 (4%) 32 30	2, 10, 70, 99	0
1	B	363/368 (98%)	0.09	22 (6%) 21 20	2, 29, 83, 97	0
1	C	368/368 (100%)	-0.08	18 (4%) 29 27	2, 13, 75, 99	0
1	D	363/368 (98%)	0.13	27 (7%) 14 14	2, 25, 83, 99	0
All	All	1462/1472 (99%)	0.00	84 (5%) 23 23	2, 17, 81, 99	0

The worst 5 of 84 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	158	ASP	8.7
1	C	76	GLY	7.1
1	A	75	MET	6.6
1	C	75	MET	6.5
1	D	98	GLY	6.4

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.