



wwPDB X-ray Structure Validation Summary Report ⓘ

Oct 5, 2024 – 04:33 pm BST

PDB ID : 4BVP
Title : Legionella pneumophila NTPDase1 crystal form II (closed) in complex with heptamolybdate and octamolybdate
Authors : Zebisch, M.; Schaefer, P.; Straeter, N.
Deposited on : 2013-06-27
Resolution : 1.49 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.4, CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	3.0
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4	:	9.0.003 (Gargrove)
Density-Fitness	:	1.0.11
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.39


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.49 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	3717 (1.50-1.50)
Clashscore	180529	4048 (1.50-1.50)
Ramachandran outliers	177936	3970 (1.50-1.50)
Sidechain outliers	177891	3967 (1.50-1.50)
RSRZ outliers	164620	3718 (1.50-1.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	368	<div> <div>%</div> <div> <div></div> <div>91%</div> <div>7% ..</div> </div> </div>
1	B	368	<div> <div>2%</div> <div> <div></div> <div>93%</div> <div>6% .</div> </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
12	8M0	A	1417	-	-	X	-
12	8M0	B	1403	-	-	X	-
7	M27	A	1408	-	-	X	-
9	6LL	B	1413	-	-	X	-

2 Entry composition

There are 13 unique types of molecules in this entry. The entry contains 6758 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ECTONUCLEOSIDE TRIPHOSPHATE DIPHOSPHOHYDROLASE I.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	365	Total	C	N	O	S	0	21	0
			3034	1931	501	586	16			
1	B	365	Total	C	N	O	S	0	14	0
			2990	1902	493	578	17			


There are 22 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	34	MET	-	expression tag	UNP Q5ZUA2
A	394	LEU	-	expression tag	UNP Q5ZUA2
A	395	GLU	-	expression tag	UNP Q5ZUA2
A	396	HIS	-	expression tag	UNP Q5ZUA2
A	397	HIS	-	expression tag	UNP Q5ZUA2
A	398	HIS	-	expression tag	UNP Q5ZUA2
A	399	HIS	-	expression tag	UNP Q5ZUA2
A	400	HIS	-	expression tag	UNP Q5ZUA2
A	401	HIS	-	expression tag	UNP Q5ZUA2
A	137	ASP	GLU	conflict	UNP Q5ZUA2
A	149	VAL	ALA	conflict	UNP Q5ZUA2
B	34	MET	-	expression tag	UNP Q5ZUA2
B	394	LEU	-	expression tag	UNP Q5ZUA2
B	395	GLU	-	expression tag	UNP Q5ZUA2
B	396	HIS	-	expression tag	UNP Q5ZUA2
B	397	HIS	-	expression tag	UNP Q5ZUA2
B	398	HIS	-	expression tag	UNP Q5ZUA2
B	399	HIS	-	expression tag	UNP Q5ZUA2
B	400	HIS	-	expression tag	UNP Q5ZUA2
B	401	HIS	-	expression tag	UNP Q5ZUA2
B	137	ASP	GLU	conflict	UNP Q5ZUA2
B	149	VAL	ALA	conflict	UNP Q5ZUA2

- Molecule 2 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Na	0	0
			1	1		

- Molecule 3 is MOLYBDATE ION (three-letter code: MOO) (formula: MoO_4).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	Mo	O	0	0
			5	1	4		
3	B	1	Total	Mo	O	0	0
			5	1	4		

- Molecule 4 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	3	Total	Mg	0	0
			3	3		
4	B	2	Total	Mg	0	0
			2	2		

- Molecule 5 is 2-(N-MORPHOLINO)-ETHANESULFONIC ACID (three-letter code: MES) (formula: $\text{C}_6\text{H}_{13}\text{NO}_4\text{S}$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	A	1	Total	C	N	O	S	0	0
			12	6	1	4	1		
5	B	1	Total	C	N	O	S	0	0
			12	6	1	4	1		

- Molecule 6 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	C	O	0	0
			6	3	3		
6	A	1	Total	C	O	0	0
			6	3	3		

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	B	1	Total	C	O	0	0
			6	3	3		
6	B	1	Total	C	O	0	0
			6	3	3		
6	B	1	Total	C	O	0	0
			6	3	3		
6	B	1	Total	C	O	0	0
			6	3	3		
6	B	1	Total	C	O	0	0
			6	3	3		

- Molecule 7 is bis(mu2-oxo)-octa-oxo-dimolybdenum (VI) (three-letter code: M27) (formula: Mo_2O_{10}).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	A	1	Total	Mo	O	0	0
			7	2	5		
7	A	1	Total	Mo	O	0	0
			8	2	6		
7	B	1	Total	Mo	O	0	0
			7	2	5		
7	B	1	Total	Mo	O	0	0
			8	2	6		

- Molecule 8 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: $\text{C}_2\text{H}_6\text{O}_2$).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
8	A	1	Total C O 4 2 2	0	0
8	A	1	Total C O 4 2 2	0	0
8	A	1	Total C O 4 2 2	0	0
8	B	1	Total C O 4 2 2	0	0

- Molecule 9 is TETRAKIS-(MU-OXO)-DECA-OXO-TRIMOLYBDENUM (three-letter code: 6LL) (formula: Mo_3O_{14}).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
9	A	1	Total	Mo	O	0	0
			11	3	8		
9	B	1	Total	Mo	O	0	0
			11	3	8		

- Molecule 10 is bis(mu4-oxo)-bis(mu3-oxo)-octakis(mu2-oxo)-dodecaoxo-heptamolybdenum (VI) (three-letter code: MO7) (formula: Mo_7O_{24}).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
10	A	1	Total	Mo	O	0	0
			31	7	24		
10	B	1	Total	Mo	O	0	0
			31	7	24		

- Molecule 11 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
11	A	1	Total	Cl	0	0
			1	1		
11	B	1	Total	Cl	0	0
			1	1		

- Molecule 12 is bis(mu4-oxo)-tetrakis(mu3-oxo)-hexakis(mu2-oxo)-hexadeca-oxo-octamolybdenum (VI) (three-letter code: 8M0) (formula: Mo_8O_{28}).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
12	A	1	Total	Mo	O	0	0
			34	8	26		
12	B	1	Total	Mo	O	0	0
			34	8	26		


- Molecule 13 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
13	A	240	Total	O	0	0
			240	240		
13	B	211	Total	O	0	1
			212	212		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: ECTONUCLEOSIDE TRIPHOSPHATE DIPHOSPHOHYDROLASE I



- Molecule 1: ECTONUCLEOSIDE TRIPHOSPHATE DIPHOSPHOHYDROLASE I



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	64.56Å 85.44Å 71.78Å 90.00° 103.94° 90.00°	Depositor
Resolution (Å)	45.34 – 1.49 45.34 – 1.49	Depositor EDS
% Data completeness (in resolution range)	96.3 (45.34-1.49) 96.7 (45.34-1.49)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.49 (at 1.48Å)	Xtriage
Refinement program	REFMAC 5.7.0029	Depositor
R, R_{free}	0.126 , 0.169 0.122 , 0.165	Depositor DCC
R_{free} test set	1236 reflections (1.01%)	wwPDB-VP
Wilson B-factor (Å ²)	9.8	Xtriage
Anisotropy	0.935	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 32.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	6758	wwPDB-VP
Average B, all atoms (Å ²)	17.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.25% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: M27, GOL, MOO, MO7, 6LL, 8M0, MG, EDO, CL, NA, MES

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.93	3/3176 (0.1%)	0.84	4/4326 (0.1%)
1	B	0.90	1/3114 (0.0%)	0.83	4/4241 (0.1%)
All	All	0.91	4/6290 (0.1%)	0.84	8/8567 (0.1%)

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	41	LYS	CE-NZ	17.45	1.92	1.49
1	A	96	ASP	CB-CG	5.34	1.62	1.51
1	B	96	ASP	CB-CG	5.30	1.62	1.51
1	A	122	ARG	CZ-NH1	5.12	1.39	1.33

The worst 5 of 8 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	234	MET	CG-SD-CE	9.39	115.22	100.20
1	A	122	ARG	NE-CZ-NH2	-6.51	117.05	120.30
1	B	158	ASP	CB-CG-OD2	-6.19	112.73	118.30
1	A	96	ASP	CB-CG-OD1	6.10	123.79	118.30
1	A	41	LYS	CD-CE-NZ	-6.08	97.71	111.70

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within

the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3034	0	2909	24	1
1	B	2990	0	2856	12	0
2	A	1	0	0	0	0
3	A	5	0	0	1	0
3	B	5	0	0	1	0
4	A	3	0	0	0	0
4	B	2	0	0	0	0
5	A	12	0	13	0	0
5	B	12	0	13	0	0
6	A	12	0	14	1	0
6	B	30	0	35	3	0
7	A	15	0	0	7	0
7	B	15	0	0	6	0
8	A	12	0	12	0	0
8	B	4	0	4	0	0
9	A	11	0	0	3	0
9	B	11	0	0	4	0
10	A	31	0	0	2	0
10	B	31	0	0	6	0
11	A	1	0	0	0	0
11	B	1	0	0	0	0
12	A	34	0	0	9	0
12	B	34	0	0	8	1
13	A	240	0	0	2	0
13	B	212	0	0	4	0
All	All	6758	0	5856	84	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

The worst 5 of 84 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:41:LYS:NZ	1:A:41:LYS:CE	1.92	1.32
1:A:323[A]:GLN:HE21	1:A:327:ASN:HD21	1.15	0.93
1:A:180:LYS:HE2	1:A:293:GLU:HG3	1.58	0.84
1:A:310:HIS:HD2	13:A:2188:HOH:O	1.71	0.74
1:A:323[A]:GLN:HE21	1:A:327:ASN:ND2	1.86	0.72

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the sym-

metry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:398:HIS:NE2	12:B:1403:8M0:MO5[2_755]	2.17	0.03

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	384/368 (104%)	375 (98%)	9 (2%)	0	100	100
1	B	377/368 (102%)	368 (98%)	9 (2%)	0	100	100
All	All	761/736 (103%)	743 (98%)	18 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	346/328 (106%)	346 (100%)	0	100	100
1	B	339/328 (103%)	338 (100%)	1 (0%)	91	82
All	All	685/656 (104%)	684 (100%)	1 (0%)	92	84

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	195	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	207	HIS
1	A	327	ASN
1	B	319	GLN
1	B	370	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 33 ligands modelled in this entry, 8 are monoatomic - leaving 25 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
7	M27	B	1409	6,1	0,6,12	-	-	-		
7	M27	A	1410	1,8	0,7,12	-	-	-		
8	EDO	A	1414	9	3,3,3	0.60	0	2,2,2	0.65	0
7	M27	A	1408	1,8	0,6,12	-	-	-		
7	M27	B	1411	6,1	0,7,12	-	-	-		
12	8M0	A	1417	1	17,46,48	2.02	6 (35%)	-		
9	6LL	A	1412	6,1,8	2,10,18	1.99	1 (50%)	-		
6	GOL	A	1407	-	5,5,5	0.83	0	5,5,5	0.63	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
8	EDO	A	1409	7	3,3,3	0.60	0	2,2,2	0.97	0
8	EDO	B	1415	9	3,3,3	0.59	0	2,2,2	0.91	0
10	MO7	B	1416	-	20,42,42	2.45	8 (40%)	-		
6	GOL	B	1408	-	5,5,5	0.83	0	5,5,5	0.40	0
3	MOO	B	1404	4,13	2,4,4	4.22	1 (50%)	-		
6	GOL	B	1412	7	5,5,5	0.65	0	5,5,5	0.91	0
6	GOL	B	1410	7	5,5,5	0.54	0	5,5,5	0.53	0
6	GOL	B	1417	-	5,5,5	0.31	0	5,5,5	0.57	0
9	6LL	B	1413	6,1,8	2,10,18	1.50	0	-		
12	8M0	B	1403	1	17,46,48	2.12	7 (41%)	-		
8	EDO	A	1411	7	3,3,3	0.75	0	2,2,2	0.96	0
6	GOL	A	1413	9	5,5,5	0.42	0	5,5,5	0.82	0
3	MOO	A	1403	4,13	2,4,4	1.67	1 (50%)	-		
6	GOL	B	1414	9	5,5,5	0.47	0	5,5,5	0.79	0
5	MES	A	1405	-	12,12,12	2.48	1 (8%)	14,16,16	1.84	4 (28%)
10	MO7	A	1415	-	20,42,42	2.93	8 (40%)	-		
5	MES	B	1406	-	12,12,12	2.09	2 (16%)	14,16,16	1.45	3 (21%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	GOL	A	1407	-	-	0/4/4/4	-
6	GOL	A	1413	9	-	1/4/4/4	-
8	EDO	A	1414	9	-	1/1/1/1	-
8	EDO	A	1409	7	-	1/1/1/1	-
6	GOL	B	1412	7	-	0/4/4/4	-
6	GOL	B	1410	7	-	2/4/4/4	-
6	GOL	B	1414	9	-	0/4/4/4	-
8	EDO	B	1415	9	-	1/1/1/1	-
6	GOL	B	1417	-	-	2/4/4/4	-
5	MES	A	1405	-	-	3/6/14/14	0/1/1/1
6	GOL	B	1408	-	-	0/4/4/4	-
8	EDO	A	1411	7	-	0/1/1/1	-
5	MES	B	1406	-	-	3/6/14/14	0/1/1/1

The worst 5 of 35 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	A	1405	MES	C8-S	-8.18	1.65	1.77
5	B	1406	MES	C8-S	-6.53	1.68	1.77
3	B	1404	MOO	O1-MO	5.96	1.86	1.73
10	A	1415	MO7	O4-MO1	5.76	1.80	1.74
10	A	1415	MO7	O15-MO5	5.52	1.78	1.68

The worst 5 of 7 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	1405	MES	C6-C5-N4	-2.96	105.61	110.10
5	A	1405	MES	O1-C2-C3	-2.91	105.40	111.80
5	B	1406	MES	C5-N4-C3	2.84	115.21	108.83
5	A	1405	MES	C6-O1-C2	2.77	119.15	109.89
5	B	1406	MES	C6-O1-C2	2.58	118.52	109.89

There are no chirality outliers.

5 of 14 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	B	1406	MES	C8-C7-N4-C5
6	B	1410	GOL	C1-C2-C3-O3
6	B	1417	GOL	C1-C2-C3-O3
5	B	1406	MES	C8-C7-N4-C3
6	B	1417	GOL	O2-C2-C3-O3

There are no ring outliers.

14 monomers are involved in 52 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
7	B	1409	M27	3	0
7	A	1410	M27	3	0
7	A	1408	M27	4	0
7	B	1411	M27	3	0
12	A	1417	8M0	9	0
9	A	1412	6LL	3	0
10	B	1416	MO7	6	0
3	B	1404	MOO	1	0
6	B	1417	GOL	3	0
9	B	1413	6LL	4	0
12	B	1403	8M0	8	1
6	A	1413	GOL	1	0
3	A	1403	MOO	1	0


Continued on next page...


Continued from previous page...

Mol	Chain	Res	Type	Clashes	Symm-Clashes
10	A	1415	MO7	2	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









Ligand 8M0 B 1403




Bond lengths




Bond angles



Torsions



Rings



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	365/368 (99%)	-0.51	4 (1%) 77 80	6, 13, 32, 63	21 (5%)
1	B	365/368 (99%)	-0.47	7 (1%) 66 69	5, 13, 32, 65	14 (3%)
All	All	730/736 (99%)	-0.49	11 (1%) 71 74	5, 13, 32, 65	35 (4%)

The worst 5 of 11 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	177	VAL	6.5
1	A	177	VAL	4.4
1	B	179	ASN	3.0
1	A	41	LYS	2.7
1	B	175	LYS	2.6

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
6	GOL	B	1417	6/6	0.93	0.10	22,35,38,51	0
8	EDO	B	1415	4/4	0.93	0.10	19,20,24,26	0
8	EDO	A	1414	4/4	0.95	0.08	20,20,25,26	0
5	MES	B	1406	12/12	0.95	0.12	21,30,35,38	12
5	MES	A	1405	12/12	0.96	0.11	19,28,33,39	12
8	EDO	A	1411	4/4	0.96	0.07	17,18,19,19	0
6	GOL	A	1413	6/6	0.96	0.09	20,20,22,22	0
6	GOL	B	1414	6/6	0.96	0.06	20,20,21,24	0
6	GOL	A	1407	6/6	0.97	0.08	13,14,16,18	0
8	EDO	A	1409	4/4	0.98	0.06	13,16,17,19	0
6	GOL	B	1412	6/6	0.98	0.06	16,24,28,29	0
4	MG	A	1418	1/1	0.98	0.14	26,26,26,26	0
6	GOL	B	1410	6/6	0.98	0.07	14,17,28,34	0
6	GOL	B	1408	6/6	0.99	0.06	13,15,16,16	0
2	NA	A	1402	1/1	0.99	0.18	32,32,32,32	0
4	MG	B	1407	1/1	0.99	0.14	20,20,20,20	1
4	MG	B	1405	1/1	1.00	0.01	9,9,9,9	0
7	M27	A	1408	7/12	1.00	0.02	10,11,15,17	0
7	M27	A	1410	8/12	1.00	0.02	15,16,18,19	0
7	M27	B	1409	7/12	1.00	0.02	10,12,15,18	0
7	M27	B	1411	8/12	1.00	0.04	17,19,25,26	8
3	MOO	B	1404	5/5	1.00	0.02	7,7,9,9	0
4	MG	A	1404	1/1	1.00	0.02	10,10,10,10	0
4	MG	A	1406	1/1	1.00	0.09	18,18,18,18	1
3	MOO	A	1403	5/5	1.00	0.02	8,8,8,9	0
9	6LL	A	1412	11/17	1.00	0.03	12,15,18,21	11
9	6LL	B	1413	11/17	1.00	0.03	13,15,19,21	11
10	MO7	A	1415	31/31	1.00	0.03	14,18,25,26	0
10	MO7	B	1416	31/31	1.00	0.03	14,17,27,29	0
11	CL	A	1416	1/1	1.00	0.02	12,12,12,12	0
11	CL	B	1402	1/1	1.00	0.01	13,13,13,13	0
12	8M0	A	1417	34/36	1.00	0.03	10,12,16,17	0
12	8M0	B	1403	34/36	1.00	0.03	10,12,17,17	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around 6LL A 1412:

$2mF_o - DF_c$ (at 0.7 rmsd) in gray
 $mF_o - DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around MO7 A 1415:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around MO7 B 1416:


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around 8M0 A 1417:

2mF_o-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers ⓘ

There are no such residues in this entry.