



wwPDB X-ray Structure Validation Summary Report ⓘ

Oct 25, 2023 – 12:19 AM EDT

PDB ID : 3DBJ
Title : Allophycocyanin from *Thermosynechococcus vulcanus*
Authors : Adir, N.; Klartag, M.; McGregor, A.; David, L.
Deposited on : 2008-06-01
Resolution : 2.90 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1957 (2.90-2.90)
Clashscore	141614	2172 (2.90-2.90)
Ramachandran outliers	138981	2115 (2.90-2.90)
Sidechain outliers	138945	2117 (2.90-2.90)
RSRZ outliers	127900	1906 (2.90-2.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	161	
1	C	161	
1	E	161	
1	G	161	
2	B	161	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain			
2	D	161		48%	44%	7%
2	F	161		48%	44%	8%
2	H	161		43%	48%	9%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	MEN	B	71	-	-	X	-
2	MEN	D	71	-	-	X	-
2	MEN	F	71	-	-	X	-
2	MEN	H	71	-	-	X	-

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 10448 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Allophycocyanin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	160	1220	762	212	240	6	0	0	0
1	C	160	1220	762	212	240	6	0	0	0
1	E	160	1220	762	212	240	6	0	0	0
1	G	160	1220	762	212	240	6	0	0	0

- Molecule 2 is a protein called Allophycocyanin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	161	1216	768	203	237	8	0	0	0
2	D	161	1216	768	203	237	8	0	0	0
2	F	161	1216	768	203	237	8	0	0	0
2	H	161	1216	768	203	237	8	0	0	0

- Molecule 3 is PHYCOCYANOBILIN (three-letter code: CYC) (formula: $C_{33}H_{40}N_4O_6$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			43	33	4	6		
3	B	1	Total	C	N	O	0	0
			43	33	4	6		
3	C	1	Total	C	N	O	0	0
			43	33	4	6		
3	D	1	Total	C	N	O	0	0
			43	33	4	6		
3	E	1	Total	C	N	O	0	0
			43	33	4	6		
3	F	1	Total	C	N	O	0	0
			43	33	4	6		
3	G	1	Total	C	N	O	0	0
			43	33	4	6		
3	H	1	Total	C	N	O	0	0
			43	33	4	6		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	38	Total	O	0	0
			38	38		
4	B	48	Total	O	0	0
			48	48		
4	C	51	Total	O	0	0
			51	51		
4	D	45	Total	O	0	0
			45	45		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	E	45	Total 45	O 45	0	0
4	F	45	Total 45	O 45	0	0
4	G	49	Total 49	O 49	0	0
4	H	39	Total 39	O 39	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Allophycocyanin



- Molecule 1: Allophycocyanin



- Molecule 1: Allophycocyanin



M173
Q174

- Molecule 1: Allophycocyanin

Chain G:  44% 51%

ME1 S3 V4 V5 V6 V7 V8 V9 V10 V11 V12 V13 V14 V15 V16 V17 V18 V19 V20 V21 V22 V23 V24 V25 V26 V27 V28 V29 V30 V31 V32 V33 V34 V35 V36 V37 V38 V39 V40 V41 V42 V43 V44 V45 V46 V47 V48 V49 V50 V51 V52 V53 V54 V55 V56 V57 V58 V59 V60 V61 V62 V63 V64 V65 V66 V67 V68 V69 V70 V71 V72 V73 V74

Y76 G77 E78 E79 E80 E81 E82 E83 E84 E85 E86 E87 E88 E89 E90 E91 E92 E93 E94 E95 E96 E97 E98 E99 V100 D103 D104 D105 D106 D107 D108 D109 D110 D111 D112 D113 D114 D115 D116 Y119 T124 T125 T126 T127 T128 E131 I132 I133 I134 I135 I136 I137 I138 I139 L143 L144 E147

D148 A149 F166 F167 F168 F169 F170 M173 Q174

- Molecule 2: Allophycocyanin

Chain B:  44% 47% 9%

M1 Q2 D3 D4 I5 T6 T7 T8 T9 T10 T11 T12 T13 T14 T15 T16 T17 T18 T19 T20 T21 T22 M24 L27 K28 K29 K30 F31 A32 A33 A34 A35 A36 A37 A38 A39 A40 I44 N47 I51 I52 I53 I54 I55 I56 L60 L61 L62 L63 L64 L65 L66 L67 L68 L69 L70 N71 N72 N73 N74 N75 N76 N77

T78 R79 R80 R81 R82 R83 R84 R85 R86 R87 Y90 Y91 Y92 Y93 Y94 Y95 Y96 Y97 Y98 Y99 G102 G103 G104 G105 G106 G107 V111 V112 V113 V114 V115 V116 V117 T118 S121 S122 S123 L122 L123 L124 L125 L126 L127 P125 P126 P127 A128 A129 A130 A131 A132 A133 A134 A135 A136 A137 T140 T141 T142 T143 T144 D147 D148 M162

Y165 F166 C170 G172 G173 S174

- Molecule 2: Allophycocyanin

Chain D:  48% 44% 7%

M1 Q2 D3 D4 I5 T6 T7 T8 T9 T10 T11 T12 T13 T14 T15 T16 T17 T18 T19 T20 T21 T22 M24 L27 K28 K29 K30 F31 A32 A33 A34 A35 A36 A37 A38 A39 A40 I44 N47 I51 I52 I53 I54 I55 I56 L60 L61 L62 L63 L64 L65 L66 L67 L68 L69 L70 N71 N72 N73 N74 N75 N76 N77


T78 R79 R80 R81 R82 R83 R84 R85 R86 R87 Y90 Y91 Y92 Y93 Y94 Y95 Y96 Y97 Y98 Y99 D103 D104 D105 D106 D107 V111 V112 V113 V114 V115 V116 V117 S121 S122 S123 P125 P126 P127 A128 A129 A130 A131 A132 A133 A134 A135 A136 A137 E138 V139 T140 T141 T142 T143 T144 D147 Y165 F166 C170 G172

L173 S174

- Molecule 2: Allophycocyanin

Chain F:  48% 44% 8%

M1 Q2 D3 D4 I5 T6 T7 T8 T9 T10 T11 T12 T13 T14 T15 T16 T17 T18 T19 T20 T21 T22 M24 L27 K28 K29 K30 F31 A32 A33 A34 A35 A36 A37 A38 A39 A40 I44 N47 I51 I52 I53 I54 I55 I56 L60 L61 L62 L63 L64 L65 L66 L67 L68 L69 L70 N71 N72 N73 N74 N75 N76 N77 R79



L173
S174

● Molecule 2: Allophycocyanin

Chain H: 43% 48% 9%



M162
Y165
F166
C170
S171
G172
L173
S174

4 Data and refinement statistics

Property	Value	Source
Space group	P 3	Depositor
Cell constants a, b, c, α , β , γ	102.60Å 102.60Å 128.86Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	20.00 – 2.90 20.13 – 2.90	Depositor EDS
% Data completeness (in resolution range)	8.7 (20.00-2.90) 91.7 (20.13-2.90)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.45 (at 2.88Å)	Xtrriage
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.212 , 0.270 0.221 , 0.271	Depositor DCC
R_{free} test set	3097 reflections (10.09%)	wwPDB-VP
Wilson B-factor (Å ²)	32.9	Xtrriage
Anisotropy	0.508	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 31.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.29$	Xtrriage
Estimated twinning fraction	0.457 for -h,-k,l 0.458 for h,-h-k,-l 0.459 for -k,-h,-l	Xtrriage
F_o, F_c correlation	0.90	EDS
Total number of atoms	10448	wwPDB-VP
Average B, all atoms (Å ²)	31.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.45% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CYC, MEN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.37	0/1236	0.60	0/1673
1	C	0.37	0/1236	0.59	0/1673
1	E	0.38	0/1236	0.59	0/1673
1	G	0.37	0/1236	0.59	0/1673
2	B	0.37	0/1222	0.64	0/1653
2	D	0.36	0/1222	0.65	0/1653
2	F	0.36	0/1222	0.64	0/1653
2	H	0.38	0/1222	0.64	0/1653
All	All	0.37	0/9832	0.62	0/13304

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1220	0	1219	103	0
1	C	1220	0	1219	100	0
1	E	1220	0	1219	92	0
1	G	1220	0	1219	102	0
2	B	1216	0	1227	119	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	D	1216	0	1228	99	0
2	F	1216	0	1228	105	0
2	H	1216	0	1228	111	0
3	A	43	0	37	5	0
3	B	43	0	37	4	0
3	C	43	0	37	5	0
3	D	43	0	37	4	0
3	E	43	0	37	5	0
3	F	43	0	37	4	0
3	G	43	0	37	5	0
3	H	43	0	38	6	0
4	A	38	0	0	4	0
4	B	48	0	0	17	0
4	C	51	0	0	6	0
4	D	45	0	0	9	0
4	E	45	0	0	7	0
4	F	45	0	0	9	0
4	G	49	0	0	10	0
4	H	39	0	0	13	0
All	All	10448	0	10084	778	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 39.

The worst 5 of 778 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:H:84:CYS:SG	3:H:202:CYC:HAC2	1.26	1.75
2:B:75:MET:HG3	4:B:210:HOH:O	1.31	1.24
1:E:68:SER:HB3	1:E:69:PRO:HD2	1.48	0.95
3:E:201:CYC:HMA1	3:E:201:CYC:HB	1.33	0.94
2:H:77:THR:HG22	2:H:79:ARG:H	1.32	0.93

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	158/161 (98%)	144 (91%)	14 (9%)	0	100	100
1	C	158/161 (98%)	143 (90%)	14 (9%)	1 (1%)	25	58
1	E	158/161 (98%)	142 (90%)	15 (10%)	1 (1%)	25	58
1	G	158/161 (98%)	143 (90%)	14 (9%)	1 (1%)	25	58
2	B	158/161 (98%)	136 (86%)	20 (13%)	2 (1%)	12	37
2	D	158/161 (98%)	137 (87%)	19 (12%)	2 (1%)	12	37
2	F	158/161 (98%)	137 (87%)	19 (12%)	2 (1%)	12	37
2	H	158/161 (98%)	137 (87%)	19 (12%)	2 (1%)	12	37
All	All	1264/1288 (98%)	1119 (88%)	134 (11%)	11 (1%)	17	48

5 of 11 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	77	THR
2	D	77	THR
2	F	77	THR
2	H	77	THR
2	D	111	VAL

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	129/130 (99%)	117 (91%)	12 (9%)	9	27

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	C	129/130 (99%)	117 (91%)	12 (9%)	9	27
1	E	129/130 (99%)	117 (91%)	12 (9%)	9	27
1	G	129/130 (99%)	117 (91%)	12 (9%)	9	27
2	B	123/123 (100%)	106 (86%)	17 (14%)	3	10
2	D	123/123 (100%)	107 (87%)	16 (13%)	4	12
2	F	123/123 (100%)	107 (87%)	16 (13%)	4	12
2	H	123/123 (100%)	106 (86%)	17 (14%)	3	10
All	All	1008/1012 (100%)	894 (89%)	114 (11%)	6	18

5 of 114 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	E	25	ASP
2	H	143	LEU
2	F	36	LEU
2	H	134	GLN
2	H	36	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 17 such sidechains are listed below:

Mol	Chain	Res	Type
1	G	120	ASN
2	H	47	ASN
1	E	47	ASN
1	E	54	GLN
1	E	120	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

4 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The

Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	MEN	B	71	2	7,8,9	2.79	1 (14%)	6,9,11	1.86	2 (33%)
2	MEN	D	71	2	7,8,9	0.88	1 (14%)	6,9,11	2.39	1 (16%)
2	MEN	H	71	2	7,8,9	1.29	1 (14%)	6,9,11	2.73	2 (33%)
2	MEN	F	71	2	7,8,9	0.67	0	6,9,11	2.39	1 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	MEN	B	71	2	-	6/7/8/10	-
2	MEN	D	71	2	-	4/7/8/10	-
2	MEN	H	71	2	-	2/7/8/10	-
2	MEN	F	71	2	-	2/7/8/10	-

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	71	MEN	CE2-ND2	7.30	1.58	1.45
2	H	71	MEN	CE2-ND2	3.21	1.51	1.45
2	D	71	MEN	CE2-ND2	2.01	1.49	1.45

The worst 5 of 6 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	H	71	MEN	CB-CA-C	-5.81	100.57	111.47
2	F	71	MEN	CB-CA-C	-5.79	100.62	111.47
2	D	71	MEN	CB-CA-C	-5.79	100.62	111.47
2	B	71	MEN	CE2-ND2-CG	3.81	151.01	121.93
2	H	71	MEN	CE2-ND2-CG	3.21	146.41	121.93

There are no chirality outliers.

5 of 14 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	71	MEN	CB-CG-ND2-CE2
2	D	71	MEN	CB-CG-ND2-CE2
2	B	71	MEN	OD1-CG-ND2-CE2
2	D	71	MEN	OD1-CG-ND2-CE2
2	B	71	MEN	CA-CB-CG-OD1

There are no ring outliers.

4 monomers are involved in 29 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	71	MEN	7	0
2	D	71	MEN	6	0
2	H	71	MEN	6	0
2	F	71	MEN	10	0

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

8 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	CYC	H	202	2	42,46,46	2.30	14 (33%)	50,67,67	3.54	19 (38%)
3	CYC	E	201	1	42,46,46	2.37	16 (38%)	50,67,67	3.53	24 (48%)
3	CYC	C	201	1	42,46,46	2.37	15 (35%)	50,67,67	3.55	24 (48%)
3	CYC	D	202	2	42,46,46	2.28	14 (33%)	50,67,67	3.53	19 (38%)
3	CYC	A	201	1	42,46,46	2.39	14 (33%)	50,67,67	3.54	24 (48%)
3	CYC	B	202	2	42,46,46	2.28	14 (33%)	50,67,67	3.56	19 (38%)
3	CYC	G	201	1	42,46,46	2.38	15 (35%)	50,67,67	3.54	24 (48%)
3	CYC	F	202	2	42,46,46	2.30	14 (33%)	50,67,67	3.55	19 (38%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	CYC	H	202	2	-	10/25/74/74	0/4/4/4
3	CYC	E	201	1	-	8/25/74/74	0/4/4/4
3	CYC	C	201	1	-	8/25/74/74	0/4/4/4
3	CYC	D	202	2	-	10/25/74/74	0/4/4/4
3	CYC	A	201	1	-	8/25/74/74	0/4/4/4
3	CYC	B	202	2	-	10/25/74/74	0/4/4/4
3	CYC	G	201	1	-	8/25/74/74	0/4/4/4
3	CYC	F	202	2	-	10/25/74/74	0/4/4/4

The worst 5 of 116 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	H	202	CYC	OB-C4B	7.34	1.37	1.23
3	F	202	CYC	OB-C4B	7.16	1.37	1.23
3	D	202	CYC	OB-C4B	7.08	1.37	1.23
3	B	202	CYC	OB-C4B	7.07	1.37	1.23
3	G	201	CYC	OB-C4B	6.95	1.36	1.23

The worst 5 of 172 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	202	CYC	C3B-C4B-NB	12.80	117.12	106.78
3	E	201	CYC	C3B-C4B-NB	12.68	117.03	106.78
3	A	201	CYC	C3B-C4B-NB	12.66	117.01	106.78
3	C	201	CYC	C3B-C4B-NB	12.62	116.98	106.78
3	H	202	CYC	C3B-C4B-NB	12.62	116.97	106.78

There are no chirality outliers.

5 of 72 torsion outliers are listed below:



Mol	Chain	Res	Type	Atoms
3	A	201	CYC	ND-C1D-CHD-C4C
3	A	201	CYC	C2D-C1D-CHD-C4C
3	B	202	CYC	NA-C4A-CHB-C1B
3	B	202	CYC	C3A-C4A-CHB-C1B
3	B	202	CYC	ND-C1D-CHD-C4C



There are no ring outliers.



8 monomers are involved in 38 short contacts:



Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	H	202	CYC	6	0
3	E	201	CYC	5	0
3	C	201	CYC	5	0
3	D	202	CYC	4	0
3	A	201	CYC	5	0
3	B	202	CYC	4	0
3	G	201	CYC	5	0
3	F	202	CYC	4	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	160/161 (99%)	-0.40	1 (0%) 89 89	9, 34, 60, 78	0
1	C	160/161 (99%)	-0.39	0 100 100	5, 33, 59, 75	1 (0%)
1	E	160/161 (99%)	-0.38	1 (0%) 89 89	3, 33, 61, 78	0
1	G	160/161 (99%)	-0.39	1 (0%) 89 89	11, 33, 60, 77	0
2	B	160/161 (99%)	-0.52	0 100 100	8, 27, 54, 68	0
2	D	160/161 (99%)	-0.52	0 100 100	9, 27, 52, 66	0
2	F	160/161 (99%)	-0.50	0 100 100	10, 28, 51, 67	0
2	H	160/161 (99%)	-0.52	0 100 100	5, 28, 50, 65	0
All	All	1280/1288 (99%)	-0.45	3 (0%) 95 95	3, 30, 55, 78	1 (0%)

All (3) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	E	30	PHE	2.3
1	A	30	PHE	2.3
1	G	30	PHE	2.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	MEN	B	71	9/10	0.90	0.22	46,49,57,58	0
2	MEN	H	71	9/10	0.91	0.22	41,50,53,54	0
2	MEN	F	71	9/10	0.92	0.24	44,54,58,60	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	MEN	D	71	9/10	0.92	0.23	38,41,55,55	0

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	CYC	H	202	43/43	0.94	0.18	15,21,30,31	0
3	CYC	F	202	43/43	0.95	0.19	22,29,35,36	0
3	CYC	B	202	43/43	0.95	0.17	15,23,28,28	0
3	CYC	C	201	43/43	0.96	0.17	20,27,32,34	0
3	CYC	D	202	43/43	0.96	0.18	20,26,34,35	0
3	CYC	A	201	43/43	0.97	0.16	18,23,28,29	0
3	CYC	E	201	43/43	0.97	0.16	20,26,31,32	0
3	CYC	G	201	43/43	0.98	0.13	1,8,13,15	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around CYC H 202:


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




Electron density around CYC F 202:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)













Electron density around CYC D 202:


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




Electron density around CYC A 201:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







6.5 Other polymers [i](#)

There are no such residues in this entry.