



# Full wwPDB X-ray Structure Validation Report ⓘ

Jun 12, 2024 – 09:35 PM EDT

PDB ID : 3DDV  
Title : The crystal structure of the transcriptional regulator (GntR family) from *Enterococcus faecalis* V583  
Authors : Zhang, R.; Zhou, M.; Bargassa, M.; Otwinowski, Z.; Joachimiak, A.; Midwest Center for Structural Genomics (MCSG)  
Deposited on : 2008-06-06  
Resolution : 2.65 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.20.1
EDS	:	2.36.2
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.36.2


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.65 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1332 (2.68-2.64)
Clashscore	141614	1374 (2.68-2.64)
Ramachandran outliers	138981	1349 (2.68-2.64)
Sidechain outliers	138945	1349 (2.68-2.64)
RSRZ outliers	127900	1318 (2.68-2.64)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	145	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 1%, green 54%, yellow 25%, orange 1%, grey 17%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> <span>%</span> <span>54%</span> <span>25%</span> <span>• •</span> <span>17%</span> </div> </div>
1	B	145	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 1%, green 70%, yellow 23%, orange 1%, grey 5%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> <span>%</span> <span>70%</span> <span>23%</span> <span>• •</span> </div> </div>
1	C	145	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 1%, green 59%, yellow 18%, orange 1%, grey 17%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> <span>%</span> <span>59%</span> <span>18%</span> <span>• •</span> <span>17%</span> </div> </div>
1	D	145	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 1%, green 81%, yellow 13%, orange 1%, grey 5%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> <span>%</span> <span>81%</span> <span>13%</span> <span>•</span> <span>5%</span> </div> </div>

## 2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 4252 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Transcriptional regulator (GntR family).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	121	Total	C	N	O	S	0	0	0
			980	624	167	186	3			
1	B	139	Total	C	N	O	S	0	0	0
			1121	713	188	217	3			
1	C	120	Total	C	N	O	S	0	0	0
			972	617	166	186	3			
1	D	138	Total	C	N	O	S	0	0	0
			1112	707	186	216	3			

- Molecule 2 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Mg	0	0
			1	1		
2	B	1	Total	Mg	0	0
			1	1		
2	D	1	Total	Mg	0	0
			1	1		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	18	Total	O	0	0
			18	18		
3	B	16	Total	O	0	0
			16	16		
3	C	12	Total	O	0	0
			12	12		
3	D	18	Total	O	0	0
			18	18		


### 3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Transcriptional regulator (GntR family)




- Molecule 1: Transcriptional regulator (GntR family)




- Molecule 1: Transcriptional regulator (GntR family)



- Molecule 1: Transcriptional regulator (GntR family)





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	86.15Å 86.15Å 240.06Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	80.06 – 2.65 46.76 – 2.65	Depositor EDS
% Data completeness (in resolution range)	98.1 (80.06-2.65) 98.1 (46.76-2.65)	Depositor EDS
$R_{merge}$	0.12	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	4.28 (at 2.65Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, $R_{free}$	0.206 , 0.251 0.225 , 0.265	Depositor DCC
$R_{free}$ test set	1529 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	72.9	Xtriage
Anisotropy	0.193	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 57.6	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.027 for -h,-k,l	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	4252	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	59.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.14% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	1.11	1/999 (0.1%)	1.03	0/1346
1	B	1.01	1/1144 (0.1%)	0.98	2/1540 (0.1%)
1	C	1.06	0/991	0.99	2/1336 (0.1%)
1	D	0.87	1/1135 (0.1%)	0.83	1/1529 (0.1%)
All	All	1.01	3/4269 (0.1%)	0.96	5/5751 (0.1%)

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	108	ARG	CG-CD	6.28	1.67	1.51
1	A	134	TYR	CB-CG	-5.31	1.43	1.51
1	D	51	GLU	CD-OE2	5.29	1.31	1.25

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	131	ARG	NE-CZ-NH1	6.28	123.44	120.30
1	B	116	ARG	NE-CZ-NH2	-5.64	117.48	120.30
1	D	114	ARG	NE-CZ-NH2	-5.62	117.49	120.30
1	C	116	ARG	CG-CD-NE	5.51	123.38	111.80
1	B	41	ARG	NE-CZ-NH1	5.08	122.84	120.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	980	0	970	47	0
1	B	1121	0	1102	29	0
1	C	972	0	954	37	0
1	D	1112	0	1089	14	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	D	1	0	0	0	0
3	A	18	0	0	0	0
3	B	16	0	0	3	0
3	C	12	0	0	1	0
3	D	18	0	0	3	0
All	All	4252	0	4115	113	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 14.

All (113) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:84:ILE:HG22	1:A:85:GLY:H	1.06	1.11
1:A:59:LEU:O	1:A:60:VAL:HG23	1.53	1.06
1:A:59:LEU:O	1:A:60:VAL:CG2	2.04	1.03
1:A:137:SER:O	1:A:138:ARG:HB3	1.50	1.03
1:C:136:GLY:HA2	1:C:137:SER:HB2	1.46	0.95
1:A:90:THR:CG2	1:B:140:GLU:OE1	2.16	0.93
1:C:137:SER:O	1:C:138:ARG:HB2	1.67	0.92
1:A:84:ILE:HG22	1:A:85:GLY:N	1.87	0.88
1:C:49:CYS:SG	3:C:209:HOH:O	2.37	0.82
1:A:84:ILE:CG2	1:A:85:GLY:H	1.89	0.81
1:B:100:ILE:HD13	1:B:133:GLN:NE2	1.95	0.81
1:C:92:SER:HB2	3:D:191:HOH:O	1.80	0.81
1:B:112:ILE:HG21	1:B:133:GLN:CG	2.11	0.80
1:A:137:SER:O	1:A:138:ARG:CB	2.30	0.79
1:A:136:GLY:HA2	1:A:137:SER:CB	2.16	0.75
1:C:137:SER:O	1:C:138:ARG:CB	2.38	0.72
1:C:2:GLN:HE21	1:C:2:GLN:N	1.88	0.72
1:D:49:CYS:SG	3:D:150:HOH:O	2.48	0.71
1:A:59:LEU:O	1:A:60:VAL:HG22	1.91	0.70
1:C:92:SER:CB	3:D:191:HOH:O	2.39	0.70

*Continued on next page...*



*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:83:LYS:N	1:A:84:ILE:HD12	2.07	0.70
1:B:112:ILE:HG21	1:B:133:GLN:HG3	1.73	0.70
1:C:136:GLY:HA2	1:C:137:SER:CB	2.19	0.70
1:B:112:ILE:HG21	1:B:133:GLN:HG2	1.72	0.69
1:A:90:THR:HG21	1:B:140:GLU:OE1	1.90	0.68
1:A:136:GLY:HA2	1:A:137:SER:HB2	1.76	0.67
1:C:13:TYR:HE2	1:C:35:LEU:HD11	1.61	0.65
1:A:18:PRO:HD3	1:A:32:ASP:O	1.99	0.63
1:C:136:GLY:CA	1:C:137:SER:HB2	2.26	0.63
1:A:94:VAL:CG1	1:A:95:GLN:N	2.61	0.62
1:C:99:GLN:NE2	1:C:103:TYR:CE1	2.67	0.62
1:C:13:TYR:CE2	1:C:35:LEU:HD11	2.35	0.62
1:A:17:LYS:NZ	1:D:41:ARG:HE	1.98	0.61
1:B:47:PRO:O	1:B:138:ARG:NH1	2.33	0.60
1:C:99:GLN:NE2	1:C:103:TYR:OH	2.32	0.60
1:A:91:ILE:HD11	1:B:143:LEU:HD11	1.83	0.59
1:A:135:ALA:O	1:A:139:PHE:CD2	2.57	0.58
1:D:14:PHE:O	1:D:35:LEU:HD12	2.04	0.58
1:B:18:PRO:HD3	1:B:32:ASP:O	2.04	0.58
1:C:123:ASN:N	1:C:123:ASN:OD1	2.37	0.56
1:A:90:THR:HG23	1:B:140:GLU:OE1	2.03	0.56
1:C:129:TYR:CZ	1:C:131:ARG:HD3	2.41	0.56
1:D:72:PHE:CZ	1:D:76:LEU:HD11	2.42	0.55
1:A:84:ILE:C	1:A:85:GLY:O	2.45	0.54
1:A:95:GLN:HE21	1:A:109:GLY:HA2	1.73	0.54
1:B:114:ARG:NH1	3:B:190:HOH:O	2.34	0.54
1:A:136:GLY:CA	1:A:137:SER:CB	2.85	0.54
1:C:57:TYR:O	1:C:60:VAL:HG22	2.08	0.53
1:C:136:GLY:CA	1:C:137:SER:CB	2.86	0.53
1:C:95:GLN:HG2	1:C:109:GLY:HA2	1.90	0.53
1:A:84:ILE:HG23	1:A:127:PHE:CE2	2.44	0.53
1:A:135:ALA:O	1:A:139:PHE:HD2	1.92	0.53
1:D:101:ALA:HB1	1:D:106:ILE:O	2.09	0.52
1:A:136:GLY:HA3	1:A:137:SER:O	2.09	0.52
1:B:76:LEU:O	1:B:80:SER:OG	2.19	0.51
1:C:131:ARG:CG	1:C:131:ARG:HH11	2.23	0.51
1:B:17:LYS:HD2	1:B:31:GLU:HA	1.93	0.51
1:B:98:GLU:HB2	1:B:108:ARG:HH11	1.76	0.50
1:B:23:MET:HE1	1:B:30:PRO:HD3	1.93	0.49
1:A:34:ILE:HG22	1:A:56:PRO:HA	1.95	0.49
1:B:94:VAL:HG21	1:B:114:ARG:NH1	2.27	0.49

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:84:ILE:O	1:A:85:GLY:O	2.30	0.49
1:C:112:ILE:HG12	1:C:134:TYR:O	2.13	0.48
1:A:84:ILE:HD12	1:A:84:ILE:N	2.28	0.48
1:C:131:ARG:HG3	1:C:131:ARG:NH1	2.29	0.48
1:C:44:ASP:O	1:C:45:ASP:HB2	2.14	0.48
1:C:112:ILE:HD13	1:C:112:ILE:HG21	1.47	0.48
1:C:90:THR:HG22	1:D:142:TYR:CD1	2.48	0.48
1:A:138:ARG:HG2	1:A:139:PHE:N	2.28	0.48
1:A:98:GLU:HB2	1:A:108:ARG:HE	1.78	0.47
1:D:49:CYS:HB3	1:D:134:TYR:CD1	2.49	0.47
1:C:131:ARG:HH11	1:C:131:ARG:HG3	1.79	0.47
1:A:94:VAL:HG12	1:A:95:GLN:N	2.30	0.47
1:B:128:GLU:HG2	1:B:130:VAL:CG2	2.45	0.47
1:C:94:VAL:CG2	1:C:112:ILE:HG22	2.44	0.47
1:B:30:PRO:O	1:B:31:GLU:HB2	2.15	0.47
1:B:100:ILE:HD13	1:B:133:GLN:CD	2.34	0.47
1:A:84:ILE:CG2	1:A:85:GLY:N	2.59	0.46
1:A:17:LYS:HZ3	1:D:41:ARG:HE	1.62	0.46
1:A:84:ILE:HD12	1:A:84:ILE:H	1.81	0.46
1:B:30:PRO:HA	3:B:154:HOH:O	2.15	0.46
1:A:100:ILE:HD13	1:A:133:GLN:CD	2.35	0.46
1:C:122:GLU:C	1:C:124:GLY:H	2.19	0.46
1:A:88:ASN:CB	1:B:145:LYS:HZ3	2.29	0.46
1:B:37:MET:CE	1:B:69:THR:HG22	2.46	0.46
1:C:91:ILE:CD1	1:D:143:LEU:HD11	2.46	0.46
1:B:95:GLN:HG3	3:B:187:HOH:O	2.17	0.45
1:A:17:LYS:HZ1	1:D:41:ARG:HE	1.65	0.45
1:C:122:GLU:C	1:C:124:GLY:N	2.70	0.45
1:D:18:PRO:HD3	1:D:32:ASP:O	2.16	0.44
1:A:101:ALA:HB1	1:A:106:ILE:O	2.17	0.44
1:B:30:PRO:O	1:B:31:GLU:CB	2.66	0.44
1:C:34:ILE:O	1:C:34:ILE:HD12	2.17	0.44
1:A:23:MET:HE3	1:A:29:GLY:HA2	1.99	0.44
1:A:94:VAL:HG13	1:A:95:GLN:N	2.31	0.44
1:A:135:ALA:C	1:A:136:GLY:O	2.54	0.44
1:B:65:LYS:O	1:B:69:THR:OG1	2.29	0.43
1:C:94:VAL:HG22	1:C:112:ILE:HG22	2.00	0.43
1:A:136:GLY:HA2	1:A:137:SER:HB3	1.98	0.43
1:A:135:ALA:O	1:A:139:PHE:CE2	2.72	0.43
1:C:2:GLN:HG3	1:C:4:ARG:HH21	1.85	0.42
1:D:129:TYR:CZ	1:D:131:ARG:HD3	2.54	0.42

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:91:ILE:CD1	1:B:143:LEU:HD11	2.50	0.42
1:B:142:TYR:CD1	1:C:27:GLN:HG2	2.55	0.42
1:A:88:ASN:CB	1:B:145:LYS:NZ	2.83	0.42
1:C:4:ARG:HG2	1:C:44:ASP:OD2	2.19	0.41
1:C:91:ILE:HD11	1:D:143:LEU:HD11	2.02	0.41
1:A:125:LEU:HA	1:A:126:PRO:HD3	1.91	0.41
1:B:43:ALA:HB3	1:B:48:ILE:HD11	2.02	0.41
1:C:122:GLU:O	1:C:124:GLY:N	2.52	0.41
1:C:99:GLN:NE2	1:C:103:TYR:HE1	2.18	0.41
1:D:129:TYR:CE2	1:D:131:ARG:HD3	2.55	0.41
1:A:94:VAL:HG12	1:A:95:GLN:O	2.21	0.41

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	117/145 (81%)	109 (93%)	5 (4%)	3 (3%)	5	7
1	B	137/145 (94%)	127 (93%)	10 (7%)	0	100	100
1	C	116/145 (80%)	110 (95%)	3 (3%)	3 (3%)	5	7
1	D	136/145 (94%)	131 (96%)	5 (4%)	0	100	100
All	All	506/580 (87%)	477 (94%)	23 (4%)	6 (1%)	13	19

All (6) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	85	GLY
1	A	138	ARG
1	C	138	ARG
1	C	123	ASN

Continued on next page...

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	136	GLY
1	C	135	ALA

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	108/129 (84%)	93 (86%)	15 (14%)	3	4
1	B	123/129 (95%)	115 (94%)	8 (6%)	17	26
1	C	107/129 (83%)	97 (91%)	10 (9%)	9	13
1	D	122/129 (95%)	118 (97%)	4 (3%)	38	54
All	All	460/516 (89%)	423 (92%)	37 (8%)	12	18

All (37) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	3	ASN
1	A	11	VAL
1	A	13	TYR
1	A	17	LYS
1	A	21	SER
1	A	37	MET
1	A	55	ILE
1	A	59	LEU
1	A	90	THR
1	A	107	LYS
1	A	130	VAL
1	A	137	SER
1	A	138	ARG
1	A	143	LEU
1	A	144	GLU
1	B	7	SER
1	B	36	ARG
1	B	45	ASP
1	B	60	VAL

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	66	SER
1	B	69	THR
1	B	133	GLN
1	B	144	GLU
1	C	2	GLN
1	C	5	VAL
1	C	12	SER
1	C	13	TYR
1	C	25	LYS
1	C	44	ASP
1	C	112	ILE
1	C	123	ASN
1	C	131	ARG
1	C	137	SER
1	D	39	ARG
1	D	60	VAL
1	D	94	VAL
1	D	106	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	95	GLN
1	B	62	GLN
1	B	86	HIS
1	B	133	GLN
1	C	99	GLN
1	C	133	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no monosaccharides in this entry.

## 5.6 Ligand geometry

Of 3 ligands modelled in this entry, 3 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	121/145 (83%)	0.33	2 (1%) 70 67	24, 61, 73, 97	0
1	B	139/145 (95%)	-0.02	1 (0%) 87 87	24, 57, 68, 81	0
1	C	120/145 (82%)	0.33	1 (0%) 86 85	24, 61, 75, 85	0
1	D	138/145 (95%)	0.06	2 (1%) 75 73	24, 60, 71, 77	0
All	All	518/580 (89%)	0.16	6 (1%) 79 77	24, 60, 72, 97	0

All (6) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	84	ILE	3.2
1	B	83	LYS	2.3
1	C	13	TYR	2.3
1	A	135	ALA	2.2
1	D	63	TYR	2.2
1	D	14	PHE	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	MG	D	146	1/1	0.75	0.25	78,78,78,78	0
2	MG	A	146	1/1	0.84	0.56	88,88,88,88	0
2	MG	B	146	1/1	0.92	0.17	61,61,61,61	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.