



wwPDB X-ray Structure Validation Summary Report ⓘ

Oct 25, 2023 – 12:39 AM EDT

PDB ID : 3EHB
Title : A D-Pathway Mutation Decouples the Paracoccus Denitrificans Cytochrome c Oxidase by Altering the side chain orientation of a distant, conserved Glutamate
Authors : Koepke, J.; Mueller, H.; Peng, G.
Deposited on : 2008-09-12
Resolution : 2.32 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtrriage (Phenix) : 1.13
EDS : 2.36
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.32 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5974 (2.34-2.30)
Clashscore	141614	6604 (2.34-2.30)
Ramachandran outliers	138981	6523 (2.34-2.30)
Sidechain outliers	138945	6523 (2.34-2.30)
RSRZ outliers	127900	5855 (2.34-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	558	 5% 62% 28% 5% 5%
2	B	298	 % 59% 22% 15%
3	C	127	 7% 64% 25% 5% 6%
4	D	120	 % 64% 22% 9%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
10	LMT	A	571	-	-	-	X
10	LMT	A	572	-	-	-	X
10	LMT	A	573	-	-	-	X
10	LMT	A	574	-	-	-	X
10	LMT	A	575	-	-	-	X
10	LMT	B	279	-	-	-	X
10	LMT	D	121	-	-	-	X
5	HEA	A	559	X	-	-	-
5	HEA	A	560	X	-	-	-
9	LDA	A	564	-	-	-	X
9	LDA	B	276	-	-	-	X

2 Entry composition [i](#)

There are 12 unique types of molecules in this entry. The entry contains 9036 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Cytochrome c oxidase subunit 1-beta.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	530	4199	2818	656	692	33	0	3	1

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	131	ASP	ASN	engineered mutation	UNP P98002

- Molecule 2 is a protein called Cytochrome c oxidase subunit 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	253	1982	1298	320	356	8	0	1	1


- Molecule 3 is a protein called FV fragment Chain H.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	C	119	941	591	159	185	6	0	2	1

- Molecule 4 is a protein called FV fragment Chain L.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
4	D	109	832	530	136	164	2	0	0	1

- Molecule 5 is HEME-A (three-letter code: HEA) (formula: C₄₉H₅₆FeN₄O₆).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
5	A	1	Total	C	Fe	N	O	0	0
			60	49	1	4	6		
5	A	1	Total	C	Fe	N	O	0	0
			60	49	1	4	6		

- Molecule 6 is COPPER (II) ION (three-letter code: CU) (formula: Cu).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	1	Total	Cu	0	0
			1	1		
6	B	2	Total	Cu	0	0
			2	2		


- Molecule 7 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	1	Total	Mg	0	0
			1	1		

- Molecule 8 is CALCIUM ION (three-letter code: CA) (formula: Ca).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	1	Total	Ca	0	0
			1	1		

- Molecule 9 is LAURYL DIMETHYLAMINE-N-OXIDE (three-letter code: LDA) (formula: C₁₄H₃₁NO).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
9	A	1	Total 16	C 14	N 1	O 1	0	0
9	A	1	Total 16	C 14	N 1	O 1	0	0
9	A	1	Total 16	C 14	N 1	O 1	0	0
9	A	1	Total 16	C 14	N 1	O 1	0	0
9	B	1	Total 16	C 14	N 1	O 1	0	0
9	B	1	Total 16	C 14	N 1	O 1	0	0
9	B	1	Total 16	C 14	N 1	O 1	0	0
9	B	1	Total 16	C 14	N 1	O 1	0	0
9	B	1	Total 16	C 14	N 1	O 1	0	0

- Molecule 10 is DODECYL-BETA-D-MALTOSE (three-letter code: LMT) (formula: $C_{24}H_{46}O_{11}$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
10	A	1	Total	C	O	0	0
			35	24	11		
10	A	1	Total	C	O	0	0
			35	24	11		
10	A	1	Total	C	O	0	0
			35	24	11		
10	A	1	Total	C	O	0	0
			35	24	11		
10	A	1	Total	C	O	0	0
			35	24	11		
10	A	1	Total	C	O	0	0
			35	24	11		
10	A	1	Total	C	O	0	0
			35	24	11		
10	B	1	Total	C	O	0	1
			47	36	11		
10	B	1	Total	C	O	0	0
			35	24	11		
10	B	1	Total	C	O	0	0
			35	24	11		
10	D	1	Total	C	O	0	0
			35	24	11		

- Molecule 11 is PEROXIDE ION (three-letter code: PER) (formula: O₂).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
11	A	1	Total O 2 2	0	0


- Molecule 12 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
12	A	117	Total O 117 117	0	0
12	B	138	Total O 138 138	0	0
12	C	70	Total O 70 70	0	0
12	D	54	Total O 54 54	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Cytochrome c oxidase subunit 1-beta




• Molecule 2: Cytochrome c oxidase subunit 2



- Molecule 3: FV fragment Chain H



- Molecule 4: FV fragment Chain L



4 Data and refinement statistics i

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	84.46Å 151.33Å 157.49Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	19.69 – 2.32 19.69 – 2.32	Depositor EDS
% Data completeness (in resolution range)	100.0 (19.69-2.32) 98.9 (19.69-2.32)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.35 (at 2.33Å)	Xtrriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.205 , 0.242 0.204 , 0.205	Depositor DCC
R_{free} test set	2629 reflections (3.02%)	wwPDB-VP
Wilson B-factor (Å ²)	43.8	Xtrriage
Anisotropy	0.268	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 64.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	0.010 for -h,l,k	Xtrriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	9036	wwPDB-VP
Average B, all atoms (Å ²)	60.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.95% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CU, LDA, CA, LMT, PER, HEA, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	1.20	6/4367 (0.1%)	1.08	18/5959 (0.3%)
2	B	1.38	7/2043 (0.3%)	1.18	10/2801 (0.4%)
3	C	1.14	0/973	1.05	1/1316 (0.1%)
4	D	1.28	2/853 (0.2%)	1.10	0/1158
All	All	1.25	15/8236 (0.2%)	1.11	29/11234 (0.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
3	C	0	1

The worst 5 of 15 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	D	23	CYS	CB-SG	-8.64	1.67	1.82
2	B	125	TYR	CE2-CZ	7.65	1.48	1.38
1	A	407	TYR	CD2-CE2	7.47	1.50	1.39
2	B	182	ALA	CA-CB	6.48	1.66	1.52
1	A	460	PHE	CE2-CZ	6.47	1.49	1.37

The worst 5 of 29 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	54	ARG	NE-CZ-NH2	-16.54	112.03	120.30
1	A	468	ARG	NE-CZ-NH2	-15.06	112.77	120.30
1	A	54	ARG	NE-CZ-NH1	8.99	124.79	120.30
1	A	263	ASP	CB-CG-OD1	8.92	126.33	118.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	249	ARG	NE-CZ-NH2	-8.12	116.24	120.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
3	C	8	GLY	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4199	0	4116	177	0
2	B	1982	0	1965	63	0
3	C	941	0	898	26	0
4	D	832	0	807	23	0
5	A	120	0	104	24	0
6	A	1	0	0	0	0
6	B	2	0	0	0	0
7	A	1	0	0	0	0
8	A	1	0	0	0	0
9	A	64	0	124	5	0
9	B	80	0	155	15	0
10	A	280	0	335	27	0
10	B	117	0	133	8	0
10	D	35	0	39	3	0
11	A	2	0	0	0	0
12	A	117	0	0	4	0
12	B	138	0	0	5	0
12	C	70	0	0	4	0
12	D	54	0	0	2	0
All	All	9036	0	8676	305	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 18.

The worst 5 of 305 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:222:ILE:CD1	2:B:222:ILE:CG1	1.85	1.55
1:A:276:HIS:NE2	1:A:280:TYR:HE2	1.07	1.44
1:A:276:HIS:NE2	1:A:280:TYR:CE2	1.96	1.31
10:A:575:LMT:O6B	10:A:575:LMT:C6B	1.81	1.28
1:A:342:LEU:HA	1:A:345:MET:CE	1.78	1.13

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	531/558 (95%)	494 (93%)	34 (6%)	3 (1%)	25	30
2	B	252/298 (85%)	240 (95%)	10 (4%)	2 (1%)	19	23
3	C	119/127 (94%)	110 (92%)	8 (7%)	1 (1%)	19	23
4	D	107/120 (89%)	101 (94%)	4 (4%)	2 (2%)	8	6
All	All	1009/1103 (92%)	945 (94%)	56 (6%)	8 (1%)	19	23

5 of 8 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	74	ASN
3	C	118	SER
1	A	79	GLU
4	D	77	SER
2	B	235	SER

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	435/454 (96%)	405 (93%)	30 (7%)	15	20
2	B	212/243 (87%)	194 (92%)	18 (8%)	10	12
3	C	103/107 (96%)	94 (91%)	9 (9%)	10	12
4	D	92/104 (88%)	82 (89%)	10 (11%)	6	7
All	All	842/908 (93%)	775 (92%)	67 (8%)	13	15

5 of 67 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
4	D	1	ASP
4	D	14	SER
4	D	107	ILE
1	A	388	VAL
1	A	380	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 25 such sidechains are listed below:

Mol	Chain	Res	Type
2	B	29	HIS
2	B	108	ASN
4	D	91	HIS
2	B	104	GLN
2	B	173	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry

Of 30 ligands modelled in this entry, 5 are monoatomic - leaving 25 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	HEA	A	559	1	57,67,67	3.10	17 (29%)	61,103,103	4.42	37 (60%)
9	LDA	B	276	-	12,15,15	2.03	1 (8%)	14,17,17	0.44	0
9	LDA	A	565	-	12,15,15	1.91	1 (8%)	14,17,17	0.50	0
9	LDA	B	274	-	12,15,15	1.98	1 (8%)	14,17,17	0.49	0
9	LDA	B	273	-	12,15,15	2.01	1 (8%)	14,17,17	0.61	0
10	LMT	A	571	-	36,36,36	4.36	16 (44%)	47,47,47	1.51	7 (14%)
10	LMT	A	570	-	36,36,36	4.32	16 (44%)	47,47,47	1.45	5 (10%)
10	LMT	B	278	-	36,36,36	4.29	13 (36%)	47,47,47	1.60	9 (19%)
10	LMT	B	277[A]	-	36,36,36	4.32	15 (41%)	47,47,47	1.90	9 (19%)
9	LDA	B	275	-	12,15,15	1.91	1 (8%)	14,17,17	0.60	0
10	LMT	A	568	-	36,36,36	4.41	14 (38%)	47,47,47	1.22	3 (6%)
10	LMT	A	574	-	36,36,36	4.33	15 (41%)	47,47,47	1.62	6 (12%)
10	LMT	A	569	-	36,36,36	4.42	17 (47%)	47,47,47	1.93	13 (27%)
9	LDA	A	566	-	12,15,15	1.90	1 (8%)	14,17,17	0.56	0
9	LDA	A	564	-	12,15,15	2.02	1 (8%)	14,17,17	0.65	0
11	PER	A	576	5	0,1,1	-	-	-	-	-
10	LMT	A	572	-	36,36,36	4.10	17 (47%)	47,47,47	1.51	5 (10%)
10	LMT	A	573	-	36,36,36	4.10	16 (44%)	47,47,47	1.63	7 (14%)
5	HEA	A	560	1,11	57,67,67	3.68	25 (43%)	61,103,103	4.31	37 (60%)
10	LMT	B	277[B]	-	36,36,36	4.32	15 (41%)	47,47,47	1.80	9 (19%)
9	LDA	A	567	-	12,15,15	2.00	1 (8%)	14,17,17	0.41	0
10	LMT	A	575	-	36,36,36	4.51	16 (44%)	47,47,47	1.55	7 (14%)
10	LMT	D	121	-	36,36,36	4.15	15 (41%)	47,47,47	2.17	12 (25%)
10	LMT	B	279	-	36,36,36	4.32	14 (38%)	47,47,47	1.23	5 (10%)
9	LDA	B	272	-	12,15,15	1.95	1 (8%)	14,17,17	0.69	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	HEA	A	559	1	1/1/7/16	13/32/76/76	-
9	LDA	B	276	-	-	8/13/13/13	-
9	LDA	A	565	-	-	8/13/13/13	-
9	LDA	B	274	-	-	10/13/13/13	-
9	LDA	B	273	-	-	9/13/13/13	-
10	LMT	A	571	-	-	15/21/61/61	0/2/2/2
10	LMT	A	570	-	-	11/21/61/61	0/2/2/2
10	LMT	B	278	-	-	14/21/61/61	0/2/2/2
10	LMT	B	277[A]	-	-	10/21/61/61	0/2/2/2
9	LDA	B	275	-	-	4/13/13/13	-
10	LMT	A	568	-	-	11/21/61/61	0/2/2/2
10	LMT	A	574	-	-	16/21/61/61	0/2/2/2
10	LMT	A	569	-	-	9/21/61/61	0/2/2/2
9	LDA	A	566	-	-	7/13/13/13	-
9	LDA	A	564	-	-	7/13/13/13	-
10	LMT	A	572	-	-	13/21/61/61	0/2/2/2
10	LMT	A	573	-	-	10/21/61/61	0/2/2/2
5	HEA	A	560	1,11	1/1/7/16	17/32/76/76	-
10	LMT	B	277[B]	-	-	5/21/61/61	0/2/2/2
9	LDA	A	567	-	-	9/13/13/13	-
10	LMT	A	575	-	-	15/21/61/61	0/2/2/2
10	LMT	D	121	-	-	14/21/61/61	0/2/2/2
10	LMT	B	279	-	-	13/21/61/61	0/2/2/2
9	LDA	B	272	-	-	6/13/13/13	-

The worst 5 of 250 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	A	560	HEA	O11-C11	-12.28	1.16	1.42
5	A	560	HEA	C14-C15	10.68	1.58	1.33
5	A	559	HEA	O11-C11	-9.38	1.22	1.42
10	A	575	LMT	O6B-C6B	9.17	1.81	1.42
10	A	569	LMT	O2B-C2B	-8.91	1.22	1.43

The worst 5 of 171 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
5	A	560	HEA	C13-C12-C11	-16.19	90.03	114.35
5	A	559	HEA	CHB-C1B-NB	9.97	135.26	124.43
5	A	559	HEA	C17-C18-C19	-9.77	104.14	127.66
5	A	559	HEA	C4A-CHB-C1B	-9.15	110.48	122.56
5	A	559	HEA	C16-C15-C14	-9.13	102.64	121.12

All (2) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
5	A	559	HEA	C11
5	A	560	HEA	C11

5 of 254 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	A	559	HEA	O11-C11-C3B-C2B
5	A	559	HEA	C12-C11-C3B-C2B
5	A	559	HEA	C12-C11-C3B-C4B
5	A	559	HEA	O11-C11-C12-C13
5	A	559	HEA	C17-C18-C19-C27

There are no ring outliers.

20 monomers are involved in 78 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	559	HEA	8	0
9	B	276	LDA	1	0
9	A	565	LDA	1	0
9	B	274	LDA	4	0
9	B	273	LDA	3	0
10	A	571	LMT	3	0
10	A	570	LMT	3	0
10	B	278	LMT	2	0
9	B	275	LDA	4	0
10	A	568	LMT	2	0
10	A	574	LMT	9	0
10	A	569	LMT	2	0
9	A	566	LDA	3	0
9	A	564	LDA	1	0
10	A	572	LMT	3	0
5	A	560	HEA	16	0


Continued on next page...


Continued from previous page...


Mol	Chain	Res	Type	Clashes	Symm-Clashes
10	A	575	LMT	5	0
10	D	121	LMT	3	0
10	B	279	LMT	6	0
9	B	272	LDA	4	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.














5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	530/558 (94%)	-0.16	28 (5%) 26 33	41, 55, 79, 125	0
2	B	253/298 (84%)	-0.46	2 (0%) 86 89	36, 50, 70, 86	0
3	C	119/127 (93%)	0.16	9 (7%) 13 18	39, 60, 82, 112	0
4	D	109/120 (90%)	-0.15	1 (0%) 84 88	42, 57, 71, 82	0
All	All	1011/1103 (91%)	-0.20	40 (3%) 38 45	36, 55, 77, 125	0

The worst 5 of 40 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	C	119	ALA	8.1
1	A	545	THR	7.9
1	A	17	GLY	7.2
1	A	18	PHE	6.4
1	A	546	LEU	6.4

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
10	LMT	A	575	35/35	0.29	0.46	97,142,149,150	0
10	LMT	A	573	35/35	0.36	0.42	65,102,131,132	0
10	LMT	A	572	35/35	0.36	0.42	84,112,123,123	0
10	LMT	A	571	35/35	0.39	0.44	81,116,145,146	0
10	LMT	A	574	35/35	0.43	0.45	64,134,147,148	0
9	LDA	B	272	16/16	0.44	0.38	46,67,114,117	0
9	LDA	A	564	16/16	0.44	0.43	68,75,106,109	0
9	LDA	B	276	16/16	0.48	0.41	121,128,130,130	0
10	LMT	D	121	35/35	0.54	0.41	93,103,108,110	0
10	LMT	A	569	35/35	0.63	0.36	84,98,111,111	0
9	LDA	A	565	16/16	0.65	0.31	72,85,94,95	0
9	LDA	B	274	16/16	0.65	0.28	83,98,111,111	0
9	LDA	A	567	16/16	0.66	0.27	96,103,104,105	0
9	LDA	A	566	16/16	0.68	0.33	92,97,113,113	0
10	LMT	A	570	35/35	0.70	0.27	69,104,122,125	0
9	LDA	B	275	16/16	0.74	0.27	55,81,113,115	0
10	LMT	B	279	35/35	0.76	0.40	68,120,136,137	0
9	LDA	B	273	16/16	0.76	0.28	81,85,98,100	0
10	LMT	A	568	35/35	0.78	0.34	83,103,108,109	0
10	LMT	B	278	35/35	0.84	0.29	68,86,99,100	0
10	LMT	B	277[B]	35/35	0.85	0.28	65,78,87,88	12
10	LMT	B	277[A]	35/35	0.85	0.28	67,81,87,88	12
5	HEA	A	559	60/60	0.95	0.16	37,47,51,56	0
5	HEA	A	560	60/60	0.96	0.13	34,52,54,62	0
8	CA	A	563	1/1	0.97	0.06	53,53,53,53	0
7	MG	A	562	1/1	0.98	0.15	28,28,28,28	0
6	CU	B	270	1/1	0.99	0.07	48,48,48,48	0
6	CU	B	271	1/1	0.99	0.06	50,50,50,50	0
11	PER	A	576	2/2	0.99	0.10	49,49,49,56	0
6	CU	A	561	1/1	1.00	0.03	54,54,54,54	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around LMT A 575:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around LMT A 573:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around LMT A 572:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around LMT A 571:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around LMT A 574:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around LMT D 121:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around LMT A 569:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around LMT A 570:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around LMT B 279:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around LMT A 568:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around LMT B 278:


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around LMT B 277 (B):**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







6.5 Other polymers [i](#)

There are no such residues in this entry.