



Full wwPDB X-ray Structure Validation Report ⓘ

Jun 12, 2024 – 04:36 PM EDT

PDB ID : 1FIW
Title : THREE-DIMENSIONAL STRUCTURE OF BETA-ACROSIN FROM RAM SPERMATOOZA
Authors : Tranter, R.; Read, J.A.; Jones, R.; Brady, R.L.
Deposited on : 2000-08-07
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36.2
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION




The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5197 (2.10-2.10)
Clashscore	141614	5710 (2.10-2.10)
Ramachandran outliers	138981	5647 (2.10-2.10)
Sidechain outliers	138945	5648 (2.10-2.10)
RSRZ outliers	127900	5083 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	290	
2	L	22	
3	B	5	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	NAG	B	2	-	-	X	-
3	BMA	B	3	-	-	X	X
3	MAN	B	4	X	-	-	-

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 2409 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called BETA-ACROSIN HEAVY CHAIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	260	Total	C	N	O	S	69	0	0
			2032	1294	364	359	15			

- Molecule 2 is a protein called BETA-ACROSIN LIGHT CHAIN.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	L	14	Total	C	N	O	S	0	0	0
			106	62	22	20	2			

- Molecule 3 is an oligosaccharide called alpha-D-mannopyranose-(1-6)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[beta-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	B	5	Total	C	N	O	0	0	0
			60	34	2	24			

- Molecule 4 is P-AMINO BENZAMIDINE (three-letter code: PBZ) (formula: C₇H₁₀N₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	N	0	0
			10	7	3		
4	A	1	Total	C	N	0	0
			10	7	3		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	181	Total	O	0	0
			181	181		
5	L	10	Total	O	0	0
			10	10		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: BETA-ACROSIN HEAVY CHAIN



• Molecule 2: BETA-ACROSIN LIGHT CHAIN



• Molecule 3: alpha-D-mannopyranose-(1-6)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[beta-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose



4 Data and refinement statistics

Property	Value	Source
Space group	P 65 2 2	Depositor
Cell constants a, b, c, α , β , γ	105.29Å 105.29Å 120.83Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	30.00 – 2.10 29.94 – 2.10	Depositor EDS
% Data completeness (in resolution range)	91.7 (30.00-2.10) 91.7 (29.94-2.10)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.35 (at 2.10Å)	Xtriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.197 , 0.242 0.195 , 0.240	Depositor DCC
R_{free} test set	1145 reflections (5.14%)	wwPDB-VP
Wilson B-factor (Å ²)	28.3	Xtriage
Anisotropy	0.618	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 48.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	2409	wwPDB-VP
Average B, all atoms (Å ²)	39.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.66% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PBZ, MAN, NAG, FUL, BMA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.14	7/2091 (0.3%)	1.08	10/2846 (0.4%)
2	L	0.80	0/107	0.98	0/143
All	All	1.13	7/2198 (0.3%)	1.08	10/2989 (0.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	202(B)	GLU	C-N	-12.74	1.04	1.34
1	A	74	GLU	CB-CG	8.62	1.68	1.52
1	A	94	TYR	CD1-CE1	-7.64	1.27	1.39
1	A	149	ARG	CA-CB	-7.58	1.37	1.53
1	A	77	LYS	CA-CB	-6.88	1.38	1.53
1	A	217	VAL	C-N	6.47	1.44	1.33
1	A	222	ALA	CA-CB	5.63	1.64	1.52

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	21	ASP	CB-CG-OD1	8.91	126.31	118.30
1	A	194	ASP	CB-CG-OD2	8.66	126.09	118.30
1	A	189	ASP	CB-CG-OD1	7.58	125.12	118.30
1	A	204	VAL	O-C-N	-6.37	112.52	122.70

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	66	PHE	O-C-N	-6.23	112.60	123.20
1	A	114	CYS	CA-CB-SG	-5.78	103.59	114.00
1	A	66	PHE	C-N-CA	5.64	134.15	122.30
1	A	77	LYS	N-CA-CB	5.50	120.49	110.60
1	A	200	MET	CG-SD-CE	5.31	108.70	100.20
1	A	64	ARG	NE-CZ-NH1	-5.05	117.78	120.30

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	126(A)	ARG	Sidechain
1	A	130	ARG	Mainchain

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2032	0	1995	49	1
2	L	106	0	96	6	0
3	B	60	0	52	9	0
4	A	20	0	20	1	0
5	A	181	0	0	4	0
5	L	10	0	0	0	0
All	All	2409	0	2163	58	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 14.

All (58) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:B:2:NAG:O3	3:B:3:BMA:H2	1.36	1.20
1:A:25:GLY:HA3	2:L:15:GLN:NE2	1.58	1.17
1:A:24:HIS:CE1	1:A:117:PHE:CD2	2.39	1.11

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:37:ILE:HD12	1:A:41:VAL:HG21	1.37	1.06
1:A:24:HIS:CE1	1:A:117:PHE:CE2	2.45	1.04
1:A:24:HIS:ND1	1:A:117:PHE:CE2	2.26	1.03
3:B:2:NAG:H4	3:B:3:BMA:O2	1.32	0.96
1:A:25:GLY:HA3	2:L:15:GLN:HE22	1.26	0.94
1:A:24:HIS:HD1	1:A:117:PHE:HE2	1.17	0.91
3:B:2:NAG:C4	3:B:3:BMA:O2	2.17	0.88
1:A:79:PRO:HG2	5:A:424:HOH:O	1.74	0.87
3:B:2:NAG:H4	3:B:3:BMA:HO2	1.40	0.86
3:B:2:NAG:C3	3:B:3:BMA:H2	1.94	0.79
3:B:2:NAG:HO3	3:B:3:BMA:H2	1.45	0.79
1:A:37:ILE:CD1	1:A:41:VAL:HG21	2.14	0.77
1:A:24:HIS:HE1	1:A:117:PHE:CD2	1.99	0.76
1:A:254:THR:HG21	5:A:465:HOH:O	1.85	0.76
1:A:24:HIS:ND1	1:A:117:PHE:CD2	2.55	0.71
1:A:187:GLY:O	1:A:188:LYS:HB2	1.89	0.71
1:A:38:ARG:HH12	1:A:75(E):PRO:HA	1.56	0.70
1:A:173:ASN:ND2	3:B:5:FUL:H3	2.08	0.69
1:A:86:GLU:HG2	5:A:472:HOH:O	1.92	0.68
1:A:75:TRP:CE2	1:A:150:ARG:HG2	2.32	0.64
1:A:71:LYS:HZ2	1:A:154:MET:HE1	1.62	0.64
1:A:37(E):ASN:C	1:A:37(E):ASN:HD22	2.05	0.60
1:A:25:GLY:CA	2:L:15:GLN:HE22	2.09	0.59
1:A:91:HIS:CE1	1:A:93:LYS:HB2	2.38	0.59
1:A:52:LEU:C	1:A:52:LEU:HD12	2.22	0.59
2:L:6:ASP:N	2:L:6:ASP:OD1	2.37	0.57
1:A:90:ILE:HD13	1:A:104:ALA:HB2	1.86	0.57
1:A:25:GLY:CA	2:L:15:GLN:NE2	2.51	0.56
1:A:65(A):ILE:HD12	1:A:76:VAL:HG11	1.89	0.55
1:A:39:TYR:O	1:A:41:VAL:HG23	2.08	0.54
1:A:71:LYS:NZ	1:A:154:MET:HE1	2.23	0.53
1:A:24:HIS:HE1	1:A:117:PHE:CE2	2.15	0.52
1:A:59:PHE:HD2	1:A:90:ILE:HD11	1.74	0.51
1:A:173:ASN:HD22	3:B:5:FUL:H3	1.76	0.48
1:A:99:GLU:HA	1:A:99:GLU:OE1	2.15	0.47
1:A:59:PHE:CD2	1:A:90:ILE:HD11	2.50	0.46
1:A:71:LYS:NZ	1:A:154:MET:CE	2.79	0.46
1:A:77:LYS:O	1:A:78:PRO:C	2.53	0.46
1:A:141:TRP:O	1:A:151:THR:HB	2.16	0.46
1:A:203:TYR:OH	5:A:359:HOH:O	2.19	0.45
1:A:90:ILE:HD13	1:A:104:ALA:CB	2.47	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:37(E):ASN:ND2	1:A:37(G):ARG:H	2.15	0.44
1:A:41:VAL:O	1:A:193:GLY:HA2	2.16	0.44
1:A:61(A):LYS:HB3	1:A:62:ASP:HB3	1.99	0.43
1:A:52:LEU:HD12	1:A:52:LEU:O	2.19	0.43
2:L:3:THR:HG23	2:L:3:THR:O	2.19	0.43
1:A:60:ARG:HD3	1:A:96:ALA:HB2	2.00	0.42
3:B:2:NAG:H81	3:B:5:FUL:H61	2.01	0.42
1:A:213:THR:HG1	1:A:228:TYR:HE2	1.65	0.42
1:A:101:ASN:ND2	1:A:234:TYR:OH	2.49	0.42
1:A:37:ILE:CG1	1:A:41:VAL:HG21	2.50	0.41
1:A:24:HIS:HE1	1:A:117:PHE:CG	2.37	0.41
1:A:51:TRP:CE2	1:A:242:ILE:HG22	2.56	0.41
1:A:61(A):LYS:HB3	1:A:62:ASP:CB	2.51	0.41
1:A:192:GLN:NE2	4:A:305:PBZ:H2	2.36	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:175:ARG:NH2	1:A:175:ARG:NH2[11_655]	2.01	0.19

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	258/290 (89%)	247 (96%)	9 (4%)	2 (1%)	19	15
2	L	12/22 (54%)	11 (92%)	0	1 (8%)	1	0
All	All	270/312 (86%)	258 (96%)	9 (3%)	3 (1%)	14	9

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	201(B)	ASP
1	A	202	SER
2	L	14	ARG

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	216/240 (90%)	211 (98%)	5 (2%)	50	55
2	L	12/18 (67%)	7 (58%)	5 (42%)	0	0
All	All	228/258 (88%)	218 (96%)	10 (4%)	28	28

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	37(E)	ASN
1	A	60	ARG
1	A	201(A)	LYS
1	A	202(D)	SER
1	A	254	THR
2	L	3	THR
2	L	6	ASP
2	L	11	VAL
2	L	15	GLN
2	L	16	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	37(E)	ASN
1	A	101	ASN
1	A	173	ASN
2	L	15	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

5 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
3	NAG	B	1	1,3	14,14,15	1.14	1 (7%)	17,19,21	1.91	5 (29%)
3	NAG	B	2	3	14,14,15	1.33	1 (7%)	17,19,21	2.27	7 (41%)
3	BMA	B	3	3	11,11,12	0.86	0	15,15,17	3.03	9 (60%)
3	MAN	B	4	3	11,11,12	0.76	0	15,15,17	3.09	4 (26%)
3	FUL	B	5	3	10,10,11	0.95	1 (10%)	14,14,16	2.66	4 (28%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	B	1	1,3	-	2/6/23/26	0/1/1/1
3	NAG	B	2	3	-	0/6/23/26	0/1/1/1
3	BMA	B	3	3	-	2/2/19/22	0/1/1/1
3	MAN	B	4	3	1/1/4/5	0/2/19/22	0/1/1/1
3	FUL	B	5	3	-	-	0/1/1/1

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	2	NAG	O5-C1	-4.58	1.36	1.43
3	B	1	NAG	C8-C7	3.01	1.56	1.50
3	B	5	FUL	O3-C3	-2.09	1.38	1.43

All (29) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	4	MAN	C1-C2-C3	-8.93	98.70	109.67
3	B	5	FUL	C1-O5-C5	-7.10	96.69	112.78
3	B	3	BMA	C1-C2-C3	6.29	117.40	109.67
3	B	4	MAN	C1-O5-C5	5.95	120.25	112.19
3	B	3	BMA	C1-O5-C5	5.78	120.03	112.19
3	B	2	NAG	C4-C3-C2	4.60	117.75	111.02
3	B	1	NAG	O5-C5-C6	4.58	114.38	107.20
3	B	2	NAG	O4-C4-C5	4.25	119.86	109.30
3	B	5	FUL	O3-C3-C2	-4.24	101.88	109.99
3	B	2	NAG	O5-C1-C2	-3.88	105.16	111.29
3	B	3	BMA	O5-C5-C4	3.88	120.25	110.83
3	B	3	BMA	C2-C3-C4	-3.51	104.83	110.89
3	B	5	FUL	C2-C3-C4	3.26	116.54	110.89
3	B	1	NAG	C6-C5-C4	-3.25	105.39	113.00
3	B	5	FUL	O5-C5-C6	3.19	114.20	107.33
3	B	4	MAN	O5-C5-C4	3.16	118.51	110.83
3	B	3	BMA	O3-C3-C2	-3.12	104.01	109.99
3	B	2	NAG	C3-C4-C5	2.85	115.32	110.24
3	B	1	NAG	C3-C4-C5	-2.69	105.44	110.24
3	B	3	BMA	O6-C6-C5	-2.57	102.46	111.29
3	B	2	NAG	O3-C3-C4	-2.45	104.70	110.35
3	B	2	NAG	O5-C5-C4	-2.30	105.24	110.83
3	B	3	BMA	C6-C5-C4	-2.30	107.63	113.00
3	B	3	BMA	O2-C2-C3	-2.29	105.54	110.14
3	B	2	NAG	O5-C5-C6	-2.27	103.64	107.20
3	B	1	NAG	O5-C5-C4	2.16	116.09	110.83
3	B	3	BMA	C3-C4-C5	2.14	114.05	110.24
3	B	4	MAN	O2-C2-C1	2.11	113.48	109.15
3	B	1	NAG	O4-C4-C5	-2.11	104.05	109.30

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
3	B	4	MAN	C5

All (4) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	B	3	BMA	O5-C5-C6-O6
3	B	3	BMA	C4-C5-C6-O6
3	B	1	NAG	C4-C5-C6-O6
3	B	1	NAG	O5-C5-C6-O6

There are no ring outliers.

3 monomers are involved in 9 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	5	FUL	3	0
3	B	2	NAG	7	0
3	B	3	BMA	6	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	PBZ	A	305	-	10,10,10	1.92	2 (20%)	9,13,13	0.29	0
4	PBZ	A	306	-	10,10,10	2.20	2 (20%)	9,13,13	2.11	5 (55%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	PBZ	A	305	-	-	1/4/4/4	0/1/1/1
4	PBZ	A	306	-	-	2/4/4/4	0/1/1/1

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	306	PBZ	C7-N3	4.96	1.49	1.28
4	A	306	PBZ	C7-N2	4.47	1.45	1.33
4	A	305	PBZ	C7-N3	4.12	1.45	1.28
4	A	305	PBZ	C7-N2	3.74	1.43	1.33

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	306	PBZ	C5-C6-C1	-3.04	116.77	120.67
4	A	306	PBZ	C2-C3-C4	-2.90	117.40	120.78
4	A	306	PBZ	C2-C1-N1	-2.88	115.55	120.91
4	A	306	PBZ	C5-C4-C3	2.76	122.52	118.59
4	A	306	PBZ	C6-C1-C2	2.30	121.71	118.15

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	305	PBZ	C5-C4-C7-N2
4	A	306	PBZ	C3-C4-C7-N3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
4	A	306	PBZ	C5-C4-C7-N3

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	305	PBZ	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
1	A	1

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	A	202(B):GLU	C	202(C):ASN	N	1.04

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	256/290 (88%)	0.07	13 (5%) 28 33	26, 34, 53, 69	8 (3%)
2	L	14/22 (63%)	0.94	4 (28%) 0 0	40, 50, 64, 66	0
All	All	270/312 (86%)	0.12	17 (6%) 20 24	26, 34, 55, 69	8 (2%)

All (17) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	254	THR	4.9
1	A	202(C)	ASN	4.6
2	L	15	GLN	3.5
2	L	6	ASP	3.2
1	A	24	HIS	3.0
2	L	3	THR	3.0
2	L	16	ASN	2.8
1	A	147	ASN	2.7
1	A	44	GLY	2.7
1	A	31	VAL	2.5
1	A	150	ARG	2.5
1	A	197	GLY	2.4
1	A	78	PRO	2.3
1	A	75(E)	PRO	2.3
1	A	37(D)	HIS	2.1
1	A	212	ILE	2.1
1	A	33	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)


There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	BMA	B	3	11/12	0.52	0.47	77,86,90,90	0
3	MAN	B	4	11/12	0.75	0.31	64,69,72,72	0
3	FUL	B	5	10/11	0.75	0.23	59,61,67,73	0
3	NAG	B	2	14/15	0.79	0.22	48,54,71,75	0
3	NAG	B	1	14/15	0.93	0.09	27,36,48,53	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.



6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
4	PBZ	A	306	10/10	0.92	0.14	36,43,45,46	0
4	PBZ	A	305	10/10	0.95	0.12	28,31,33,35	0

6.5 Other polymers [i](#)

There are no such residues in this entry.