



wwPDB X-ray Structure Validation Summary Report ⓘ

Jun 13, 2024 – 08:02 AM EDT

PDB ID : 4FQU
Title : Glutathionyl-Hydroquinone Reductase PcpF of *Sphingobium chlorophenolicum*
Authors : Green, A.R.; Hayes, R.P.; Xun, L.; Kang, C.
Deposited on : 2012-06-25
Resolution : 3.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.20.1
EDS	:	2.36.2
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.36.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2092 (3.00-3.00)
Clashscore	141614	2416 (3.00-3.00)
Ramachandran outliers	138981	2333 (3.00-3.00)
Sidechain outliers	138945	2336 (3.00-3.00)
RSRZ outliers	127900	1990 (3.00-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	313	<div> <div>2%</div> <div> <div></div> <div>55%</div> <div>33%</div> <div>10%</div> <div>.</div> </div> </div>
1	B	313	<div> <div>2%</div> <div> <div></div> <div>52%</div> <div>35%</div> <div>8%</div> <div>..</div> </div> </div>
1	C	313	<div> <div>2%</div> <div> <div></div> <div>45%</div> <div>43%</div> <div>9%</div> <div>.</div> </div> </div>
1	D	313	<div> <div>3%</div> <div> <div></div> <div>41%</div> <div>46%</div> <div>12%</div> <div>.</div> </div> </div>
1	E	313	<div> <div>3%</div> <div> <div></div> <div>45%</div> <div>40%</div> <div>13%</div> <div>.</div> </div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	313	<div><div></div><div>2%</div><div>55%</div><div>33%</div><div>7%</div><div></div><div></div></div>
1	G	313	<div><div></div><div>2%</div><div>46%</div><div>41%</div><div>9%</div><div></div><div></div></div>
1	H	313	<div><div></div><div>3%</div><div>38%</div><div>46%</div><div>13%</div><div></div><div></div></div>

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 19729 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Putative glutathione transferase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	312	Total	C	N	O	S	0	0	0
			2491	1592	435	458	6			
1	B	303	Total	C	N	O	S	0	0	0
			2420	1549	424	441	6			
1	C	312	Total	C	N	O	S	0	0	0
			2482	1584	435	457	6			
1	D	312	Total	C	N	O	S	0	0	0
			2481	1586	432	457	6			
1	E	312	Total	C	N	O	S	0	0	0
			2484	1587	435	456	6			
1	F	308	Total	C	N	O	S	0	0	0
			2444	1559	429	450	6			
1	G	305	Total	C	N	O	S	0	0	0
			2428	1551	426	445	6			
1	H	309	Total	C	N	O	S	0	0	0
			2444	1561	430	447	6			

- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	21	Total	O	0	0
			21	21		
2	B	5	Total	O	0	0
			5	5		
2	C	9	Total	O	0	0
			9	9		
2	D	3	Total	O	0	0
			3	3		
2	E	1	Total	O	0	0
			1	1		
2	F	13	Total	O	0	0
			13	13		

Continued on next page...


Continued from previous page...


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	G	2	Total	O	0	0
			2	2		
2	H	1	Total	O	0	0
			1	1		

3 Residue-property plots


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Putative glutathione transferase








• Molecule 1: Putative glutathione transferase




• Molecule 1: Putative glutathione transferase






● Molecule 1: Putative glutathione transferase



● Molecule 1: Putative glutathione transferase



● Molecule 1: Putative glutathione transferase





4 Data and refinement statistics

Property	Value	Source
Space group	P 21 3	Depositor
Cell constants a, b, c, α , β , γ	242.84Å 242.84Å 242.84Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	49.57 – 3.00 49.57 – 2.83	Depositor EDS
% Data completeness (in resolution range)	87.9 (49.57-3.00) 84.0 (49.57-2.83)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	0.36 (at 2.81Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.6.1_357)	Depositor
R, R_{free}	0.202 , 0.256 0.195 , 0.250	Depositor DCC
R_{free} test set	2007 reflections (1.78%)	wwPDB-VP
Wilson B-factor (Å ²)	43.0	Xtriage
Anisotropy	0.000	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 46.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.024 for l,k,h	Xtriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	19729	wwPDB-VP
Average B, all atoms (Å ²)	45.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 1.35% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.94	5/2562 (0.2%)	0.97	4/3490 (0.1%)
1	B	0.82	1/2488 (0.0%)	0.94	6/3387 (0.2%)
1	C	0.80	2/2552 (0.1%)	0.95	4/3476 (0.1%)
1	D	0.72	1/2552 (0.0%)	0.88	3/3478 (0.1%)
1	E	0.80	0/2554	0.92	4/3479 (0.1%)
1	F	0.93	2/2511 (0.1%)	0.95	2/3418 (0.1%)
1	G	0.68	0/2495	0.84	2/3397 (0.1%)
1	H	0.63	0/2511	0.79	0/3419
All	All	0.80	11/20225 (0.1%)	0.91	25/27544 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	3
1	C	0	2
1	D	0	1
1	E	0	1
1	F	0	2
All	All	0	9

The worst 5 of 11 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	18	SER	CB-OG	6.75	1.51	1.42
1	B	53	CYS	CB-SG	-6.33	1.71	1.82
1	A	53	CYS	CB-SG	-6.18	1.71	1.82
1	F	70	GLU	CG-CD	6.14	1.61	1.51
1	A	70	GLU	CG-CD	5.71	1.60	1.51

The worst 5 of 25 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	39	ARG	NE-CZ-NH1	8.07	124.34	120.30
1	E	76	MET	CG-SD-CE	-7.11	88.83	100.20
1	F	91	GLY	N-CA-C	6.47	129.29	113.10
1	C	91	GLY	N-CA-C	6.35	128.98	113.10
1	B	306	LEU	CA-CB-CG	6.27	129.72	115.30

There are no chirality outliers.

5 of 9 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	148	ALA	Peptide
1	A	39	ARG	Peptide
1	A	40	GLY	Peptide
1	C	36	ALA	Peptide
1	C	6	ASP	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2491	0	2406	178	0
1	B	2420	0	2340	152	0
1	C	2482	0	2392	196	0
1	D	2481	0	2383	236	0
1	E	2484	0	2385	189	0
1	F	2444	0	2354	147	0
1	G	2428	0	2344	153	0
1	H	2444	0	2351	207	0
2	A	21	0	0	3	0
2	B	5	0	0	0	0
2	C	9	0	0	4	0
2	D	3	0	0	1	0
2	E	1	0	0	0	0
2	F	13	0	0	1	0
2	G	2	0	0	0	0
2	H	1	0	0	0	0
All	All	19729	0	18955	1412	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 37.

The worst 5 of 1412 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:39:ARG:HE	1:A:148:ALA:C	1.39	1.27
1:A:143:ARG:HG3	1:A:143:ARG:HH11	1.06	1.14
1:A:221:GLY:HA3	2:A:401:HOH:O	1.46	1.12
1:C:213:LEU:HD21	1:C:219:LEU:HD23	1.29	1.12
1:H:23:PHE:CB	1:H:24:VAL:HA	1.73	1.12

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	310/313 (99%)	263 (85%)	33 (11%)	14 (4%)	2	14
1	B	299/313 (96%)	252 (84%)	39 (13%)	8 (3%)	5	26
1	C	310/313 (99%)	238 (77%)	52 (17%)	20 (6%)	1	7
1	D	310/313 (99%)	238 (77%)	52 (17%)	20 (6%)	1	7
1	E	310/313 (99%)	247 (80%)	41 (13%)	22 (7%)	1	5
1	F	304/313 (97%)	262 (86%)	27 (9%)	15 (5%)	2	13
1	G	301/313 (96%)	241 (80%)	39 (13%)	21 (7%)	1	6
1	H	305/313 (97%)	213 (70%)	60 (20%)	32 (10%)	0	2
All	All	2449/2504 (98%)	1954 (80%)	343 (14%)	152 (6%)	1	8

5 of 152 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	13	TRP

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	36	ALA
1	A	38	PHE
1	A	147	SER
1	A	307	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	258/260 (99%)	223 (86%)	35 (14%)	3	17
1	B	250/260 (96%)	212 (85%)	38 (15%)	3	14
1	C	256/260 (98%)	221 (86%)	35 (14%)	3	17
1	D	255/260 (98%)	217 (85%)	38 (15%)	3	14
1	E	254/260 (98%)	216 (85%)	38 (15%)	3	14
1	F	252/260 (97%)	216 (86%)	36 (14%)	3	15
1	G	250/260 (96%)	214 (86%)	36 (14%)	3	15
1	H	249/260 (96%)	209 (84%)	40 (16%)	2	12
All	All	2024/2080 (97%)	1728 (85%)	296 (15%)	3	15

5 of 296 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	G	53	CYS
1	H	235	LEU
1	G	145	LEU
1	H	31	ARG
1	C	181	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 52 such sidechains are listed below:

Mol	Chain	Res	Type
1	E	289	HIS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	F	245	HIS
1	H	280	HIS
1	E	304	GLN
1	F	136	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	312/313 (99%)	-0.57	7 (2%) 62 33	15, 25, 76, 137	0
1	B	303/313 (96%)	-0.52	5 (1%) 70 41	20, 38, 68, 147	0
1	C	312/313 (99%)	-0.55	7 (2%) 62 33	21, 36, 90, 132	0
1	D	312/313 (99%)	-0.44	8 (2%) 56 27	26, 45, 94, 153	0
1	E	312/313 (99%)	-0.46	8 (2%) 56 27	19, 41, 87, 146	0
1	F	308/313 (98%)	-0.60	6 (1%) 66 37	17, 29, 78, 139	0
1	G	305/313 (97%)	-0.42	5 (1%) 72 44	32, 49, 86, 154	0
1	H	309/313 (98%)	-0.15	10 (3%) 47 20	36, 63, 99, 164	0
All	All	2473/2504 (98%)	-0.46	56 (2%) 60 31	15, 41, 88, 164	0

The worst 5 of 56 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	H	310	THR	10.2
1	G	312	GLN	10.1
1	B	310	THR	8.5
1	F	310	THR	7.9
1	G	310	THR	7.7

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.