



wwPDB X-ray Structure Validation Summary Report ⓘ

Oct 18, 2023 – 02:26 PM EDT

PDB ID : 2FSH
Title : Complex SecA:AMP-PNP from Escherichia coli
Authors : Papanikolaou, Y.; Petratos, K.; Economou, A.
Deposited on : 2006-01-23
Resolution : 2.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	8085 (2.00-2.00)
Clashscore	141614	9178 (2.00-2.00)
Ramachandran outliers	138981	9054 (2.00-2.00)
Sidechain outliers	138945	9053 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Mol	Chain	Length	Quality of chain
1	A	853	
1	B	853	

2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 11728 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Preprotein translocase secA subunit.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	691	Total	C	N	O	S	0	0	0
			5500	3445	971	1058	26			
1	B	722	Total	C	N	O	S	0	0	0
			5735	3596	1008	1102	29			

- Molecule 2 is PHOSPHOAMINOPHOSPHONIC ACID-ADENYLATE ESTER (three-letter code: ANP) (formula: $C_{10}H_{17}N_6O_{12}P_3$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
2	A	1	Total	C	N	O	P	0	0
			31	10	6	12	3		
2	B	1	Total	C	N	O	P	0	0
			31	10	6	12	3		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	189	Total 189	O 189	0	0
3	B	242	Total 242	O 242	0	0

GLU	VAL	ILE	ILE	VAL	ASP	GLU	HIS	THR	GLY	ARG	THR	MET	GLN	GLY	ARG	ARG	TRP	SER	ASP	GLY	LEU	HIS	GLN	ALA	VAL	GLU	GLN	ILE	Q366	N367	E368	N369	Q370	T371	L372	F377	R382	E385	G392	T393	A394	D395	T396	T409	K421	E431		
K448	I457	S458	I459	E463	N477	V478	F483	H484	A485	N486	E487	I490	Y496	P497	A503	T504	N505	M506	A507	G508	R509	G510	T511	D512	S518	W519	Q520	A521	E522	V523	L526	E527	N528	A531	E535	K538	E560	R561	H562	E563	R566	I567	D568	N569	Q570			
I571	R572	G573	R574	R577	L594	M595	R596	I597	S600	V603	M607	L610	G611	M612	K613	P614	I618	E619	N629	R632	K633	V634	E635	S636	R637	I641	R642	E647	V651	R657	S661	N664	E665	L666	D671	V672	S673	I679	K685	Y691								
I682	P683	S696	D702	I703	L706	Q707	E708	D713	L718	P719	I720	W723	L724	D725	K726	E727	L730	H731	E732	L740	A741	Q742	S743	I744	E745	W746	Y747	Q748	R749	V754	E757	M758	M759	E763	K764	M767	L771	L779	A780	A781	M782	D783	Y784	L785	R786			
I789	R792	G793	Y794	A795	Q796	K797	D798	F799	K800	O801	K804	F808	S809	M810	Y820	I823	V831	R832	M833	PRO	GLU	GLU	VAL	GLU	GLU	LEU	LEU	GLN	GLN	ARG	ARG	MET	GLU	ALA	GLN	MET	GLN	GLN	LEU	ALA	GLN	MET	GLN	LEU	SER	HIS	GLN	ASP

4 Data and refinement statistics i

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	75.02Å 90.01Å 163.02Å 90.00° 100.52° 90.00°	Depositor
Resolution (Å)	19.97 – 2.00 19.97 – 2.00	Depositor EDS
% Data completeness (in resolution range)	90.5 (19.97-2.00) 90.5 (19.97-2.00)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.73 (at 2.01Å)	Xtrriage
Refinement program	REFMAC 5.2.0005	Depositor
R, R_{free}	0.204 , 0.255 0.256 , 0.296	Depositor DCC
R_{free} test set	6538 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å ²)	41.0	Xtrriage
Anisotropy	0.068	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 57.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	0.022 for h,-k,-h-l	Xtrriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	11728	wwPDB-VP
Average B, all atoms (Å ²)	54.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.86% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ANP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	1.19	15/5590 (0.3%)	1.00	15/7542 (0.2%)
1	B	1.29	33/5827 (0.6%)	1.07	17/7861 (0.2%)
All	All	1.25	48/11417 (0.4%)	1.04	32/15403 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	3
1	B	0	4
All	All	0	7

The worst 5 of 48 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	698	GLU	C-O	17.67	1.56	1.23
1	A	699	GLU	CD-OE2	14.21	1.41	1.25
1	A	98	CYS	CB-SG	-11.81	1.62	1.82
1	B	23	LYS	CE-NZ	10.56	1.75	1.49
1	A	696	SER	CB-OG	10.43	1.55	1.42

The worst 5 of 32 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	657	ARG	NE-CZ-NH2	11.13	125.87	120.30
1	A	72	ARG	NE-CZ-NH2	8.82	124.71	120.30
1	B	23	LYS	CD-CE-NZ	8.12	130.37	111.70
1	A	805	ARG	NE-CZ-NH2	-7.96	116.32	120.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	72	ARG	NE-CZ-NH1	-7.21	116.70	120.30

There are no chirality outliers.

5 of 7 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	366	GLN	Peptide
1	A	393	THR	Peptide
1	A	800	LYS	Peptide
1	B	229	ALA	Peptide
1	B	393	THR	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5500	0	5481	107	0
1	B	5735	0	5724	152	0
2	A	31	0	13	1	0
2	B	31	0	13	7	0
3	A	189	0	0	18	1
3	B	242	0	0	25	1
All	All	11728	0	11231	256	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

The worst 5 of 256 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:23:LYS:CE	1:B:23:LYS:NZ	1.75	1.49
1:B:369:ASN:C	1:B:369:ASN:HD22	1.39	1.24
1:B:757:GLU:HG3	3:B:1056:HOH:O	1.51	1.08
1:A:731:HIS:CE1	1:A:733:GLU:HB3	1.94	1.01
1:B:369:ASN:C	1:B:369:ASN:ND2	2.18	0.96

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:972:HOH:O	3:A:999:HOH:O[1_455]	2.05	0.15
3:B:1057:HOH:O	3:B:1069:HOH:O[1_655]	2.16	0.04

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	687/853 (80%)	640 (93%)	32 (5%)	15 (2%)	6 2
1	B	716/853 (84%)	661 (92%)	38 (5%)	17 (2%)	6 2
All	All	1403/1706 (82%)	1301 (93%)	70 (5%)	32 (2%)	6 2

5 of 32 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	230	GLU
1	A	368	GLU
1	A	394	ALA
1	A	532	GLU
1	A	599	ALA

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	586/728 (80%)	549 (94%)	37 (6%)	18	13
1	B	612/728 (84%)	560 (92%)	52 (8%)	10	6
All	All	1198/1456 (82%)	1109 (93%)	89 (7%)	13	9

5 of 89 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	511	THR
1	B	708	GLU
1	B	518	SER
1	B	612	MET
1	B	725	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 23 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	528	ASN
1	B	638	ASN
1	B	629	ASN
1	B	644	GLN
1	A	707	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	ANP	B	901	-	29,33,33	4.09	14 (48%)	31,52,52	3.08	14 (45%)
2	ANP	A	900	-	29,33,33	2.45	6 (20%)	31,52,52	2.14	14 (45%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	ANP	B	901	-	-	4/14/38/38	0/3/3/3
2	ANP	A	900	-	-	8/14/38/38	0/3/3/3

The worst 5 of 20 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	901	ANP	PG-O1G	14.77	1.69	1.46
2	B	901	ANP	PG-N3B	10.58	1.91	1.63
2	A	900	ANP	PG-N3B	6.77	1.81	1.63
2	A	900	ANP	PB-N3B	5.92	1.78	1.63
2	A	900	ANP	O4'-C1'	5.33	1.48	1.41

The worst 5 of 28 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	901	ANP	O1G-PG-N3B	6.82	121.81	111.77
2	B	901	ANP	O3A-PB-N3B	-6.72	87.95	106.59
2	A	900	ANP	PB-O3A-PA	-4.94	115.22	132.62
2	B	901	ANP	O2G-PG-O3G	4.77	120.35	107.64
2	B	901	ANP	O5'-C5'-C4'	4.71	125.21	108.99

There are no chirality outliers.

5 of 12 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	900	ANP	PB-N3B-PG-O1G
2	A	900	ANP	PA-O3A-PB-O1B
2	A	900	ANP	C5'-O5'-PA-O1A
2	A	900	ANP	C5'-O5'-PA-O2A
2	A	900	ANP	C5'-O5'-PA-O3A


There are no ring outliers.

2 monomers are involved in 8 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	901	ANP	7	0
2	A	900	ANP	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

Unable to reproduce the depositors R factor - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

Unable to reproduce the depositors R factor - this section is therefore empty.


6.3 Carbohydrates [i](#)


Unable to reproduce the depositors R factor - this section is therefore empty.

6.4 Ligands [i](#)

Unable to reproduce the depositors R factor - this section is therefore empty.

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





6.5 Other polymers [i](#)

Unable to reproduce the depositors R factor - this section is therefore empty.