



wwPDB X-ray Structure Validation Summary Report

Jun 13, 2024 – 10:28 AM EDT

PDB ID : 4H4L
Title : Crystal Structure of ternary complex of HutP(HutP-L-His-Zn)
Authors : Dhakshnamoorthy, B.; Misono, T.S.; Mizuno, H.; Kumar, P.K.R.
Deposited on : 2012-09-17
Resolution : 2.50 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the  symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references](#) ) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 1.20.1
EDS : 2.36.2
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4661 (2.50-2.50)
Clashscore	141614	5346 (2.50-2.50)
Ramachandran outliers	138981	5231 (2.50-2.50)
Sidechain outliers	138945	5233 (2.50-2.50)
RSRZ outliers	127900	4559 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	148	 3% 69% 26% 2%
1	B	148	 2% 68% 26% 4%
1	C	148	 0% 66% 26% 8%
1	D	148	 3% 71% 24% 2%
1	E	148	 0% 66% 28% 6%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	148	
1	G	148	
1	H	148	
1	I	148	
1	J	148	
1	K	148	
1	L	148	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	HIS	A	201	-	-	X	-
2	HIS	B	201	-	-	X	-
2	HIS	F	201	-	-	X	-
2	HIS	J	201	-	-	X	-

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 13047 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Hut operon positive regulatory protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	143	1072	676	192	200	4	1	0	0
1	B	143	1057	668	189	196	4	1	0	0
1	C	139	1062	669	191	198	4	9	0	0
1	D	142	1065	672	191	198	4	1	0	0
1	E	143	1061	670	189	198	4	1	0	0
1	F	143	1088	685	196	203	4	9	0	0
1	G	142	1065	672	191	198	4	1	0	0
1	H	143	1065	672	189	200	4	1	0	0
1	I	141	1078	679	193	202	4	9	0	0
1	J	142	1065	672	191	198	4	0	0	0
1	K	143	1061	670	189	198	4	1	0	0
1	L	142	1083	682	195	202	4	9	0	0

There are 12 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	51	ILE	VAL	engineered mutation	UNP P10943
B	51	ILE	VAL	engineered mutation	UNP P10943
C	51	ILE	VAL	engineered mutation	UNP P10943
D	51	ILE	VAL	engineered mutation	UNP P10943
E	51	ILE	VAL	engineered mutation	UNP P10943

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
F	51	ILE	VAL	engineered mutation	UNP P10943
G	51	ILE	VAL	engineered mutation	UNP P10943
H	51	ILE	VAL	engineered mutation	UNP P10943
I	51	ILE	VAL	engineered mutation	UNP P10943
J	51	ILE	VAL	engineered mutation	UNP P10943
K	51	ILE	VAL	engineered mutation	UNP P10943
L	51	ILE	VAL	engineered mutation	UNP P10943

- Molecule 2 is HISTIDINE (three-letter code: HIS) (formula: C₆H₁₀N₃O₂).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			11	6	3	2		
2	B	1	Total	C	N	O	0	0
			11	6	3	2		
2	C	1	Total	C	N	O	0	0
			11	6	3	2		
2	D	1	Total	C	N	O	0	0
			11	6	3	2		
2	D	1	Total	C	N	O	0	0
			11	6	3	2		
2	F	1	Total	C	N	O	0	0
			11	6	3	2		
2	G	1	Total	C	N	O	0	0
			11	6	3	2		
2	G	1	Total	C	N	O	0	0
			11	6	3	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	I	1	Total	C	N	O	0	0
			11	6	3	2		
2	J	1	Total	C	N	O	0	0
			11	6	3	2		
2	J	1	Total	C	N	O	0	0
			11	6	3	2		
2	L	1	Total	C	N	O	0	0
			11	6	3	2		

- Molecule 3 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Zn	0	0
			1	1		
3	B	1	Total	Zn	0	0
			1	1		
3	C	1	Total	Zn	0	0
			1	1		
3	D	1	Total	Zn	0	0
			1	1		
3	E	1	Total	Zn	0	0
			1	1		
3	F	1	Total	Zn	0	0
			1	1		
3	G	1	Total	Zn	0	0
			1	1		
3	H	1	Total	Zn	0	0
			1	1		
3	I	1	Total	Zn	0	0
			1	1		
3	J	1	Total	Zn	0	0
			1	1		
3	K	1	Total	Zn	0	0
			1	1		
3	L	1	Total	Zn	0	0
			1	1		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	10	Total	O	0	0
			10	10		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	B	8	Total O 8 8	0	0
4	C	10	Total O 10 10	0	0
4	D	6	Total O 6 6	0	0
4	E	6	Total O 6 6	0	0
4	F	7	Total O 7 7	0	0
4	G	6	Total O 6 6	0	0
4	H	9	Total O 9 9	0	0
4	I	5	Total O 5 5	0	0
4	J	5	Total O 5 5	0	0
4	K	3	Total O 3 3	0	0
4	L	6	Total O 6 6	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Hut operon positive regulatory protein



- Molecule 1: Hut operon positive regulatory protein



- Molecule 1: Hut operon positive regulatory protein




- Molecule 1: Hut operon positive regulatory protein








- Molecule 1: Hut operon positive regulatory protein




- Molecule 1: Hut operon positive regulatory protein




- Molecule 1: Hut operon positive regulatory protein




- Molecule 1: Hut operon positive regulatory protein




- Molecule 1: Hut operon positive regulatory protein




- Molecule 1: Hut operon positive regulatory protein



- Molecule 1: Hut operon positive regulatory protein



- Molecule 1: Hut operon positive regulatory protein



4 Data and refinement statistics

Property	Value	Source
Space group	P 3	Depositor
Cell constants a, b, c, α , β , γ	129.53Å 129.53Å 76.67Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	36.28 – 2.50 32.99 – 2.50	Depositor EDS
% Data completeness (in resolution range)	98.1 (36.28-2.50) 98.1 (32.99-2.50)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.40 (at 2.51Å)	Xtrriage
Refinement program	REFMAC 5.6.0117	Depositor
R, R_{free}	0.219 , 0.298 0.220 , 0.298	Depositor DCC
R_{free} test set	2507 reflections (5.13%)	wwPDB-VP
Wilson B-factor (Å ²)	51.1	Xtrriage
Anisotropy	0.072	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 24.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	0.478 for -h,-k,l 0.054 for h,-h-k,-l 0.054 for -k,-h,-l	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	13047	wwPDB-VP
Average B, all atoms (Å ²)	58.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 53.79 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 4.0059e-05. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section:
ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.68	2/1089 (0.2%)	0.79	0/1471
1	B	0.66	1/1073 (0.1%)	0.79	0/1450
1	C	0.73	1/1079 (0.1%)	0.80	0/1453
1	D	0.68	2/1082 (0.2%)	0.82	0/1461
1	E	0.64	0/1077	0.81	1/1455 (0.1%)
1	F	0.73	1/1105 (0.1%)	0.82	0/1489
1	G	0.67	0/1082	0.83	0/1461
1	H	0.64	1/1081 (0.1%)	0.85	0/1460
1	I	0.68	1/1095 (0.1%)	0.79	1/1475 (0.1%)
1	J	0.67	0/1082	0.82	0/1461
1	K	0.66	1/1077 (0.1%)	0.82	0/1455
1	L	0.75	1/1100 (0.1%)	0.89	3/1482 (0.2%)
All	All	0.68	11/13022 (0.1%)	0.82	5/17573 (0.0%)

The worst 5 of 11 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	L	120	TRP	CD2-CE2	5.76	1.48	1.41
1	C	134	LYS	CB-CG	-5.65	1.37	1.52
1	K	35	TRP	CD2-CE2	5.26	1.47	1.41
1	F	120	TRP	CD2-CE2	5.20	1.47	1.41
1	D	120	TRP	CD2-CE2	5.20	1.47	1.41

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	I	141	PHE	CB-CA-C	-5.63	99.14	110.40
1	E	66	SER	CA-CB-OG	5.44	125.90	111.20
1	L	39	LEU	CB-CG-CD1	-5.32	101.96	111.00
1	L	30	LEU	N-CA-C	-5.06	97.33	111.00
1	L	134	LYS	N-CA-CB	5.02	119.64	110.60

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1072	0	1058	30	0
1	B	1057	0	1041	38	0
1	C	1062	0	1062	29	0
1	D	1065	0	1051	27	0
1	E	1061	0	1045	36	0
1	F	1088	0	1086	26	0
1	G	1065	0	1051	55	0
1	H	1065	0	1049	44	0
1	I	1078	0	1077	36	0
1	J	1065	0	1051	56	0
1	K	1061	0	1045	32	0
1	L	1083	0	1081	36	0
2	A	11	0	6	8	0
2	B	11	0	6	9	0
2	C	11	0	6	4	0
2	D	22	0	12	5	0
2	F	11	0	6	6	0
2	G	22	0	12	5	0
2	I	11	0	6	4	0
2	J	22	0	12	12	0
2	L	11	0	6	4	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
3	C	1	0	0	0	0
3	D	1	0	0	0	0
3	E	1	0	0	0	0
3	F	1	0	0	0	0
3	G	1	0	0	0	0
3	H	1	0	0	0	0
3	I	1	0	0	0	0
3	J	1	0	0	0	0
3	K	1	0	0	0	0
3	L	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	A	10	0	0	1	0
4	B	8	0	0	0	0
4	C	10	0	0	0	0
4	D	6	0	0	0	0
4	E	6	0	0	2	0
4	F	7	0	0	0	0
4	G	6	0	0	3	0
4	H	9	0	0	1	0
4	I	5	0	0	0	0
4	J	5	0	0	2	0
4	K	3	0	0	0	0
4	L	6	0	0	2	0
All	All	13047	0	12769	393	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 15.

The worst 5 of 393 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:I:130:GLY:HA3	1:I:137:GLU:HB3	1.21	1.14
1:C:29:GLU:HG2	1:C:30:LEU:H	0.97	1.10
1:J:63:VAL:HG12	1:J:64:ILE:HD12	1.25	1.09
1:G:98:ARG:HD2	1:G:129:ILE:HD11	1.36	1.06
1:D:26:GLN:HG3	1:D:27:VAL:H	1.19	1.00

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	139/148 (94%)	129 (93%)	10 (7%)	0	100	100
1	B	139/148 (94%)	128 (92%)	11 (8%)	0	100	100
1	C	135/148 (91%)	123 (91%)	11 (8%)	1 (1%)	22	39
1	D	138/148 (93%)	127 (92%)	11 (8%)	0	100	100
1	E	139/148 (94%)	128 (92%)	11 (8%)	0	100	100
1	F	139/148 (94%)	130 (94%)	9 (6%)	0	100	100
1	G	138/148 (93%)	123 (89%)	15 (11%)	0	100	100
1	H	139/148 (94%)	127 (91%)	12 (9%)	0	100	100
1	I	137/148 (93%)	123 (90%)	12 (9%)	2 (2%)	10	18
1	J	138/148 (93%)	125 (91%)	13 (9%)	0	100	100
1	K	139/148 (94%)	125 (90%)	13 (9%)	1 (1%)	22	39
1	L	138/148 (93%)	126 (91%)	11 (8%)	1 (1%)	22	39
All	All	1658/1776 (93%)	1514 (91%)	139 (8%)	5 (0%)	41	61

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	I	29	GLU
1	L	29	GLU
1	I	95	SER
1	C	132	PRO
1	K	62	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	108/119 (91%)	101 (94%)	7 (6%)	17	33
1	B	104/119 (87%)	97 (93%)	7 (7%)	16	31
1	C	109/119 (92%)	95 (87%)	14 (13%)	4	8
1	D	107/119 (90%)	101 (94%)	6 (6%)	21	40

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	E	105/119 (88%)	95 (90%)	10 (10%)	8	17
1	F	111/119 (93%)	101 (91%)	10 (9%)	9	19
1	G	107/119 (90%)	94 (88%)	13 (12%)	5	9
1	H	106/119 (89%)	91 (86%)	15 (14%)	3	6
1	I	111/119 (93%)	98 (88%)	13 (12%)	5	10
1	J	107/119 (90%)	88 (82%)	19 (18%)	2	3
1	K	105/119 (88%)	91 (87%)	14 (13%)	4	7
1	L	111/119 (93%)	99 (89%)	12 (11%)	6	12
All	All	1291/1428 (90%)	1151 (89%)	140 (11%)	6	12

5 of 140 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	K	3	LEU
1	K	45	MET
1	L	20	GLU
1	F	32	ARG
1	F	27	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 27 such sidechains are listed below:

Mol	Chain	Res	Type
1	G	4	HIS
1	G	65	GLN
1	J	84	HIS
1	G	48	HIS
1	G	84	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 24 ligands modelled in this entry, 12 are monoatomic - leaving 12 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	HIS	J	203	3	6,11,11	1.00	1 (16%)	5,14,14	1.63	2 (40%)
2	HIS	A	201	3	6,11,11	0.86	0	5,14,14	1.55	1 (20%)
2	HIS	G	201	3	6,11,11	1.03	0	5,14,14	1.53	1 (20%)
2	HIS	I	201	3	6,11,11	0.94	1 (16%)	5,14,14	1.70	2 (40%)
2	HIS	L	201	3	6,11,11	1.05	1 (16%)	5,14,14	1.72	3 (60%)
2	HIS	C	201	3	6,11,11	0.90	1 (16%)	5,14,14	1.70	2 (40%)
2	HIS	D	203	3	6,11,11	1.12	1 (16%)	5,14,14	1.74	2 (40%)
2	HIS	D	201	3	6,11,11	0.89	0	5,14,14	1.47	1 (20%)
2	HIS	F	201	3	6,11,11	1.09	1 (16%)	5,14,14	1.67	2 (40%)
2	HIS	G	203	3	6,11,11	0.98	1 (16%)	5,14,14	1.60	2 (40%)
2	HIS	J	201	3	6,11,11	0.99	1 (16%)	5,14,14	1.74	2 (40%)
2	HIS	B	201	3	6,11,11	0.96	1 (16%)	5,14,14	1.58	2 (40%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	HIS	J	203	3	-	3/8/8/8	0/1/1/1
2	HIS	A	201	3	-	6/8/8/8	0/1/1/1
2	HIS	G	201	3	-	6/8/8/8	0/1/1/1
2	HIS	I	201	3	-	2/8/8/8	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	HIS	L	201	3	-	2/8/8/8	0/1/1/1
2	HIS	C	201	3	-	6/8/8/8	0/1/1/1
2	HIS	D	203	3	-	6/8/8/8	0/1/1/1
2	HIS	D	201	3	-	6/8/8/8	0/1/1/1
2	HIS	F	201	3	-	4/8/8/8	0/1/1/1
2	HIS	G	203	3	-	6/8/8/8	0/1/1/1
2	HIS	J	201	3	-	6/8/8/8	0/1/1/1
2	HIS	B	201	3	-	4/8/8/8	0/1/1/1

The worst 5 of 9 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	203	HIS	OXT-C	-2.65	1.22	1.30
2	F	201	HIS	OXT-C	-2.44	1.22	1.30
2	J	203	HIS	OXT-C	-2.37	1.23	1.30
2	J	201	HIS	OXT-C	-2.34	1.23	1.30
2	L	201	HIS	OXT-C	-2.21	1.23	1.30

The worst 5 of 22 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	201	HIS	CD2-NE2-CE1	2.95	110.38	105.72
2	D	201	HIS	CD2-NE2-CE1	2.86	110.25	105.72
2	A	201	HIS	CD2-NE2-CE1	2.83	110.20	105.72
2	D	203	HIS	CD2-NE2-CE1	2.71	110.00	105.72
2	F	201	HIS	CD2-NE2-CE1	2.65	109.91	105.72

There are no chirality outliers.

5 of 57 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	201	HIS	CA-CB-CG-ND1
2	A	201	HIS	CA-CB-CG-CD2
2	B	201	HIS	N-CA-CB-CG
2	B	201	HIS	C-CA-CB-CG
2	C	201	HIS	CA-CB-CG-ND1

There are no ring outliers.

12 monomers are involved in 57 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	J	203	HIS	5	0
2	A	201	HIS	8	0
2	G	201	HIS	3	0
2	I	201	HIS	4	0
2	L	201	HIS	4	0
2	C	201	HIS	4	0
2	D	203	HIS	2	0
2	D	201	HIS	3	0
2	F	201	HIS	6	0
2	G	203	HIS	2	0
2	J	201	HIS	7	0
2	B	201	HIS	9	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	143/148 (96%)	-0.20	5 (3%) 44 47	32, 53, 97, 135	1 (0%)
1	B	143/148 (96%)	-0.23	3 (2%) 63 66	33, 54, 89, 110	1 (0%)
1	C	139/148 (93%)	-0.40	2 (1%) 75 77	30, 44, 74, 108	2 (1%)
1	D	142/148 (95%)	-0.20	4 (2%) 53 56	31, 54, 93, 130	1 (0%)
1	E	143/148 (96%)	-0.23	2 (1%) 75 77	33, 56, 90, 108	1 (0%)
1	F	143/148 (96%)	-0.37	3 (2%) 63 66	26, 44, 76, 105	2 (1%)
1	G	142/148 (95%)	0.08	7 (4%) 29 31	39, 63, 95, 118	1 (0%)
1	H	143/148 (96%)	0.13	9 (6%) 20 21	39, 59, 101, 145	1 (0%)
1	I	141/148 (95%)	-0.15	4 (2%) 53 56	34, 51, 95, 133	2 (1%)
1	J	142/148 (95%)	0.20	6 (4%) 36 39	35, 63, 92, 120	0
1	K	143/148 (96%)	0.33	9 (6%) 20 21	37, 59, 101, 157	1 (0%)
1	L	142/148 (95%)	-0.12	5 (3%) 44 47	29, 51, 100, 123	2 (1%)
All	All	1706/1776 (96%)	-0.10	59 (3%) 44 47	26, 55, 96, 157	15 (0%)

The worst 5 of 59 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	K	4	HIS	19.2
1	K	2	THR	11.0
1	G	2	THR	8.1
1	H	4	HIS	8.1
1	L	2	THR	7.7

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	HIS	I	201	11/11	0.90	0.23	42,55,60,62	0
2	HIS	J	203	11/11	0.90	0.23	43,46,52,52	0
2	HIS	A	201	11/11	0.91	0.20	41,46,51,52	0
2	HIS	F	201	11/11	0.91	0.17	38,39,44,50	0
2	HIS	C	201	11/11	0.92	0.20	41,48,51,51	0
2	HIS	D	201	11/11	0.92	0.17	41,47,59,63	0
2	HIS	D	203	11/11	0.92	0.18	41,45,49,49	0
2	HIS	J	201	11/11	0.93	0.16	68,72,84,88	0
2	HIS	G	203	11/11	0.93	0.18	42,44,48,53	0
2	HIS	B	201	11/11	0.94	0.15	36,39,47,50	0
2	HIS	L	201	11/11	0.94	0.17	43,54,60,62	0
2	HIS	G	201	11/11	0.95	0.14	76,80,85,92	0
3	ZN	C	202	1/1	0.95	0.12	48,48,48,48	0
3	ZN	A	202	1/1	0.96	0.15	47,47,47,47	0
3	ZN	I	202	1/1	0.96	0.15	56,56,56,56	0
3	ZN	D	202	1/1	0.97	0.12	46,46,46,46	0
3	ZN	B	202	1/1	0.97	0.14	50,50,50,50	0
3	ZN	F	202	1/1	0.98	0.12	43,43,43,43	0
3	ZN	E	201	1/1	0.98	0.15	48,48,48,48	0
3	ZN	L	202	1/1	0.98	0.14	54,54,54,54	0
3	ZN	J	202	1/1	0.99	0.11	60,60,60,60	0
3	ZN	K	201	1/1	0.99	0.12	46,46,46,46	0
3	ZN	H	201	1/1	0.99	0.14	46,46,46,46	0
3	ZN	G	202	1/1	1.00	0.12	61,61,61,61	0

6.5 Other polymers [i](#)

There are no such residues in this entry.