



# wwPDB X-ray Structure Validation Summary Report ⓘ

Jun 12, 2024 – 11:15 PM EDT

PDB ID : 3HO8  
Title : Crystal Structure of S. aureus Pyruvate Carboxylase in complex with Coenzyme A  
Authors : Tong, L.; Yu, L.P.C.  
Deposited on : 2009-06-01  
Resolution : 2.90 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	1.20.1
EDS	:	2.36.2
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.36.2


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1957 (2.90-2.90)
Clashscore	141614	2172 (2.90-2.90)
Ramachandran outliers	138981	2115 (2.90-2.90)
Sidechain outliers	138945	2117 (2.90-2.90)
RSRZ outliers	127900	1906 (2.90-2.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	1150	<div> <div>4%</div> <div>57%</div> <div>26%</div> <div>•</div> <div>14%</div> </div>
1	B	1150	<div> <div>%</div> <div>52%</div> <div>29%</div> <div>5%</div> <div>14%</div> </div>
1	C	1150	<div> <div>2%</div> <div>56%</div> <div>26%</div> <div>•</div> <div>13%</div> </div>
1	D	1150	<div> <div>6%</div> <div>54%</div> <div>23%</div> <div>•</div> <div>19%</div> </div>

## 2 Entry composition i


There are 4 unique types of molecules in this entry. The entry contains 31229 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Pyruvate carboxylase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	994	Total	C	N	O	S	0	0	0
			7880	5000	1329	1525	26			
1	D	934	Total	C	N	O	S	0	0	0
			7396	4696	1250	1426	24			
1	C	995	Total	C	N	O	S	0	0	0
			7889	5005	1330	1528	26			
1	B	989	Total	C	N	O	S	0	0	0
			7838	4975	1321	1516	26			

- Molecule 2 is COENZYME A (three-letter code: COA) (formula: C<sub>21</sub>H<sub>36</sub>N<sub>7</sub>O<sub>16</sub>P<sub>3</sub>S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P S	0	0
			48	21	7	16	3 1		
2	D	1	Total	C	N	O	P S	0	0
			48	21	7	16	3 1		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
2	C	1	Total	C	N	O	P	S	0	0
			48	21	7	16	3	1		
2	B	1	Total	C	N	O	P	S	0	0
			48	21	7	16	3	1		

- Molecule 3 is MANGANESE (II) ION (three-letter code: MN) (formula: Mn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Mn	0	0
			1	1		
3	D	1	Total	Mn	0	0
			1	1		
3	C	1	Total	Mn	0	0
			1	1		
3	B	1	Total	Mn	0	0
			1	1		

- Molecule 4 is 5-(HEXAHYDRO-2-OXO-1H-THIENO[3,4-D]IMIDAZOL-6-YL)PENTANAL (three-letter code: BTI) (formula: C<sub>10</sub>H<sub>16</sub>N<sub>2</sub>O<sub>2</sub>S).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	C	1	Total	C	N	O	S	0	0
			15	10	2	2	1		
4	B	1	Total	C	N	O	S	0	0
			15	10	2	2	1		


### 3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


#### • Molecule 1: Pyruvate carboxylase








### • Molecule 1: Pyruvate carboxylase



- Molecule 1: Pyruvate carboxylase





M1042	R1043	N1044	E1049	D1053	I1059	I1060	I1065	E1070	N1071	I1076	Y1077	M1080	N1081	R1086	I1089	E1092	N1093	VAL	HIS	THR	ASN	ALA	ASN	ASN	VAL	LYS	PRO	LYS	ALA	ASP	LYS	SER	THR	ASN	PRO	SER	SER	HIS	ILE	GLY	LEU	ALA	GLN	ILE	GLU	ILE	PRO	GLY	LYS	SER	ALA	VAL	THR	THR	GLU	VAL	LYS	VAL
SER	VAL	GLY	THR	VAL	LYS	ALA	ASN	GLN	PRO	LEU	LEU	ILE	THR	GLU	ALA	MET	LYS	MET	GLU	THR	THR	ILE	GLN	ALA	PRO	PHE	ASP	GLY	VAL	ILE	LYS	GLN	VAL	THR	VAL	ASN	ASN	GLY	ASP	LYS	THR	ILE	ALA	THR	GLY	ASP	ILE	LEU	LEU	ALA	ALA	THR	THR	ASP				

## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	96.59Å 164.47Å 373.35Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	29.97 – 2.90 29.98 – 2.90	Depositor EDS
% Data completeness (in resolution range)	89.8 (29.97-2.90) 89.8 (29.98-2.90)	Depositor EDS
$R_{merge}$	0.14	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.03 (at 2.90Å)	Xtriage
Refinement program	REFMAC 5.5.0072	Depositor
R, $R_{free}$	0.264 , 0.328 0.266 , 0.322	Depositor DCC
$R_{free}$ test set	5969 reflections (5.04%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	64.2	Xtriage
Anisotropy	0.098	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.30 , 42.5	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.88	EDS
Total number of atoms	31229	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	67.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 22.90 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 5.1935e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MN, BTI, COA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.43	0/8033	0.60	1/10868 (0.0%)
1	B	0.49	0/7990	0.64	0/10810
1	C	0.50	0/8042	0.65	2/10880 (0.0%)
1	D	0.43	0/7537	0.60	0/10192
All	All	0.47	0/31602	0.62	3/42750 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	C	0	1

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	526	LEU	CA-CB-CG	5.56	128.09	115.30
1	C	828	ILE	CB-CA-C	-5.22	101.16	111.60
1	C	526	LEU	CA-CB-CG	5.00	126.81	115.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	C	880	SER	Peptide

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	7880	0	7807	199	0
1	B	7838	0	7771	292	0
1	C	7889	0	7813	232	0
1	D	7396	0	7347	158	0
2	A	48	0	32	0	0
2	B	48	0	32	1	0
2	C	48	0	32	2	0
2	D	48	0	32	6	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
3	C	1	0	0	0	0
3	D	1	0	0	0	0
4	B	15	0	16	0	0
4	C	15	0	16	3	0
All	All	31229	0	30898	870	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 14.

The worst 5 of 870 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:504:ILE:HG21	1:C:1042:MET:HE2	1.18	1.12
1:C:704:ILE:HG12	1:C:726:LEU:HD23	1.22	1.12
1:A:304:TYR:HE2	1:A:307:ALA:O	1.35	1.10
1:A:1085:ARG:HG2	1:A:1085:ARG:HH11	1.12	1.06
1:C:743:MET:HG3	1:C:907:VAL:HG13	1.38	1.04

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	990/1150 (86%)	862 (87%)	105 (11%)	23 (2%)	6	23
1	B	985/1150 (86%)	875 (89%)	89 (9%)	21 (2%)	7	26
1	C	991/1150 (86%)	887 (90%)	79 (8%)	25 (2%)	5	21
1	D	924/1150 (80%)	777 (84%)	123 (13%)	24 (3%)	5	20
All	All	3890/4600 (85%)	3401 (87%)	396 (10%)	93 (2%)	6	22

5 of 93 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	868	GLY
1	A	939	ASP
1	D	87	GLY
1	D	163	ASP
1	D	272	GLN

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	860/987 (87%)	783 (91%)	77 (9%)	9	29
1	B	856/987 (87%)	762 (89%)	94 (11%)	6	19
1	C	861/987 (87%)	759 (88%)	102 (12%)	5	16
1	D	806/987 (82%)	711 (88%)	95 (12%)	5	16
All	All	3383/3948 (86%)	3015 (89%)	368 (11%)	6	19

5 of 368 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	671	ILE
1	B	250	ILE
1	C	781	LEU
1	C	991	ARG
1	B	426	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 122 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	330	GLN
1	B	873	ASN
1	C	870	GLN
1	B	864	HIS
1	B	1025	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 10 ligands modelled in this entry, 4 are monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	COA	C	2001	-	43,50,50	1.08	1 (2%)	56,75,75	1.57	6 (10%)
4	BTI	B	2000	-	15,16,16	1.62	2 (13%)	20,21,21	1.95	3 (15%)
2	COA	A	2001	-	43,50,50	1.39	3 (6%)	56,75,75	1.67	7 (12%)
2	COA	B	2001	-	43,50,50	1.26	4 (9%)	56,75,75	1.72	8 (14%)
4	BTI	C	2000	-	15,16,16	1.70	2 (13%)	20,21,21	2.23	3 (15%)
2	COA	D	2001	-	43,50,50	1.07	1 (2%)	56,75,75	1.64	5 (8%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	COA	C	2001	-	-	14/44/64/64	0/3/3/3
4	BTI	B	2000	-	-	3/6/27/27	0/2/2/2
2	COA	A	2001	-	-	16/44/64/64	0/3/3/3
2	COA	B	2001	-	-	17/44/64/64	0/3/3/3
4	BTI	C	2000	-	-	5/6/27/27	0/2/2/2
2	COA	D	2001	-	-	4/44/64/64	0/3/3/3

The worst 5 of 13 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	2001	COA	P2A-O3A	5.00	1.64	1.59
4	C	2000	BTI	O3-C3	4.72	1.33	1.23
4	B	2000	BTI	O3-C3	4.41	1.32	1.23
2	A	2001	COA	P1A-O3A	4.10	1.63	1.59
4	C	2000	BTI	C2-S1	-3.68	1.76	1.82

The worst 5 of 32 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	2001	COA	O4B-C1B-N9A	-7.91	98.26	108.75
4	B	2000	BTI	C6-C5-N3	-6.36	104.99	113.18
4	C	2000	BTI	C2-C4-N2	-6.31	106.66	113.34
4	C	2000	BTI	C6-C5-N3	-6.26	105.12	113.18
2	A	2001	COA	O4B-C1B-N9A	-6.19	100.54	108.75

There are no chirality outliers.

5 of 59 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	2001	COA	CCP-O6A-P2A-O3A
2	A	2001	COA	CCP-O6A-P2A-O4A
2	A	2001	COA	CCP-O6A-P2A-O5A
2	A	2001	COA	C9P-CAP-CBP-CCP
2	D	2001	COA	S1P-C2P-C3P-N4P



There are no ring outliers.


4 monomers are involved in 12 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	C	2001	COA	2	0
2	B	2001	COA	1	0
4	C	2000	BTI	3	0
2	D	2001	COA	6	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å <sup>2</sup> )	Q<0.9
1	A	994/1150 (86%)	0.21	48 (4%)	30	27	39, 76, 117, 151	0
1	B	989/1150 (86%)	-0.20	12 (1%)	79	79	26, 52, 83, 102	0
1	C	995/1150 (86%)	-0.17	19 (1%)	66	65	29, 52, 83, 107	0
1	D	934/1150 (81%)	0.41	64 (6%)	16	13	41, 83, 128, 173	0
All	All	3912/4600 (85%)	0.06	143 (3%)	41	37	26, 64, 111, 173	0

The worst 5 of 143 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	962	ASP	9.6
1	D	965	ALA	6.8
1	D	981	TYR	6.4
1	D	975	THR	5.7
1	D	966	VAL	5.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	MN	A	2002	1/1	0.91	0.08	86,86,86,86	0
2	COA	D	2001	48/48	0.94	0.15	45,51,55,56	0
3	MN	B	2002	1/1	0.94	0.08	60,60,60,60	0
4	BTI	B	2000	15/15	0.94	0.13	47,53,54,55	0
4	BTI	C	2000	15/15	0.95	0.12	36,40,42,43	0
3	MN	C	2002	1/1	0.96	0.07	65,65,65,65	0
2	COA	B	2001	48/48	0.96	0.14	37,40,43,45	0
2	COA	C	2001	48/48	0.97	0.12	32,36,48,51	0
3	MN	D	2002	1/1	0.97	0.10	65,65,65,65	0
2	COA	A	2001	48/48	0.97	0.13	44,48,57,59	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




**Electron density around COA D 2001:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around COA B 2001:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



**Electron density around COA C 2001:**




$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





**Electron density around COA A 2001:**

2mF<sub>o</sub>-DF<sub>c</sub> (at 0.7 rmsd) in gray  
mF<sub>o</sub>-DF<sub>c</sub> (at 3 rmsd) in purple (negative)  
and green (positive)



## 6.5 Other polymers [i](#)

There are no such residues in this entry.