



wwPDB X-ray Structure Validation Summary Report ⓘ

Jun 18, 2024 – 09:15 PM EDT

PDB ID : 4JW1
Title : Crystal structure of N-terminal 618-residue fragment of LepB from Legionella pneumophila
Authors : Hu, L.; Yao, Q.; Zhu, Y.; Shao, F.
Deposited on : 2013-03-26
Resolution : 3.16 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	1.20.1
EDS	:	2.37.1
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.37.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.16 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1665 (3.20-3.12)
Clashscore	141614	1804 (3.20-3.12)
Ramachandran outliers	138981	1770 (3.20-3.12)
Sidechain outliers	138945	1769 (3.20-3.12)
RSRZ outliers	127900	1616 (3.20-3.12)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	626	
1	B	626	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	FLC	A	702	-	X	-	-

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 6745 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Effector protein B.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	515	Total	C	N	O	S	Se	0	1	0
			4130	2679	676	758	7	10			
1	B	319	Total	C	N	O	S	Se	0	0	0
			2569	1670	426	467	4	2			

There are 22 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-7	HIS	-	EXPRESSION TAG	UNP Q6X1Y7
A	-6	HIS	-	EXPRESSION TAG	UNP Q6X1Y7
A	-5	HIS	-	EXPRESSION TAG	UNP Q6X1Y7
A	-4	HIS	-	EXPRESSION TAG	UNP Q6X1Y7
A	-3	HIS	-	EXPRESSION TAG	UNP Q6X1Y7
A	-2	HIS	-	EXPRESSION TAG	UNP Q6X1Y7
A	-1	GLY	-	EXPRESSION TAG	UNP Q6X1Y7
A	0	PRO	-	EXPRESSION TAG	UNP Q6X1Y7
A	13	ALA	LYS	CONFLICT	UNP Q6X1Y7
A	14	ALA	GLU	CONFLICT	UNP Q6X1Y7
A	15	ALA	LYS	CONFLICT	UNP Q6X1Y7
B	-7	HIS	-	EXPRESSION TAG	UNP Q6X1Y7
B	-6	HIS	-	EXPRESSION TAG	UNP Q6X1Y7
B	-5	HIS	-	EXPRESSION TAG	UNP Q6X1Y7
B	-4	HIS	-	EXPRESSION TAG	UNP Q6X1Y7
B	-3	HIS	-	EXPRESSION TAG	UNP Q6X1Y7
B	-2	HIS	-	EXPRESSION TAG	UNP Q6X1Y7
B	-1	GLY	-	EXPRESSION TAG	UNP Q6X1Y7
B	0	PRO	-	EXPRESSION TAG	UNP Q6X1Y7
B	13	ALA	LYS	CONFLICT	UNP Q6X1Y7
B	14	ALA	GLU	CONFLICT	UNP Q6X1Y7
B	15	ALA	LYS	CONFLICT	UNP Q6X1Y7

- Molecule 2 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			6	3	3		
2	B	1	Total	C	O	0	0
			6	3	3		
2	B	1	Total	C	O	0	0
			6	3	3		
2	B	1	Total	C	O	0	0
			6	3	3		
2	B	1	Total	C	O	0	0
			6	3	3		


- Molecule 3 is CITRATE ANION (three-letter code: FLC) (formula: $C_6H_5O_7$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			13	6	7		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	2	Total	O	0	0
			2	2		
4	B	1	Total	O	0	0
			1	1		



4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	101.54Å 159.42Å 181.15Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	19.95 – 3.16 19.95 – 3.16	Depositor EDS
% Data completeness (in resolution range)	99.5 (19.95-3.16) 99.5 (19.95-3.16)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.94 (at 3.15Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.8_1069)	Depositor
R, R_{free}	0.264 , 0.283 0.251 , 0.290	Depositor DCC
R_{free} test set	1269 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å ²)	35.4	Xtriage
Anisotropy	0.155	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.24 , 47.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.83	EDS
Total number of atoms	6745	wwPDB-VP
Average B, all atoms (Å ²)	43.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.40% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: FLC, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.61	8/4207 (0.2%)	0.61	4/5644 (0.1%)
1	B	0.62	1/2615 (0.0%)	0.60	0/3512
All	All	0.62	9/6822 (0.1%)	0.60	4/9156 (0.0%)

The worst 5 of 9 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	183	TYR	CD2-CE2	-7.53	1.28	1.39
1	A	183	TYR	CD1-CE1	-7.23	1.28	1.39
1	A	159	VAL	CB-CG2	-7.20	1.37	1.52
1	B	394	GLU	CB-CG	-5.91	1.41	1.52
1	A	183	TYR	CE2-CZ	-5.82	1.30	1.38

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	174	ARG	NE-CZ-NH1	6.86	123.73	120.30
1	A	174	ARG	NE-CZ-NH2	-6.36	117.12	120.30
1	A	167	GLU	OE1-CD-OE2	5.94	130.43	123.30
1	A	546	LEU	CB-CG-CD2	-5.28	102.03	111.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within

the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4130	0	4151	160	0
1	B	2569	0	2573	73	0
2	A	6	0	8	0	0
2	B	24	0	32	0	0
3	A	13	0	5	2	0
4	A	2	0	0	0	0
4	B	1	0	0	0	0
All	All	6745	0	6769	234	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 17.

The worst 5 of 234 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:168:LYS:NZ	1:B:172:PHE:CE2	1.88	1.34
1:A:431:ARG:CG	1:A:431:ARG:HH21	1.67	1.06
1:A:161:LEU:HD12	1:A:165:GLU:HG2	1.07	1.06
1:A:98:GLU:HA	1:A:159:VAL:HG12	1.38	1.05
1:A:431:ARG:HH21	1:A:431:ARG:HG2	1.20	1.05

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	498/626 (80%)	465 (93%)	26 (5%)	7 (1%)	11	43
1	B	293/626 (47%)	267 (91%)	22 (8%)	4 (1%)	11	43
All	All	791/1252 (63%)	732 (92%)	48 (6%)	11 (1%)	11	43

5 of 11 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	85	LYS
1	A	187	PRO
1	A	390	VAL
1	B	251	LYS
1	A	211	ASP

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	446/526 (85%)	389 (87%)	57 (13%)	4	18
1	B	278/526 (53%)	236 (85%)	42 (15%)	3	13
All	All	724/1052 (69%)	625 (86%)	99 (14%)	3	16

5 of 99 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	49	THR
1	B	254	PRO
1	B	52	PHE
1	B	170	LYS
1	B	340	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	265	GLN
1	B	157	ASN
1	B	479	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	GOL	B	702	-	5,5,5	0.36	0	5,5,5	0.25	0
3	FLC	A	702	-	12,12,12	3.67	7 (58%)	17,17,17	2.72	6 (35%)
2	GOL	B	704	-	5,5,5	0.37	0	5,5,5	0.30	0
2	GOL	B	701	-	5,5,5	0.35	0	5,5,5	0.38	0
2	GOL	B	703	-	5,5,5	0.37	0	5,5,5	0.31	0
2	GOL	A	701	-	5,5,5	0.38	0	5,5,5	0.27	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GOL	B	702	-	-	2/4/4/4	-
3	FLC	A	702	-	-	8/16/16/16	-
2	GOL	B	704	-	-	3/4/4/4	-
2	GOL	B	701	-	-	1/4/4/4	-
2	GOL	B	703	-	-	2/4/4/4	-
2	GOL	A	701	-	-	4/4/4/4	-

The worst 5 of 7 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	702	FLC	CB-CBC	6.20	1.59	1.53
3	A	702	FLC	CG-CB	-5.99	1.46	1.54
3	A	702	FLC	CA-CB	5.69	1.61	1.54
3	A	702	FLC	OB1-CBC	4.30	1.35	1.22
3	A	702	FLC	OHB-CB	3.62	1.50	1.43

The worst 5 of 6 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	702	FLC	OHB-CB-CBC	7.79	120.01	108.96
3	A	702	FLC	CG-CB-CBC	-4.52	100.05	110.03
3	A	702	FLC	CB-CG-CGC	-2.80	106.28	113.92
3	A	702	FLC	OG2-CGC-CG	2.74	123.02	114.35
3	A	702	FLC	OG1-CGC-CG	-2.43	116.07	122.95

There are no chirality outliers.

5 of 20 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	701	GOL	O1-C1-C2-C3
2	B	702	GOL	O1-C1-C2-C3
2	B	704	GOL	O1-C1-C2-C3
3	A	702	FLC	CAC-CA-CB-CBC
3	A	702	FLC	CBC-CB-CG-CGC

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	702	FLC	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	505/626 (80%)	-0.33	8 (1%) 72 59	9, 37, 71, 104	0
1	B	317/626 (50%)	-0.23	4 (1%) 77 66	15, 47, 80, 92	0
All	All	822/1252 (65%)	-0.29	12 (1%) 73 61	9, 41, 75, 104	0

The worst 5 of 12 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	391	TRP	3.6
1	A	31	ASN	3.0
1	B	335	SER	3.0
1	A	30	ASP	2.7
1	B	392	ALA	2.6

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	GOL	B	704	6/6	0.73	0.25	49,52,73,78	0
2	GOL	B	702	6/6	0.82	0.27	36,51,55,65	0
3	FLC	A	702	13/13	0.83	0.25	28,36,60,67	0
2	GOL	B	703	6/6	0.85	0.35	35,58,65,66	0
2	GOL	B	701	6/6	0.88	0.27	21,38,43,59	0
2	GOL	A	701	6/6	0.90	0.27	39,48,56,59	0

6.5 Other polymers [i](#)

There are no such residues in this entry.