



# wwPDB X-ray Structure Validation Summary Report ⓘ

May 25, 2020 – 05:14 am BST

PDB ID : 3LYU  
Title : Crystal Structure of the C-terminal domain (residues 83-215) of PF1911 hydrogenase from *Pyrococcus furiosus*, Northeast Structural Genomics Consortium Target PfR246A  
Authors : Forouhar, F.; Abashidze, M.; Seetharaman, J.; Sahdev, S.; Xiao, R.; Foote, E.L.; Ciccocanti, C.; Belote, R.L.; Everett, J.K.; Nair, R.; Acton, T.B.; Rost, B.; Montelione, G.T.; Tong, L.; Hunt, J.F.; Northeast Structural Genomics Consortium (NESG)  
Deposited on : 2010-02-28  
Resolution : 2.30 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	5042 (2.30-2.30)
Clashscore	141614	5643 (2.30-2.30)
Ramachandran outliers	138981	5575 (2.30-2.30)
Sidechain outliers	138945	5575 (2.30-2.30)
RSRZ outliers	127900	4938 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	142	
1	B	142	
1	C	142	
1	D	142	
1	E	142	
1	F	142	

## 2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 6280 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Putative hydrogenase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	Se			
1	A	129	1021	664	164	188	5	0	0	0
1	B	131	1040	675	168	192	5	0	0	0
1	C	130	1030	669	165	191	5	0	0	0
1	D	131	1040	675	168	192	5	0	0	0
1	E	131	1040	675	168	192	5	0	0	0
1	F	131	1040	675	168	192	5	0	0	0

There are 54 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	82	MSE	-	INITIATING METHIONINE	UNP Q8TZS3
A	216	LEU	-	EXPRESSION TAG	UNP Q8TZS3
A	217	GLU	-	EXPRESSION TAG	UNP Q8TZS3
A	218	HIS	-	EXPRESSION TAG	UNP Q8TZS3
A	219	HIS	-	EXPRESSION TAG	UNP Q8TZS3
A	220	HIS	-	EXPRESSION TAG	UNP Q8TZS3
A	221	HIS	-	EXPRESSION TAG	UNP Q8TZS3
A	222	HIS	-	EXPRESSION TAG	UNP Q8TZS3
A	223	HIS	-	EXPRESSION TAG	UNP Q8TZS3
B	82	MSE	-	INITIATING METHIONINE	UNP Q8TZS3
B	216	LEU	-	EXPRESSION TAG	UNP Q8TZS3
B	217	GLU	-	EXPRESSION TAG	UNP Q8TZS3
B	218	HIS	-	EXPRESSION TAG	UNP Q8TZS3
B	219	HIS	-	EXPRESSION TAG	UNP Q8TZS3
B	220	HIS	-	EXPRESSION TAG	UNP Q8TZS3
B	221	HIS	-	EXPRESSION TAG	UNP Q8TZS3
B	222	HIS	-	EXPRESSION TAG	UNP Q8TZS3

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
B	223	HIS	-	EXPRESSION TAG	UNP Q8TZS3
C	82	MSE	-	INITIATING METHIONINE	UNP Q8TZS3
C	216	LEU	-	EXPRESSION TAG	UNP Q8TZS3
C	217	GLU	-	EXPRESSION TAG	UNP Q8TZS3
C	218	HIS	-	EXPRESSION TAG	UNP Q8TZS3
C	219	HIS	-	EXPRESSION TAG	UNP Q8TZS3
C	220	HIS	-	EXPRESSION TAG	UNP Q8TZS3
C	221	HIS	-	EXPRESSION TAG	UNP Q8TZS3
C	222	HIS	-	EXPRESSION TAG	UNP Q8TZS3
C	223	HIS	-	EXPRESSION TAG	UNP Q8TZS3
D	82	MSE	-	INITIATING METHIONINE	UNP Q8TZS3
D	216	LEU	-	EXPRESSION TAG	UNP Q8TZS3
D	217	GLU	-	EXPRESSION TAG	UNP Q8TZS3
D	218	HIS	-	EXPRESSION TAG	UNP Q8TZS3
D	219	HIS	-	EXPRESSION TAG	UNP Q8TZS3
D	220	HIS	-	EXPRESSION TAG	UNP Q8TZS3
D	221	HIS	-	EXPRESSION TAG	UNP Q8TZS3
D	222	HIS	-	EXPRESSION TAG	UNP Q8TZS3
D	223	HIS	-	EXPRESSION TAG	UNP Q8TZS3
E	82	MSE	-	INITIATING METHIONINE	UNP Q8TZS3
E	216	LEU	-	EXPRESSION TAG	UNP Q8TZS3
E	217	GLU	-	EXPRESSION TAG	UNP Q8TZS3
E	218	HIS	-	EXPRESSION TAG	UNP Q8TZS3
E	219	HIS	-	EXPRESSION TAG	UNP Q8TZS3
E	220	HIS	-	EXPRESSION TAG	UNP Q8TZS3
E	221	HIS	-	EXPRESSION TAG	UNP Q8TZS3
E	222	HIS	-	EXPRESSION TAG	UNP Q8TZS3
E	223	HIS	-	EXPRESSION TAG	UNP Q8TZS3
F	82	MSE	-	INITIATING METHIONINE	UNP Q8TZS3
F	216	LEU	-	EXPRESSION TAG	UNP Q8TZS3
F	217	GLU	-	EXPRESSION TAG	UNP Q8TZS3
F	218	HIS	-	EXPRESSION TAG	UNP Q8TZS3
F	219	HIS	-	EXPRESSION TAG	UNP Q8TZS3
F	220	HIS	-	EXPRESSION TAG	UNP Q8TZS3
F	221	HIS	-	EXPRESSION TAG	UNP Q8TZS3
F	222	HIS	-	EXPRESSION TAG	UNP Q8TZS3
F	223	HIS	-	EXPRESSION TAG	UNP Q8TZS3

- Molecule 2 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	7	Total O 7 7	0	0

*Continued on next page...*


*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Residues</b>	<b>Atoms</b>	<b>ZeroOcc</b>	<b>AltConf</b>
2	B	14	Total O 14 14	0	0
2	C	10	Total O 10 10	0	0
2	D	14	Total O 14 14	0	0
2	E	16	Total O 16 16	0	0
2	F	8	Total O 8 8	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Putative hydrogenase




- Molecule 1: Putative hydrogenase



- Molecule 1: Putative hydrogenase



- Molecule 1: Putative hydrogenase





- Molecule 1: Putative hydrogenase



- Molecule 1: Putative hydrogenase



## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 31	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	77.65Å 77.65Å 117.51Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	19.59 – 2.30 29.18 – 2.30	Depositor EDS
% Data completeness (in resolution range)	93.2 (19.59-2.30) 99.7 (29.18-2.30)	Depositor EDS
$R_{merge}$	0.09	Depositor
$R_{sym}$	0.09	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.20 (at 2.29Å)	Xtriage
Refinement program	CNS 1.2, REFMAC	Depositor
R, $R_{free}$	0.210 , 0.237 0.212 , 0.238	Depositor DCC
$R_{free}$ test set	6466 reflections (9.20%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	38.4	Xtriage
Anisotropy	0.419	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.40 , 32.5	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.37$ , $\langle L^2 \rangle = 0.20$	Xtriage
Estimated twinning fraction	0.119 for -h,-k,l 0.449 for h,-h-k,-l 0.125 for -k,-h,-l	Xtriage
Reported twinning fraction	0.497 for H, K, L 0.503 for K, H, -L	Depositor
Outliers	0 of 70298 reflections	Xtriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	6280	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	45.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.73% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup> Intensities estimated from amplitudes.

<sup>2</sup> Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.44	0/1039	0.69	2/1404 (0.1%)
1	B	0.44	0/1059	0.90	3/1431 (0.2%)
1	C	0.44	0/1048	0.68	2/1416 (0.1%)
1	D	0.48	0/1059	0.91	3/1431 (0.2%)
1	E	0.46	0/1059	0.70	3/1431 (0.2%)
1	F	0.50	0/1059	0.68	2/1431 (0.1%)
All	All	0.46	0/6323	0.77	15/8544 (0.2%)

There are no bond length outliers.

The worst 5 of 15 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	151	ARG	NE-CZ-NH2	17.50	129.05	120.30
1	B	151	ARG	NE-CZ-NH1	-17.02	111.79	120.30
1	D	151	ARG	NE-CZ-NH1	-16.88	111.86	120.30
1	B	151	ARG	NE-CZ-NH2	16.56	128.58	120.30
1	A	151	ARG	NE-CZ-NH2	-8.58	116.01	120.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1021	0	1043	74	0
1	B	1040	0	1056	75	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	C	1030	0	1049	73	0
1	D	1040	0	1056	79	0
1	E	1040	0	1056	86	0
1	F	1040	0	1056	86	0
2	A	7	0	0	0	0
2	B	14	0	0	3	0
2	C	10	0	0	3	0
2	D	14	0	0	4	0
2	E	16	0	0	2	0
2	F	8	0	0	0	0
All	All	6280	0	6316	430	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 34.

The worst 5 of 430 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:122:GLN:HE21	1:C:128:VAL:HG12	1.28	0.97
1:E:122:GLN:HE21	1:E:128:VAL:HG12	1.28	0.97
1:D:122:GLN:HE21	1:D:128:VAL:HG12	1.30	0.97
1:A:122:GLN:HE21	1:A:128:VAL:HG12	1.30	0.96
1:B:151:ARG:HD3	1:B:153:ILE:HD11	1.47	0.96

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	127/142 (89%)	122 (96%)	5 (4%)	0	100	100
1	B	129/142 (91%)	124 (96%)	5 (4%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	C	128/142 (90%)	123 (96%)	5 (4%)	0	100	100
1	D	129/142 (91%)	123 (95%)	6 (5%)	0	100	100
1	E	129/142 (91%)	123 (95%)	6 (5%)	0	100	100
1	F	129/142 (91%)	122 (95%)	7 (5%)	0	100	100
All	All	771/852 (90%)	737 (96%)	34 (4%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	114/120 (95%)	105 (92%)	9 (8%)	12	15
1	B	116/120 (97%)	107 (92%)	9 (8%)	12	16
1	C	115/120 (96%)	106 (92%)	9 (8%)	12	16
1	D	116/120 (97%)	107 (92%)	9 (8%)	12	16
1	E	116/120 (97%)	107 (92%)	9 (8%)	12	16
1	F	116/120 (97%)	105 (90%)	11 (10%)	8	10
All	All	693/720 (96%)	637 (92%)	56 (8%)	11	15

5 of 56 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	187	GLU
1	D	160	ASN
1	F	179	LYS
1	C	209	LYS
1	D	124	ILE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 29 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	168	ASN
1	D	162	ASN
1	F	168	ASN
1	D	122	GLN
1	D	168	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

### 5.7 Other polymers [i](#)

There are no such residues in this entry.

### 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	124/142 (87%)	0.25	2 (1%) 72 77	24, 48, 64, 69	0
1	B	126/142 (88%)	0.17	1 (0%) 86 89	27, 49, 63, 71	0
1	C	125/142 (88%)	0.14	1 (0%) 86 89	23, 49, 62, 69	0
1	D	126/142 (88%)	0.08	1 (0%) 86 89	22, 46, 62, 68	0
1	E	126/142 (88%)	-0.05	0 100 100	20, 44, 58, 70	0
1	F	126/142 (88%)	-0.03	1 (0%) 86 89	22, 43, 58, 66	0
All	All	753/852 (88%)	0.09	6 (0%) 86 89	20, 47, 62, 71	0

The worst 5 of 6 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	F	162	ASN	2.5
1	D	208	VAL	2.3
1	A	166	LEU	2.3
1	C	180	VAL	2.3
1	B	213	VAL	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers

There are no such residues in this entry.