



wwPDB EM Validation Summary Report ⓘ

Nov 20, 2022 – 02:41 PM EST

PDB ID : 7MIX
EMDB ID : EMD-23867
Title : Human N-type voltage-gated calcium channel Cav2.2 in the presence of ziconotide at 3.0 Angstrom resolution
Authors : Yan, N.; Gao, S.; Yao, X.
Deposited on : 2021-04-18
Resolution : 3.00 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev43
Mogul : 1.8.5 (274361), CSD as541be (2020)
MolProbity : 4.02b-467
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
MapQ : 1.9.9
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.31.2


1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

ELECTRON MICROSCOPY

The reported resolution of this entry is 3.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)
Clashscore	158937	4297
Ramachandran outliers	154571	4023
Sidechain outliers	154315	3826

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	2339	
2	B	25	
3	D	1103	
4	C	484	
5	G	3	
6	J	2	
6	P	2	
6	S	2	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
7	L	4	 <p>A horizontal bar chart representing the quality of chain. The bar is divided into four segments: a red segment (top) and a green segment (bottom) each labeled '25%', followed by a yellow segment (top) labeled '25%', and an orange segment (bottom) labeled '50%'. The segments are stacked horizontally to represent a total of 100%.</p>

2 Entry composition i

There are 13 unique types of molecules in this entry. The entry contains 21652 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Voltage-dependent N-type calcium channel subunit alpha-1B.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
1	A	1326	10740	7049	1746	1868	77	1	0

- Molecule 2 is a protein called Omega-conotoxin MVIIA.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
2	B	25	176	102	35	32	7	0	0

- Molecule 3 is a protein called Voltage-dependent calcium channel subunit alpha-2/delta-1.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
3	D	948	7570	4803	1269	1467	31	0	0

- Molecule 4 is a protein called Voltage-dependent L-type calcium channel subunit beta-3.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
4	C	324	2575	1619	467	479	10	0	0

- Molecule 5 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				AltConf	Trace
			Total	C	N	O		
5	G	3	42	24	3	15	0	0

- Molecule 6 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				AltConf	Trace
			Total	C	N	O		
6	J	2	28	16	2	10	0	0
6	P	2	28	16	2	10	0	0
6	S	2	28	16	2	10	0	0

- Molecule 7 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				AltConf	Trace
			Total	C	N	O		
7	L	4	56	32	4	20	0	0

- Molecule 8 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: $C_8H_{15}NO_6$).




Mol	Chain	Residues	Atoms				AltConf
			Total	C	N	O	
8	A	1	14	8	1	5	0
8	D	1	28	16	2	10	0
8	D	1	28	16	2	10	0

- Molecule 9 is CALCIUM ION (three-letter code: CA) (formula: Ca) (labeled as "Ligand of Interest" by depositor).


Mol	Chain	Residues	Atoms		AltConf
			Total	Ca	
9	A	1	1	1	0
9	D	1	1	1	0

- Molecule 10 is CHOLESTEROL HEMISUCCINATE (three-letter code: Y01) (formula: C₃₁H₅₀O₄) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms			AltConf
10	A	1	Total	C	O	0
			70	62	8	
10	A	1	Total	C	O	0
			70	62	8	

- Molecule 11 is CHOLESTEROL (three-letter code: CLR) (formula: $C_{27}H_{46}O$) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms			AltConf
11	A	1	Total	C	O	0
			140	135	5	

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms			AltConf
			Total	C	O	
11	A	1	140	135	5	0
11	A	1	140	135	5	0
11	A	1	140	135	5	0
11	A	1	140	135	5	0

- Molecule 12 is 1,2-Distearoyl-sn-glycerophosphoethanolamine (three-letter code: 3PE) (formula: $C_{41}H_{82}NO_8P$).



Mol	Chain	Residues	Atoms					AltConf
			Total	C	N	O	P	
12	A	1	91	71	2	16	2	0
12	A	1	91	71	2	16	2	0

- Molecule 13 is [(2R)-1-octadecanoyloxy-3-[oxidanyl-[(1R,2R,3S,4R,5R,6S)-2,3,6-tris(oxidanyl)-4,5-diphosphonoxy-cyclohexyl]oxy-phosphoryl]oxy-propan-2-yl] (8Z)-icosa-5,8,11,14-tetraenoate (three-letter code: PT5) (formula: $C_{47}H_{85}O_{19}P_3$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				AltConf
			Total	C	O	P	
13	A	1	64	42	19	3	0

SER
TYR
HIS
HIS
PRO
ASP
GLN
ASP
HIS
TRP
CYS


• Molecule 2: Omega-conotoxin MVIIIA



• Molecule 3: Voltage-dependent calcium channel subunit alpha-2/delta-1



- Molecule 4: Voltage-dependent L-type calcium channel subunit beta-3



- Molecule 5: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 6: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose




- Molecule 6: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 6: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 7: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	170839	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	50	Depositor
Minimum defocus (nm)	Not provided	
Maximum defocus (nm)	Not provided	
Magnification	Not provided	
Image detector	GATAN K2 SUMMIT (4k x 4k)	Depositor
Maximum map value	0.200	Depositor
Minimum map value	-0.128	Depositor
Average map value	0.000	Depositor
Map value standard deviation	0.006	Depositor
Recommended contour level	0.02	Depositor
Map size (Å)	311.91998, 311.91998, 311.91998	wwPDB
Map dimensions	280, 280, 280	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	1.114, 1.114, 1.114	Depositor

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CA, NAG, PT5, 3PE, CLR, Y01

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.27	0/11002	0.48	2/14897 (0.0%)
2	B	0.26	0/176	0.50	0/228
3	D	0.25	0/7728	0.49	0/10477
4	C	0.25	0/2624	0.51	0/3544
All	All	0.26	0/21530	0.48	2/29146 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	596[A]	ARG	CA-C-O	5.86	132.40	120.10
1	A	596[B]	ARG	CA-C-O	5.86	132.40	120.10

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	10740	0	10851	212	0
2	B	176	0	176	5	0
3	D	7570	0	7369	118	0
4	C	2575	0	2619	54	0
5	G	42	0	37	3	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
6	J	28	0	25	0	0
6	P	28	0	25	0	0
6	S	28	0	25	1	0
7	L	56	0	49	4	0
8	A	14	0	13	0	0
8	D	28	0	26	1	0
9	A	1	0	0	0	0
9	D	1	0	0	0	0
10	A	70	0	98	16	0
11	A	140	0	230	11	0
12	A	91	0	139	5	0
13	A	64	0	67	12	0
All	All	21652	0	21749	410	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

The worst 5 of 410 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1187:ASP:O	1:A:1191:THR:HG22	1.67	0.94
1:A:1195:THR:HG22	1:A:1227:VAL:HG13	1.48	0.94
1:A:254:PHE:HB3	1:A:284:ARG:NH2	1.83	0.92
1:A:160:SER:HA	1:A:163:ARG:HG3	1.52	0.90
1:A:476:ARG:HH11	1:A:476:ARG:HB3	1.35	0.89

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	1317/2339 (56%)	1237 (94%)	77 (6%)	3 (0%)	47	82
2	B	23/25 (92%)	23 (100%)	0	0	100	100
3	D	936/1103 (85%)	895 (96%)	41 (4%)	0	100	100
4	C	322/484 (66%)	307 (95%)	15 (5%)	0	100	100
All	All	2598/3951 (66%)	2462 (95%)	133 (5%)	3 (0%)	54	85

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	154	PHE
1	A	286	TYR
1	A	255	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	1165/1991 (58%)	1139 (98%)	26 (2%)	52	81
2	B	20/20 (100%)	19 (95%)	1 (5%)	24	60
3	D	837/971 (86%)	835 (100%)	2 (0%)	93	98
4	C	287/426 (67%)	282 (98%)	5 (2%)	60	85
All	All	2309/3408 (68%)	2275 (98%)	34 (2%)	66	87

5 of 34 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	D	1029	ARG
4	C	198	LYS
4	C	296	SER
1	A	606	MET
1	A	590	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	702	ASN
1	A	1673	GLN
1	A	1725	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

13 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	NAG	G	1	3,5	14,14,15	0.53	0	17,19,21	0.56	0
5	NAG	G	2	5	14,14,15	0.45	0	17,19,21	1.27	1 (5%)
5	NAG	G	3	5	14,14,15	0.35	0	17,19,21	0.37	0
6	NAG	J	1	3,6	14,14,15	0.34	0	17,19,21	0.50	0
6	NAG	J	2	6	14,14,15	0.49	0	17,19,21	0.77	0
7	NAG	L	1	3,7	14,14,15	0.26	0	17,19,21	0.44	0
7	NAG	L	2	7	14,14,15	0.26	0	17,19,21	0.40	0
7	NAG	L	3	7	14,14,15	0.98	1 (7%)	17,19,21	1.66	2 (11%)
7	NAG	L	4	7	14,14,15	0.52	0	17,19,21	1.26	1 (5%)
6	NAG	P	1	3,6	14,14,15	0.26	0	17,19,21	0.47	0
6	NAG	P	2	6	14,14,15	0.44	0	17,19,21	0.54	0
6	NAG	S	1	3,6	14,14,15	0.88	1 (7%)	17,19,21	1.25	1 (5%)
6	NAG	S	2	6	14,14,15	0.21	0	17,19,21	0.45	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral

centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '2' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	NAG	G	1	3,5	-	2/6/23/26	0/1/1/1
5	NAG	G	2	5	-	5/6/23/26	0/1/1/1
5	NAG	G	3	5	-	4/6/23/26	0/1/1/1
6	NAG	J	1	3,6	-	3/6/23/26	0/1/1/1
6	NAG	J	2	6	-	3/6/23/26	0/1/1/1
7	NAG	L	1	3,7	-	2/6/23/26	0/1/1/1
7	NAG	L	2	7	-	2/6/23/26	0/1/1/1
7	NAG	L	3	7	-	5/6/23/26	0/1/1/1
7	NAG	L	4	7	-	5/6/23/26	0/1/1/1
6	NAG	P	1	3,6	-	2/6/23/26	0/1/1/1
6	NAG	P	2	6	-	2/6/23/26	0/1/1/1
6	NAG	S	1	3,6	-	3/6/23/26	0/1/1/1
6	NAG	S	2	6	-	4/6/23/26	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
7	L	3	NAG	O5-C1	3.16	1.48	1.43
6	S	1	NAG	O5-C1	2.57	1.47	1.43

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	S	1	NAG	C1-O5-C5	4.81	118.71	112.19
7	L	3	NAG	C1-O5-C5	4.60	118.42	112.19
7	L	3	NAG	C2-N2-C7	4.34	129.09	122.90
7	L	4	NAG	C2-N2-C7	4.29	129.02	122.90
5	G	2	NAG	C2-N2-C7	4.26	128.97	122.90

There are no chirality outliers.

5 of 42 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	G	2	NAG	O5-C5-C6-O6
7	L	3	NAG	C4-C5-C6-O6
6	P	1	NAG	O5-C5-C6-O6

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
6	P	2	NAG	O5-C5-C6-O6
7	L	3	NAG	O5-C5-C6-O6


There are no ring outliers.


6 monomers are involved in 8 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	G	2	NAG	2	0
7	L	4	NAG	1	0
7	L	1	NAG	2	0
5	G	1	NAG	2	0
7	L	3	NAG	1	0
6	S	1	NAG	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.











5.6 Ligand geometry [i](#)

Of 15 ligands modelled in this entry, 2 are monoatomic - leaving 13 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
11	CLR	A	2404	-	31,31,31	1.08	2 (6%)	48,48,48	1.30	6 (12%)
11	CLR	A	2409	-	31,31,31	1.07	1 (3%)	48,48,48	1.28	7 (14%)
8	NAG	A	2401	1	14,14,15	0.33	0	17,19,21	0.47	0
11	CLR	A	2407	-	31,31,31	1.06	1 (3%)	48,48,48	1.33	7 (14%)
11	CLR	A	2408	-	31,31,31	1.04	2 (6%)	48,48,48	1.36	7 (14%)
13	PT5	A	2412	-	64,64,69	0.85	2 (3%)	78,82,87	1.01	3 (3%)
10	Y01	A	2406	-	38,38,38	2.71	11 (28%)	57,57,57	4.40	33 (57%)
11	CLR	A	2405	-	31,31,31	1.04	2 (6%)	48,48,48	1.37	9 (18%)
8	NAG	D	1202	3	14,14,15	0.87	1 (7%)	17,19,21	0.88	1 (5%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
10	Y01	A	2403	-	38,38,38	2.43	11 (28%)	57,57,57	4.36	33 (57%)
12	3PE	A	2411	-	50,50,50	0.51	0	53,55,55	0.54	1 (1%)
12	3PE	A	2410	-	39,39,50	0.57	0	42,44,55	0.56	1 (2%)
8	NAG	D	1201	3	14,14,15	0.36	0	17,19,21	0.46	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
11	CLR	A	2404	-	-	7/10/68/68	0/4/4/4
11	CLR	A	2409	-	-	8/10/68/68	0/4/4/4
8	NAG	A	2401	1	-	2/6/23/26	0/1/1/1
11	CLR	A	2407	-	-	7/10/68/68	0/4/4/4
11	CLR	A	2408	-	-	6/10/68/68	0/4/4/4
13	PT5	A	2412	-	-	25/61/85/90	0/1/1/1
10	Y01	A	2406	-	-	13/19/77/77	0/4/4/4
11	CLR	A	2405	-	-	2/10/68/68	0/4/4/4
8	NAG	D	1202	3	-	2/6/23/26	0/1/1/1
10	Y01	A	2403	-	-	8/19/77/77	0/4/4/4
12	3PE	A	2411	-	-	25/54/54/54	-
12	3PE	A	2410	-	-	9/43/43/54	-
8	NAG	D	1201	3	-	1/6/23/26	0/1/1/1

The worst 5 of 33 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
10	A	2406	Y01	CAK-CBD	10.83	1.71	1.53
10	A	2403	Y01	CAK-CBD	8.36	1.67	1.53
10	A	2406	Y01	CBD-CBG	-6.36	1.41	1.53
10	A	2403	Y01	CBD-CBG	-6.09	1.42	1.53
10	A	2403	Y01	CAI-CAZ	4.33	1.42	1.33

The worst 5 of 108 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
10	A	2406	Y01	CAD-CBH-CAT	11.66	127.85	109.43
10	A	2403	Y01	CAD-CBH-CAT	11.29	127.26	109.43

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
10	A	2403	Y01	CAK-CBD-CBG	-9.58	97.02	110.91
10	A	2403	Y01	CAD-CBH-CAZ	-9.49	92.99	108.34
10	A	2406	Y01	CAD-CBH-CAZ	-9.03	93.72	108.34

There are no chirality outliers.

5 of 115 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
10	A	2403	Y01	OAG-CAY-OAW-CBC
10	A	2403	Y01	CAX-CAL-CAM-CAY
10	A	2406	Y01	OAG-CAY-OAW-CBC
10	A	2406	Y01	CAM-CAY-OAW-CBC
12	A	2410	3PE	C1-O11-P-O12



There are no ring outliers.


10 monomers are involved in 45 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
11	A	2404	CLR	3	0
11	A	2409	CLR	2	0
11	A	2408	CLR	4	0
13	A	2412	PT5	12	0
10	A	2406	Y01	5	0
11	A	2405	CLR	2	0
8	D	1202	NAG	1	0
10	A	2403	Y01	11	0
12	A	2411	3PE	4	0
12	A	2410	3PE	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.











5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-23867. These allow visual inspection of the internal detail of the map and identification of artifacts.

No raw map or half-maps were deposited for this entry and therefore no images, graphs, etc. pertaining to the raw map can be shown.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map




The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 136



Y Index: 141



Z Index: 141

The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal surface views [i](#)

6.4.1 Primary map



X



Y



Z

The images above show the 3D surface view of the map at the recommended contour level 0.02. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.


6.5 Mask visualisation

This section was not generated. No masks/segmentation were deposited.

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 135 nm³; this corresponds to an approximate mass of 122 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum [\(i\)](#)



*Reported resolution corresponds to spatial frequency of 0.333\AA^{-1}


8 Fourier-Shell correlation

This section was not generated. No FSC curve or half-maps provided.

9 Map-model fit [i](#)


This section contains information regarding the fit between EMDB map EMD-23867 and PDB model 7MIX. Per-residue inclusion information can be found in section 3 on page 10.

9.1 Map-model overlay [i](#)




The images above show the 3D surface view of the map at the recommended contour level 0.02 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.02).

9.4 Atom inclusion [i](#)



At the recommended contour level, 75% of all backbone atoms, 69% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary

The table lists the average atom inclusion at the recommended contour level (0.02) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	 0.6946	 0.4960
A	 0.7230	 0.5090
B	 0.9181	 0.6050
C	 0.2908	 0.3270
D	 0.7885	 0.5320
G	 0.5000	 0.3530
J	 0.2857	 0.3440
L	 0.7143	 0.4910
P	 0.4286	 0.3640
S	 0.4643	 0.3820

