



Full wwPDB X-ray Structure Validation Report ⓘ

Jun 11, 2024 – 05:24 PM EDT

PDB ID : 6MTW
Title : Lysosomal Phospholipase A2 in complex with Zinc
Authors : Bouley, R.; Tesmer, J.J.G.
Deposited on : 2018-10-22
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36.2
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	8085 (2.00-2.00)
Clashscore	141614	9178 (2.00-2.00)
Ramachandran outliers	138981	9054 (2.00-2.00)
Sidechain outliers	138945	9053 (2.00-2.00)
RSRZ outliers	127900	7900 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	380	<div> <div>6%</div> <div> <div></div> <div>87%</div> <div>12%</div> <div>.</div> </div> </div>
1	B	380	<div> <div>9%</div> <div> <div></div> <div>91%</div> <div>8%</div> <div>.</div> </div> </div>

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Group XV phospholipase A2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	376	Total 3019	C 1949	N 507	O 550	S 13	0	0	0
1	B	376	Total 3019	C 1949	N 507	O 550	S 13	0	0	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	GLY	-	expression tag	UNP Q8NCC3
B	0	GLY	-	expression tag	UNP Q8NCC3

- Molecule 2 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: $\text{C}_8\text{H}_{15}\text{NO}_6$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			14	8	1	5		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	B	1	Total	C	N	O	0	0
			14	8	1	5		
2	B	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 3 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Cl	0	0
			1	1		
3	B	1	Total	Cl	0	0
			1	1		

- Molecule 4 is ZINC ION (three-letter code: ZN) (formula: Zn) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	4	Total	Zn	0	0
			4	4		
4	B	4	Total	Zn	0	0
			4	4		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	174	Total	O	0	0
			174	174		
5	B	168	Total	O	0	0
			168	168		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Group XV phospholipase A2



- Molecule 1: Group XV phospholipase A2



4 Data and refinement statistics

Property	Value	Source
Space group	H 3	Depositor
Cell constants a, b, c, α , β , γ	84.57Å 84.57Å 321.74Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	39.34 – 2.00 39.34 – 2.00	Depositor EDS
% Data completeness (in resolution range)	98.8 (39.34-2.00) 89.2 (39.34-2.00)	Depositor EDS
R_{merge}	0.15	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	0.53 (at 2.00Å)	Xtriage
Refinement program	PHENIX (1.13_2998: ???)	Depositor
R, R_{free}	0.247 , 0.285 0.247 , 0.285	Depositor DCC
R_{free} test set	1969 reflections (3.45%)	wwPDB-VP
Wilson B-factor (Å ²)	25.5	Xtriage
Anisotropy	0.294	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 44.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.033 for -h-k,k,-l	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	6446	wwPDB-VP
Average B, all atoms (Å ²)	34.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 28.82 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 1.7236e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CL, NAG, ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.26	0/3108	0.46	1/4236 (0.0%)
1	B	0.26	0/3108	0.47	0/4236
All	All	0.26	0/6216	0.47	1/8472 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	91	LEU	CA-CB-CG	5.05	126.92	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3019	0	2954	27	0
1	B	3019	0	2954	16	0
2	A	28	0	26	0	0
2	B	28	0	26	0	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
4	A	4	0	0	0	0
4	B	4	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	A	174	0	0	3	0
5	B	168	0	0	2	0
All	All	6446	0	5960	43	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (43) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:52:VAL:HG13	1:B:53:ILE:HG23	1.82	0.61
1:B:6:PRO:HG3	1:B:155:TYR:HB2	1.85	0.59
1:A:52:VAL:HG13	1:A:53:ILE:HG23	1.84	0.59
1:A:67:LYS:NZ	1:A:273:ASP:OD1	2.33	0.58
1:A:301:THR:HG22	1:A:332:LEU:HD13	1.89	0.55
1:B:67:LYS:NZ	1:B:273:ASP:OD1	2.31	0.54
1:B:202:LYS:NZ	5:B:508:HOH:O	2.42	0.53
1:B:301:THR:HG22	1:B:332:LEU:HD13	1.91	0.53
1:A:23:ASP:OD2	5:A:502:HOH:O	2.19	0.52
1:A:6:PRO:HG3	1:A:155:TYR:HB2	1.90	0.52
1:A:4:HIS:N	5:A:507:HOH:O	2.42	0.52
1:A:197:TRP:NE1	1:A:338:CYS:SG	2.83	0.51
1:A:39:TYR:OH	1:A:79:ASP:OD2	2.19	0.51
1:A:165:SER:OG	1:A:166:MET:N	2.46	0.49
1:B:213:ASN:ND2	5:B:506:HOH:O	2.45	0.48
1:B:165:SER:OG	1:B:166:MET:N	2.46	0.48
1:A:275:TRP:O	1:A:279:GLN:HG2	2.14	0.47
1:A:202:LYS:HA	1:A:309:PHE:CZ	2.50	0.47
1:A:238:PRO:HB3	1:A:243:TRP:CD1	2.50	0.46
1:B:238:PRO:HB3	1:B:243:TRP:CD1	2.51	0.46
1:A:342:GLN:HG2	1:A:349:VAL:HB	1.97	0.46
1:B:202:LYS:HA	1:B:309:PHE:CZ	2.52	0.45
1:A:310:TYR:HE2	1:A:312:GLU:HG2	1.82	0.45
1:B:4:HIS:NE2	1:B:122:ASP:OD2	2.33	0.44
1:B:281:THR:HA	1:B:284:LEU:HG	1.99	0.44
1:A:91:LEU:HG	1:A:105:PHE:CD2	2.52	0.44
1:A:149:GLU:OE2	5:A:503:HOH:O	2.20	0.43
1:A:70:ARG:NH2	1:A:280:ASP:OD1	2.52	0.43
1:A:333:LYS:HB3	1:A:333:LYS:HE3	1.87	0.43
1:A:43:TRP:HA	1:A:44:LEU:HA	1.72	0.43
1:A:213:ASN:HD22	1:A:213:ASN:HA	1.65	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:91:LEU:O	1:A:105:PHE:HB2	2.19	0.42
1:A:281:THR:HA	1:A:284:LEU:HG	2.01	0.42
1:A:313:SER:HB3	1:A:317:ARG:HG3	2.01	0.42
1:B:202:LYS:HB2	1:B:202:LYS:HE3	1.59	0.42
1:A:253:PRO:HD3	1:A:311:TYR:O	2.20	0.42
1:A:174:PHE:O	1:A:178:GLN:HG2	2.20	0.41
1:A:202:LYS:HA	1:A:309:PHE:HZ	1.85	0.41
1:A:363:LEU:HD23	1:A:363:LEU:HA	1.89	0.41
1:B:43:TRP:HA	1:B:44:LEU:HA	1.72	0.41
1:B:342:GLN:HG2	1:B:349:VAL:HB	2.04	0.40
1:B:174:PHE:O	1:B:178:GLN:HG2	2.22	0.40
1:B:290:PRO:HB2	1:B:345:GLN:OE1	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	374/380 (98%)	364 (97%)	10 (3%)	0	100	100
1	B	374/380 (98%)	364 (97%)	10 (3%)	0	100	100
All	All	748/760 (98%)	728 (97%)	20 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	328/329 (100%)	324 (99%)	4 (1%)	71	76
1	B	328/329 (100%)	324 (99%)	4 (1%)	71	76
All	All	656/658 (100%)	648 (99%)	8 (1%)	71	76

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	16	ASN
1	A	91	LEU
1	A	190	PHE
1	A	333	LYS
1	B	16	ASN
1	B	91	LEU
1	B	190	PHE
1	B	333	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	213	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 14 ligands modelled in this entry, 10 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	NAG	A	401	1	14,14,15	1.55	2 (14%)	17,19,21	1.54	4 (23%)
2	NAG	B	402	1	14,14,15	1.67	3 (21%)	17,19,21	1.51	4 (23%)
2	NAG	A	402	1	14,14,15	1.63	3 (21%)	17,19,21	1.53	3 (17%)
2	NAG	B	401	1	14,14,15	1.57	2 (14%)	17,19,21	1.53	4 (23%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	A	401	1	-	0/6/23/26	0/1/1/1
2	NAG	B	402	1	-	0/6/23/26	0/1/1/1
2	NAG	A	402	1	-	0/6/23/26	0/1/1/1
2	NAG	B	401	1	-	1/6/23/26	0/1/1/1

All (10) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	401	NAG	C1-C2	3.95	1.58	1.52
2	B	402	NAG	C1-C2	3.43	1.57	1.52
2	A	401	NAG	C1-C2	3.39	1.57	1.52
2	A	402	NAG	C1-C2	3.29	1.57	1.52
2	B	402	NAG	O5-C5	-2.80	1.37	1.43
2	A	402	NAG	O5-C5	-2.51	1.38	1.43
2	A	401	NAG	O5-C5	-2.38	1.38	1.43
2	A	402	NAG	C3-C2	2.21	1.57	1.52
2	B	402	NAG	C3-C2	2.03	1.56	1.52
2	B	401	NAG	O5-C5	-2.03	1.39	1.43

All (15) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	402	NAG	C2-N2-C7	-3.29	118.22	122.90
2	A	401	NAG	C6-C5-C4	-3.08	105.79	113.00
2	B	402	NAG	C2-N2-C7	-3.04	118.58	122.90
2	A	401	NAG	C2-N2-C7	-2.99	118.65	122.90
2	B	401	NAG	C2-N2-C7	-2.91	118.75	122.90
2	B	401	NAG	C6-C5-C4	-2.88	106.25	113.00
2	B	402	NAG	C1-C2-N2	-2.58	106.08	110.49
2	A	401	NAG	O4-C4-C3	-2.45	104.69	110.35
2	A	402	NAG	C6-C5-C4	-2.33	107.55	113.00
2	B	402	NAG	C6-C5-C4	-2.27	107.68	113.00
2	A	402	NAG	C1-C2-N2	-2.25	106.64	110.49
2	A	401	NAG	C1-C2-N2	-2.22	106.70	110.49
2	B	401	NAG	O4-C4-C3	-2.18	105.30	110.35
2	B	401	NAG	C1-C2-N2	-2.06	106.97	110.49
2	B	402	NAG	O4-C4-C3	-2.05	105.61	110.35

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	401	NAG	O5-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2		OWAB(Å ²)	Q < 0.9
1	A	376/380 (98%)	0.82	22 (5%)	22 21	19, 31, 53, 83	0
1	B	376/380 (98%)	0.90	36 (9%)	8 7	20, 32, 56, 76	0
All	All	752/760 (98%)	0.86	58 (7%)	13 12	19, 32, 54, 83	0

All (58) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	255	ILE	6.9
1	B	255	ILE	5.9
1	B	101	VAL	4.6
1	A	379	PRO	4.5
1	B	91	LEU	4.5
1	B	314	PHE	3.9
1	A	314	PHE	3.9
1	A	254	THR	3.8
1	B	336	LEU	3.6
1	B	316	ASP	3.4
1	B	275	TRP	3.3
1	A	91	LEU	3.3
1	B	312	GLU	3.2
1	A	275	TRP	3.2
1	A	95	ASP	3.1
1	A	312	GLU	3.1
1	B	286	GLU	3.1
1	A	37	GLU	3.0
1	A	336	LEU	3.0
1	B	254	THR	3.0
1	B	287	ALA	3.0
1	B	220	PRO	2.9
1	B	100	SER	2.9
1	B	340	ALA	2.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	337	GLN	2.8
1	A	271	PHE	2.8
1	B	95	ASP	2.7
1	A	133	ALA	2.7
1	B	37	GLU	2.7
1	B	334	SER	2.7
1	A	100	SER	2.7
1	B	26	THR	2.7
1	A	263	ARG	2.6
1	B	318	ASP	2.6
1	B	322	CYS	2.5
1	B	4	HIS	2.5
1	A	264	LYS	2.5
1	A	340	ALA	2.4
1	B	282	GLU	2.4
1	B	317	ARG	2.4
1	A	338	CYS	2.3
1	B	338	CYS	2.3
1	B	221	LEU	2.3
1	B	313	SER	2.3
1	B	276	LEU	2.3
1	B	96	PRO	2.2
1	B	99	SER	2.2
1	B	335	ALA	2.2
1	B	289	MET	2.1
1	B	102	GLY	2.1
1	A	259	LEU	2.1
1	A	96	PRO	2.1
1	B	238	PRO	2.1
1	B	264	LYS	2.1
1	A	201	ALA	2.1
1	A	276	LEU	2.1
1	B	170	TYR	2.0
1	A	204	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates

There are no monosaccharides in this entry.

6.4 Ligands




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	NAG	B	402	14/15	0.86	0.15	35,38,52,55	0
2	NAG	A	402	14/15	0.88	0.15	33,37,46,47	0
2	NAG	B	401	14/15	0.89	0.14	34,40,43,47	0
2	NAG	A	401	14/15	0.89	0.13	29,34,36,41	0
3	CL	A	403	1/1	0.96	0.15	48,48,48,48	0
3	CL	B	403	1/1	0.96	0.09	56,56,56,56	0
4	ZN	A	406	1/1	0.97	0.08	39,39,39,39	0
4	ZN	B	406	1/1	0.97	0.05	38,38,38,38	0
4	ZN	A	407	1/1	0.98	0.05	51,51,51,51	0
4	ZN	A	405	1/1	0.99	0.09	38,38,38,38	0
4	ZN	B	405	1/1	0.99	0.08	37,37,37,37	0
4	ZN	A	404	1/1	0.99	0.09	32,32,32,32	0
4	ZN	B	407	1/1	0.99	0.06	47,47,47,47	0
4	ZN	B	404	1/1	1.00	0.07	33,33,33,33	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around ZN A 406:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around ZN B 406:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around ZN A 407:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around ZN A 405:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




Electron density around ZN B 405:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around ZN A 404:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around ZN B 407:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around ZN B 404:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers ⓘ

There are no such residues in this entry.