



wwPDB EM Validation Summary Report ⓘ

Nov 19, 2022 – 03:32 pm GMT

PDB ID : 5N97
EMDB ID : EMD-3604
Title : Structure of the *C. crescentus* S-layer
Authors : Bharat, T.A.; Hagen, W.J.; Briggs, J.A.; Lowe, J.
Deposited on : 2017-02-24
Resolution : 7.40 Å (reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev43
MolProbity : 4.02b-467
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
MapQ : 1.9.9
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.31.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

ELECTRON MICROSCOPY

The reported resolution of this entry is 7.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)
Clashscore	158937	4297
Ramachandran outliers	154571	4023
Sidechain outliers	154315	3826

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	778	
1	B	778	
1	C	778	
1	D	778	
1	E	778	
1	F	778	

2 Entry composition [i](#)

There are 2 unique types of molecules in this entry. The entry contains 30912 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called S-layer protein rsaA.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
1	A	778	5133	3124	862	1145	2	0	0
1	B	778	5133	3124	862	1145	2	0	0
1	C	778	5133	3124	862	1145	2	0	0
1	D	778	5133	3124	862	1145	2	0	0
1	E	778	5133	3124	862	1145	2	0	0
1	F	778	5133	3124	862	1145	2	0	0


- Molecule 2 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		AltConf
2	A	19	Total	Ca	0
			19	19	
2	B	19	Total	Ca	0
			19	19	
2	C	19	Total	Ca	0
			19	19	
2	D	19	Total	Ca	0
			19	19	
2	E	19	Total	Ca	0
			19	19	
2	F	19	Total	Ca	0
			19	19	


3 Residue-property plots


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: S-layer protein *rsaA*






- Molecule 1: S-layer protein *rsaA*







• Molecule 1: S-layer protein rsaA








• Molecule 1: S-layer protein rsaA





• Molecule 1: S-layer protein rsaA





• Molecule 1: S-layer protein rsaA



4 Experimental information

Property	Value	Source
EM reconstruction method	SUBTOMOGRAM AVERAGING	Depositor
Imposed symmetry	2D CRYSTAL, $a=220 \text{ \AA}$, $b=220 \text{ \AA}$, $c=220 \text{ \AA}$, $\gamma=1^\circ$, space group=P 1 21	Depositor
Number of subtomograms used	51866	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION; Following Schur et al, 2016	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	3.4	Depositor
Minimum defocus (nm)	2000	Depositor
Maximum defocus (nm)	5000	Depositor
Magnification	105000	Depositor
Image detector	GATAN K2 SUMMIT (4k x 4k)	Depositor
Maximum map value	0.505	Depositor
Minimum map value	-0.252	Depositor
Average map value	-0.010	Depositor
Map value standard deviation	0.057	Depositor
Recommended contour level	0.11	Depositor
Map size (\AA)	270.0, 270.0, 270.0	wwPDB
Map dimensions	200, 200, 200	wwPDB
Map angles ($^\circ$)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (\AA)	1.35, 1.35, 1.35	Depositor

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.70	1/5166 (0.0%)	0.87	3/7112 (0.0%)
1	B	0.68	1/5166 (0.0%)	0.86	2/7112 (0.0%)
1	C	0.70	1/5166 (0.0%)	0.88	3/7112 (0.0%)
1	D	0.70	1/5166 (0.0%)	0.88	5/7112 (0.1%)
1	E	0.69	2/5166 (0.0%)	0.87	4/7112 (0.1%)
1	F	0.69	1/5166 (0.0%)	0.87	3/7112 (0.0%)
All	All	0.69	7/30996 (0.0%)	0.87	20/42672 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	B	0	1
1	C	0	1
1	D	0	1
1	E	0	1
1	F	0	1
All	All	0	6

The worst 5 of 7 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	F	842	THR	CA-CB	6.14	1.69	1.53
1	B	842	THR	CA-CB	5.75	1.68	1.53
1	A	842	THR	CA-CB	5.65	1.68	1.53
1	E	842	THR	CA-CB	5.54	1.67	1.53
1	C	842	THR	CA-CB	5.33	1.67	1.53

The worst 5 of 20 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	840	THR	N-CA-C	6.93	129.71	111.00
1	E	840	THR	N-CA-C	6.87	129.56	111.00
1	C	840	THR	N-CA-C	6.73	129.18	111.00
1	D	840	THR	N-CA-C	6.68	129.05	111.00
1	F	840	THR	N-CA-C	6.67	129.00	111.00

There are no chirality outliers.

5 of 6 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	594	GLY	Peptide
1	B	594	GLY	Peptide
1	C	594	GLY	Peptide
1	D	594	GLY	Peptide
1	E	594	GLY	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5133	0	5071	61	0
1	B	5133	0	5071	67	0
1	C	5133	0	5071	67	0
1	D	5133	0	5071	63	0
1	E	5133	0	5071	60	0
1	F	5133	0	5072	69	0
2	A	19	0	0	0	0
2	B	19	0	0	0	0
2	C	19	0	0	0	0
2	D	19	0	0	0	0
2	E	19	0	0	0	0
2	F	19	0	0	0	0
All	All	30912	0	30427	383	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 383 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:265:ALA:HA	1:B:291:ALA:H	1.21	1.05
1:E:265:ALA:HA	1:E:291:ALA:H	1.22	1.05
1:C:265:ALA:HA	1:C:291:ALA:H	1.22	1.04
1:A:265:ALA:HA	1:A:291:ALA:H	1.22	1.03
1:F:265:ALA:HA	1:F:291:ALA:H	1.22	1.00

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	776/778 (100%)	709 (91%)	52 (7%)	15 (2%)	8	38
1	B	776/778 (100%)	710 (92%)	52 (7%)	14 (2%)	8	40
1	C	776/778 (100%)	711 (92%)	51 (7%)	14 (2%)	8	40
1	D	776/778 (100%)	710 (92%)	51 (7%)	15 (2%)	8	38
1	E	776/778 (100%)	712 (92%)	49 (6%)	15 (2%)	8	38
1	F	776/778 (100%)	712 (92%)	50 (6%)	14 (2%)	8	40
All	All	4656/4668 (100%)	4264 (92%)	305 (7%)	87 (2%)	11	38

5 of 87 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	574	THR
1	A	936	SER
1	A	942	ALA
1	A	1019	ALA
1	B	574	THR

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	526/526 (100%)	473 (90%)	53 (10%)	7 25
1	B	526/526 (100%)	471 (90%)	55 (10%)	7 24
1	C	526/526 (100%)	472 (90%)	54 (10%)	7 25
1	D	526/526 (100%)	473 (90%)	53 (10%)	7 25
1	E	526/526 (100%)	472 (90%)	54 (10%)	7 25
1	F	526/526 (100%)	471 (90%)	55 (10%)	7 24
All	All	3156/3156 (100%)	2832 (90%)	324 (10%)	11 25

5 of 324 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	E	380	VAL
1	F	388	THR
1	E	491	THR
1	E	873	SER
1	F	620	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 60 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	685	ASN
1	F	359	ASN
1	D	359	ASN
1	F	358	GLN
1	F	709	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 114 ligands modelled in this entry, 114 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-3604. These allow visual inspection of the internal detail of the map and identification of artifacts.


No raw map or half-maps were deposited for this entry and therefore no images, graphs, etc. pertaining to the raw map can be shown.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 100



Y Index: 100




Z Index: 100


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 118



Y Index: 96




Z Index: 128

The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal surface views [i](#)


6.4.1 Primary map



X



Y



Z

The images above show the 3D surface view of the map at the recommended contour level 0.11. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.


6.5 Mask visualisation

This section was not generated. No masks/segmentation were deposited.

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [\(i\)](#)



The volume at the recommended contour level is 905 nm³; this corresponds to an approximate mass of 817 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum [i](#)



*Reported resolution corresponds to spatial frequency of 0.135 Å⁻¹


8 Fourier-Shell correlation

This section was not generated. No FSC curve or half-maps provided.


9 Map-model fit [i](#)

This section contains information regarding the fit between EMDB map EMD-3604 and PDB model 5N97. Per-residue inclusion information can be found in section 3 on page 4.


9.1 Map-model overlay [i](#)



X




Y



Z


The images above show the 3D surface view of the map at the recommended contour level 0.11 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.11).

9.4 Atom inclusion [i](#)



At the recommended contour level, 82% of all backbone atoms, 79% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary [i](#)

The table lists the average atom inclusion at the recommended contour level (0.11) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	 0.7922	 0.1010
A	 0.8266	 0.1050
B	 0.8079	 0.1180
C	 0.8145	 0.0980
D	 0.7713	 0.0960
E	 0.7407	 0.0830
F	 0.7921	 0.1030

