



# Full wwPDB X-ray Structure Validation Report ⓘ

Sep 12, 2023 – 10:39 AM EDT

PDB ID : 4OD8  
Title : Crystal structure of the vaccinia virus DNA polymerase holoenzyme subunit D4 in complex with the A20 N-terminus  
Authors : Contesto-Richefeu, C.; Tarbouriech, N.; Brazzolotto, X.; Burmeister, W.P.; Iseni, F.  
Deposited on : 2014-01-10  
Resolution : 1.85 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.35.1  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.35.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.85 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	2469 (1.86-1.86)
Clashscore	141614	2625 (1.86-1.86)
Ramachandran outliers	138981	2592 (1.86-1.86)
Sidechain outliers	138945	2592 (1.86-1.86)
RSRZ outliers	127900	2436 (1.86-1.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	232	81% 11% • 5%
1	B	232	82% 11% • 5%
2	C	52	77% 13% • • •
2	D	52	87% 8% • •

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	GOL	B	304	-	-	X	-

## 2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 4769 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Uracil-DNA glycosylase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	B	221	1797	1163	297	331	6	16	0	0
1	A	220	1788	1158	295	329	6	18	0	0

There are 28 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	-13	MET	-	expression tag	UNP P20536
B	-12	GLY	-	expression tag	UNP P20536
B	-11	SER	-	expression tag	UNP P20536
B	-10	SER	-	expression tag	UNP P20536
B	-9	HIS	-	expression tag	UNP P20536
B	-8	HIS	-	expression tag	UNP P20536
B	-7	HIS	-	expression tag	UNP P20536
B	-6	HIS	-	expression tag	UNP P20536
B	-5	HIS	-	expression tag	UNP P20536
B	-4	HIS	-	expression tag	UNP P20536
B	-3	SER	-	expression tag	UNP P20536
B	-2	GLN	-	expression tag	UNP P20536
B	-1	ASP	-	expression tag	UNP P20536
B	0	PRO	-	expression tag	UNP P20536
A	-13	MET	-	expression tag	UNP P20536
A	-12	GLY	-	expression tag	UNP P20536
A	-11	SER	-	expression tag	UNP P20536
A	-10	SER	-	expression tag	UNP P20536
A	-9	HIS	-	expression tag	UNP P20536
A	-8	HIS	-	expression tag	UNP P20536
A	-7	HIS	-	expression tag	UNP P20536
A	-6	HIS	-	expression tag	UNP P20536
A	-5	HIS	-	expression tag	UNP P20536
A	-4	HIS	-	expression tag	UNP P20536
A	-3	SER	-	expression tag	UNP P20536

*Continued on next page...*

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	GLN	-	expression tag	UNP P20536
A	-1	ASP	-	expression tag	UNP P20536
A	0	PRO	-	expression tag	UNP P20536


- Molecule 2 is a protein called DNA polymerase processivity factor component A20.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	D	51	Total	C	N	O	S	0	0	0
			403	259	64	79	1			
2	C	50	Total	C	N	O	S	0	0	0
			398	256	63	78	1			

There are 6 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
D	-1	GLY	-	expression tag	UNP P20995
D	0	ALA	-	expression tag	UNP P20995
D	2	ALA	THR	engineered mutation	UNP P20995
C	-1	GLY	-	expression tag	UNP P20995
C	0	ALA	-	expression tag	UNP P20995
C	2	ALA	THR	engineered mutation	UNP P20995

- Molecule 3 is SULFATE ION (three-letter code: SO<sub>4</sub>) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	B	1	Total O S 5 4 1	0	0
3	B	1	Total O S 5 4 1	0	0
3	B	1	Total O S 5 4 1	0	0
3	A	1	Total O S 5 4 1	0	0
3	A	1	Total O S 5 4 1	0	0
3	A	1	Total O S 5 4 1	0	0

- Molecule 4 is GLYCEROL (three-letter code: GOL) (formula:  $C_3H_8O_3$ ).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	B	1	Total C O 6 3 3	0	0
4	A	1	Total C O 6 3 3	0	0

- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	B	140	Total O 140 140	0	0
5	D	35	Total O 35 35	0	0

*Continued on next page...*


*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Residues</b>	<b>Atoms</b>		<b>ZeroOcc</b>	<b>AltConf</b>
5	A	139	Total 139	O 139	0	0
5	C	27	Total 27	O 27	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Uracil-DNA glycosylase




- Molecule 1: Uracil-DNA glycosylase



- Molecule 2: DNA polymerase processivity factor component A20



- Molecule 2: DNA polymerase processivity factor component A20



## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	92.62Å 92.62Å 145.86Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	46.31 – 1.85 44.14 – 1.85	Depositor EDS
% Data completeness (in resolution range)	99.9 (46.31-1.85) 99.9 (44.14-1.85)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.55 (at 1.86Å)	Xtrriage
Refinement program	REFMAC 5.8.0049	Depositor
R, $R_{free}$	0.175 , 0.218 0.183 , 0.228	Depositor DCC
$R_{free}$ test set	3134 reflections (5.06%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	29.8	Xtrriage
Anisotropy	0.035	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.37 , 41.5	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	0.023 for -h,-k,l	Xtrriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	4769	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	37.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.19% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality i

### 5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	1.03	1/1838 (0.1%)	1.11	13/2500 (0.5%)
1	B	1.14	8/1847 (0.4%)	1.39	15/2512 (0.6%)
2	C	1.08	1/405 (0.2%)	1.11	1/545 (0.2%)
2	D	1.25	0/410	1.11	3/552 (0.5%)
All	All	1.10	10/4500 (0.2%)	1.24	32/6109 (0.5%)

All (10) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	-2	GLN	CA-CB	-12.20	1.27	1.53
1	B	185	ARG	CG-CD	-9.12	1.29	1.51
1	B	36	TRP	CB-CG	-7.53	1.36	1.50
1	B	58	ARG	CD-NE	-6.68	1.35	1.46
1	B	3	SER	CB-OG	-6.18	1.34	1.42
1	B	7	SER	CB-OG	-6.00	1.34	1.42
1	B	32	GLU	CD-OE2	5.51	1.31	1.25
2	C	4	SER	CB-OG	-5.40	1.35	1.42
1	B	1	MET	C-O	-5.17	1.13	1.23
1	A	203	GLU	CD-OE2	5.02	1.31	1.25

All (32) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	58	ARG	NE-CZ-NH2	-30.88	104.86	120.30
1	B	58	ARG	NE-CZ-NH1	23.36	131.98	120.30
1	B	61	ARG	NE-CZ-NH1	15.04	127.82	120.30
1	B	61	ARG	NE-CZ-NH2	-10.58	115.01	120.30
1	B	193	ARG	NE-CZ-NH1	9.76	125.18	120.30
1	B	58	ARG	CD-NE-CZ	8.75	135.85	123.60
1	B	37	LEU	CA-CB-CG	8.12	133.98	115.30

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	21	ARG	NE-CZ-NH1	7.97	124.29	120.30
1	A	22	VAL	CG1-CB-CG2	7.96	123.64	110.90
1	A	193	ARG	NE-CZ-NH1	7.93	124.26	120.30
1	A	186	ASP	N-CA-CB	7.72	124.50	110.60
1	B	192	ASP	CB-CG-OD2	-7.46	111.59	118.30
1	B	-2	GLN	CB-CA-C	7.34	125.09	110.40
2	D	1	MET	CA-CB-CG	-7.25	100.97	113.30
1	A	37	LEU	CA-CB-CG	7.19	131.84	115.30
1	A	68	ASP	CB-CG-OD1	7.15	124.74	118.30
1	A	185	ARG	CB-CA-C	-7.02	96.36	110.40
1	B	61	ARG	CD-NE-CZ	6.64	132.90	123.60
1	B	58	ARG	CG-CD-NE	-6.38	98.40	111.80
1	A	47	ASP	CB-CG-OD1	6.34	124.00	118.30
1	B	39	ARG	NE-CZ-NH2	-6.04	117.28	120.30
1	A	192	ASP	CB-CG-OD2	-5.81	113.07	118.30
1	A	138	ASP	CB-CG-OD2	-5.64	113.22	118.30
1	B	61	ARG	CG-CD-NE	-5.54	100.16	111.80
1	B	192	ASP	N-CA-CB	-5.44	100.81	110.60
2	D	50	VAL	N-CA-CB	-5.44	99.54	111.50
2	D	50	VAL	CB-CA-C	5.34	121.55	111.40
1	B	185	ARG	CB-CG-CD	5.28	125.33	111.60
1	A	68	ASP	CB-CG-OD2	-5.24	113.59	118.30
1	A	98	ARG	NE-CZ-NH2	-5.12	117.74	120.30
1	A	166	ILE	CB-CA-C	-5.07	101.46	111.60
1	A	17	ASP	CB-CG-OD1	5.02	122.81	118.30

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1788	0	1779	13	3
1	B	1797	0	1789	15	3
2	C	398	0	404	7	0
2	D	403	0	409	2	0

Continued on next page...

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	A	15	0	0	0	0
3	B	15	0	0	0	0
4	A	6	0	8	1	0
4	B	6	0	8	4	0
5	A	139	0	0	2	0
5	B	140	0	0	4	0
5	C	27	0	0	0	0
5	D	35	0	0	0	0
All	All	4769	0	4397	37	3

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (37) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:68:ASP:O	4:B:304:GOL:H12	1.63	0.96
4:B:304:GOL:H11	5:B:531:HOH:O	1.77	0.83
1:B:16:HIS:HD2	1:B:18:ASP:H	1.27	0.82
1:A:16:HIS:HD2	1:A:18:ASP:H	1.34	0.76
1:B:171:GLU:O	5:B:535:HOH:O	2.09	0.69
1:A:16:HIS:CD2	1:A:18:ASP:H	2.14	0.65
1:B:16:HIS:CD2	1:B:18:ASP:H	2.12	0.64
1:B:58:ARG:HD2	5:A:444:HOH:O	2.00	0.60
1:B:160:LYS:NZ	5:B:540:HOH:O	2.32	0.60
1:B:181:HIS:CD2	1:B:183:ALA:H	2.21	0.59
2:C:21:ARG:HH11	2:C:21:ARG:HG2	1.70	0.57
2:D:0:ALA:O	2:D:1:MET:HB2	2.04	0.57
1:A:171:GLU:OE2	1:A:171:GLU:HA	2.05	0.56
1:A:163:PHE:O	1:A:166:ILE:HG13	2.07	0.55
1:A:139:LYS:HD3	1:A:139:LYS:H	1.73	0.53
2:C:20:LEU:HD21	2:C:28:ILE:HD13	1.91	0.52
2:C:20:LEU:CD2	2:C:28:ILE:HD13	2.42	0.50
1:A:139:LYS:H	1:A:139:LYS:CD	2.25	0.49
1:A:87:LYS:NZ	1:A:87:LYS:HB2	2.27	0.49
1:B:68:ASP:O	4:B:304:GOL:C1	2.51	0.48
1:A:16:HIS:HE1	5:A:532:HOH:O	1.95	0.48
1:B:109:ASN:ND2	1:B:111:ASN:H	2.12	0.47
1:B:79:PHE:HA	4:B:304:GOL:H32	1.97	0.46
1:B:2:ASN:HB2	1:B:15:TYR:CZ	2.51	0.46
1:A:4:VAL:HG21	1:A:27:VAL:HG21	1.98	0.46

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:160:LYS:CE	5:B:540:HOH:O	2.65	0.45
1:A:68:ASP:HB2	1:A:69:PRO:CD	2.47	0.44
2:C:14:LEU:O	2:C:18:LYS:HG3	2.18	0.44
1:A:120:ASN:OD1	4:A:304:GOL:H32	2.18	0.44
1:B:181:HIS:HD2	1:B:183:ALA:HB3	1.84	0.43
2:C:20:LEU:HD23	2:C:20:LEU:HA	1.89	0.42
2:C:32:ASN:HD21	2:C:50:VAL:CG2	2.33	0.41
1:A:149:THR:HA	1:A:152:VAL:HG22	2.01	0.41
1:B:87:LYS:HA	1:B:87:LYS:HD2	1.85	0.41
1:B:181:HIS:HD2	1:B:183:ALA:H	1.67	0.41
2:D:0:ALA:O	2:D:1:MET:CB	2.69	0.41
1:A:177:ILE:HD11	2:C:2:ALA:HB2	2.02	0.41

All (3) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:185:ARG:CZ	1:A:58:ARG:NH1[6_455]	1.46	0.74
1:B:185:ARG:NH2	1:A:58:ARG:NH1[6_455]	1.50	0.70
1:B:185:ARG:NE	1:A:58:ARG:NH1[6_455]	1.92	0.28

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	218/232 (94%)	213 (98%)	4 (2%)	1 (0%)	29	15
1	B	219/232 (94%)	210 (96%)	9 (4%)	0	100	100
2	C	48/52 (92%)	47 (98%)	1 (2%)	0	100	100
2	D	49/52 (94%)	46 (94%)	3 (6%)	0	100	100
All	All	534/568 (94%)	516 (97%)	17 (3%)	1 (0%)	47	33

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	138	ASP

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	202/213 (95%)	192 (95%)	10 (5%)	24	9
1	B	203/213 (95%)	195 (96%)	8 (4%)	32	15
2	C	44/44 (100%)	40 (91%)	4 (9%)	9	1
2	D	44/44 (100%)	40 (91%)	4 (9%)	9	1
All	All	493/514 (96%)	467 (95%)	26 (5%)	22	8

All (26) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	-1	ASP
1	B	38	LEU
1	B	61	ARG
1	B	82	PRO
1	B	87	LYS
1	B	94	SER
1	B	109	ASN
1	B	165	ASN
2	D	19	SER
2	D	20	LEU
2	D	21	ARG
2	D	50	VAL
1	A	-1	ASP
1	A	22	VAL
1	A	37	LEU
1	A	38	LEU
1	A	39	ARG
1	A	139	LYS
1	A	165	ASN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	166	ILE
1	A	169	LYS
1	A	171	GLU
2	C	14	LEU
2	C	20	LEU
2	C	21	ARG
2	C	48	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (11) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	16	HIS
1	B	109	ASN
1	B	146	GLN
1	B	165	ASN
1	B	181	HIS
1	B	206	ASN
1	A	2	ASN
1	A	16	HIS
1	A	146	GLN
1	A	151	HIS
1	A	165	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

8 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	GOL	A	304	-	5,5,5	0.58	0	5,5,5	0.85	0
3	SO4	B	301	-	4,4,4	0.48	0	6,6,6	0.86	0
3	SO4	A	301	-	4,4,4	0.33	0	6,6,6	0.28	0
4	GOL	B	304	-	5,5,5	0.95	0	5,5,5	1.88	3 (60%)
3	SO4	A	302	-	4,4,4	0.41	0	6,6,6	0.62	0
3	SO4	B	302	-	4,4,4	0.30	0	6,6,6	0.84	0
3	SO4	A	303	-	4,4,4	0.44	0	6,6,6	0.83	0
3	SO4	B	303	-	4,4,4	0.46	0	6,6,6	1.62	2 (33%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '2' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GOL	A	304	-	-	3/4/4/4	-
4	GOL	B	304	-	-	2/4/4/4	-

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	303	SO4	O4-S-O3	2.98	121.78	109.06
4	B	304	GOL	O3-C3-C2	2.66	122.95	110.20
3	B	303	SO4	O4-S-O1	-2.25	97.58	109.31
4	B	304	GOL	O2-C2-C1	-2.23	99.31	109.12
4	B	304	GOL	O2-C2-C3	2.09	118.33	109.12

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	B	304	GOL	O1-C1-C2-C3

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms
4	A	304	GOL	C1-C2-C3-O3
4	B	304	GOL	O1-C1-C2-O2
4	A	304	GOL	O2-C2-C3-O3
4	A	304	GOL	O1-C1-C2-C3

There are no ring outliers.

2 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	304	GOL	1	0
4	B	304	GOL	4	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	220/232 (94%)	-0.20	3 (1%) 75 76	23, 33, 53, 77	3 (1%)
1	B	221/232 (95%)	0.10	6 (2%) 54 53	20, 34, 62, 76	4 (1%)
2	C	50/52 (96%)	-0.36	0 100 100	25, 34, 55, 62	0
2	D	51/52 (98%)	0.06	4 (7%) 13 13	22, 32, 62, 88	0
All	All	542/568 (95%)	-0.07	13 (2%) 59 57	20, 34, 61, 88	7 (1%)

All (13) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	103	ILE	4.4
2	D	0	ALA	4.3
1	B	218	TYR	3.5
1	A	184	ALA	2.9
1	A	39	ARG	2.5
1	B	206	ASN	2.5
1	A	36	TRP	2.4
2	D	23	SER	2.3
1	B	-1	ASP	2.3
1	B	104	ASP	2.2
1	B	98	ARG	2.1
2	D	21	ARG	2.1
2	D	20	LEU	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
4	GOL	B	304	6/6	0.93	0.12	33,49,52,60	0
3	SO4	A	303	5/5	0.95	0.14	33,39,42,45	5
3	SO4	B	302	5/5	0.96	0.15	35,35,37,37	5
4	GOL	A	304	6/6	0.96	0.11	28,44,51,54	0
3	SO4	B	301	5/5	0.97	0.20	51,52,65,67	0
3	SO4	B	303	5/5	0.97	0.08	34,36,43,43	5
3	SO4	A	302	5/5	0.97	0.08	34,35,40,42	5
3	SO4	A	301	5/5	0.98	0.18	56,58,61,64	0

### 6.5 Other polymers [i](#)

There are no such residues in this entry.