



# wwPDB X-ray Structure Validation Summary Report ⓘ

Aug 15, 2023 – 01:20 AM EDT

PDB ID : 1RLT  
Title : Transition State Analogue of ybiV from E. coli K12  
Authors : Roberts, A.; Lee, S.Y.; McCullagh, E.; Silversmith, R.E.; Wemmer, D.E.  
Deposited on : 2003-11-26  
Resolution : 2.20 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.35  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.35


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	4898 (2.20-2.20)
Clashscore	141614	5594 (2.20-2.20)
Ramachandran outliers	138981	5503 (2.20-2.20)
Sidechain outliers	138945	5504 (2.20-2.20)
RSRZ outliers	127900	4800 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	271	 67% 29% ..
1	B	271	 5% 65% 30% ..
1	C	271	 11% 60% 35% ..
1	D	271	 67% 29% ..

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	GOL	C	759	-	X	-	X
5	GOL	D	758	-	X	X	-

## 2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 8949 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Phosphatase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	268	2129	1366	361	394	8	0	0	0
1	B	268	2129	1366	361	394	8	0	0	0
1	C	268	2129	1366	361	394	8	0	0	0
1	D	268	2129	1366	361	394	8	0	0	0


There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	2	ALA	SER	engineered mutation	UNP P75792
A	267	TYR	SER	engineered mutation	UNP P75792
B	2	ALA	SER	engineered mutation	UNP P75792
B	267	TYR	SER	engineered mutation	UNP P75792
C	2	ALA	SER	engineered mutation	UNP P75792
C	267	TYR	SER	engineered mutation	UNP P75792
D	2	ALA	SER	engineered mutation	UNP P75792
D	267	TYR	SER	engineered mutation	UNP P75792

- Molecule 2 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total 1	Mg 1	0	0
2	B	1	Total 1	Mg 1	0	0
2	C	1	Total 1	Mg 1	0	0
2	D	1	Total 1	Mg 1	0	0

- Molecule 3 is ACETATE ION (three-letter code: ACT) (formula:  $C_2H_3O_2$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			4	2	2		

- Molecule 4 is ALUMINUM FLUORIDE (three-letter code: AF3) (formula:  $AlF_3$ ).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	Al	F	0	0
			4	1	3		
4	B	1	Total	Al	F	0	0
			4	1	3		

*Continued on next page...*

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	Al	F		
4	C	1	4	1	3	0	0
4	D	1	4	1	3	0	0

- Molecule 5 is GLYCEROL (three-letter code: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
5	C	1	6	3	3	0	0
5	D	1	6	3	3	0	0


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
6	A	116	116	116	0	0
6	B	111	111	111	0	0
6	C	76	76	76	0	0
6	D	94	94	94	0	0





- Molecule 1: Phosphatase



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	71.76Å 91.17Å 183.85Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	19.93 – 2.20 20.05 – 2.20	Depositor EDS
% Data completeness (in resolution range)	96.5 (19.93-2.20) 96.5 (20.05-2.20)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.08	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.03 (at 2.19Å)	Xtrriage
Refinement program	CNS 1.1	Depositor
R, $R_{free}$	0.221 , 0.264 0.209 , 0.250	Depositor DCC
$R_{free}$ test set	6223 reflections (10.12%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	31.3	Xtrriage
Anisotropy	0.726	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 47.2	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	8949	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	40.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.17% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ACT, AF3, MG, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.46	0/2174	0.74	3/2939 (0.1%)
1	B	0.48	1/2174 (0.0%)	0.82	3/2939 (0.1%)
1	C	0.45	0/2174	0.81	3/2939 (0.1%)
1	D	0.41	0/2174	0.81	2/2939 (0.1%)
All	All	0.45	1/8696 (0.0%)	0.79	11/11756 (0.1%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	268	PRO	N-CD	-7.01	1.38	1.47

The worst 5 of 11 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
1	D	267	TYR	C-N-CD	-22.70	70.66	120.60
1	B	267	TYR	C-N-CD	-22.55	70.98	120.60
1	C	154	PRO	CA-N-CD	-15.42	89.92	111.50
1	C	267	TYR	C-N-CD	-13.43	91.05	120.60
1	D	267	TYR	C-N-CA	13.05	176.83	122.00

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2129	0	2119	75	0
1	B	2129	0	2119	80	0
1	C	2129	0	2119	103	0
1	D	2129	0	2119	73	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
2	D	1	0	0	0	0
3	A	4	0	3	0	0
4	A	4	0	0	1	0
4	B	4	0	0	1	0
4	C	4	0	0	0	0
4	D	4	0	0	1	0
5	C	6	0	4	1	0
5	D	6	0	4	4	0
6	A	116	0	0	4	0
6	B	111	0	0	3	0
6	C	76	0	0	5	0
6	D	94	0	0	4	0
All	All	8949	0	8487	331	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 19.

The worst 5 of 331 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:153:LEU:HD13	1:C:161:VAL:HG21	1.24	1.09
1:A:84:LEU:HD21	1:A:187:ILE:HD11	1.38	1.06
1:C:245:TYR:CE2	1:C:268:PRO:HB3	1.91	1.05
1:C:153:LEU:HD12	1:C:161:VAL:HG11	1.37	1.02
1:B:236:ALA:HB3	1:B:239:ILE:HD13	1.42	1.02

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	266/271 (98%)	254 (96%)	11 (4%)	1 (0%)	34	37
1	B	266/271 (98%)	248 (93%)	16 (6%)	2 (1%)	19	19
1	C	266/271 (98%)	244 (92%)	19 (7%)	3 (1%)	14	12
1	D	266/271 (98%)	252 (95%)	12 (4%)	2 (1%)	19	19
All	All	1064/1084 (98%)	998 (94%)	58 (6%)	8 (1%)	19	19

5 of 8 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	268	PRO
1	D	268	PRO
1	A	100	LYS
1	C	218	ASN
1	C	159	PRO

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	228/230 (99%)	215 (94%)	13 (6%)	20	24
1	B	228/230 (99%)	218 (96%)	10 (4%)	28	35
1	C	228/230 (99%)	214 (94%)	14 (6%)	18	21
1	D	228/230 (99%)	219 (96%)	9 (4%)	32	41
All	All	912/920 (99%)	866 (95%)	46 (5%)	24	30

5 of 46 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	109	LEU
1	C	225	MET
1	C	147	PHE

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	C	177	THR
1	D	13	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 17 such sidechains are listed below:

Mol	Chain	Res	Type
1	D	101	GLN
1	D	131	HIS
1	B	129	HIS
1	B	208	GLN
1	C	46	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

Of 11 ligands modelled in this entry, 4 are monoatomic - leaving 7 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
5	GOL	C	759	1	5,5,5	5.30	5 (100%)	5,5,5	5.90	3 (60%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	AF3	B	802	2,1,6	0,3,3	-	-	-		
4	AF3	D	804	2,1,6	0,3,3	-	-	-		
4	AF3	A	801	2,1,6	0,3,3	-	-	-		
3	ACT	A	800	-	3,3,3	5.05	1 (33%)	3,3,3	0.76	0
5	GOL	D	758	-	5,5,5	4.56	5 (100%)	5,5,5	5.67	3 (60%)
4	AF3	C	803	2,1,6	0,3,3	-	-	-		

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	GOL	C	759	1	-	3/4/4/4	-
5	GOL	D	758	-	-	2/4/4/4	-

The worst 5 of 11 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	800	ACT	O-C	8.74	1.62	1.22
5	C	759	GOL	C3-C2	-8.63	1.16	1.51
5	D	758	GOL	C3-C2	-8.08	1.18	1.51
5	C	759	GOL	O2-C2	-5.78	1.26	1.43
5	C	759	GOL	O1-C1	4.79	1.62	1.42

The worst 5 of 6 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	C	759	GOL	O3-C3-C2	11.78	166.68	110.20
5	D	758	GOL	O3-C3-C2	10.24	159.29	110.20
5	D	758	GOL	O2-C2-C3	6.70	138.65	109.12
5	C	759	GOL	O2-C2-C3	5.25	132.24	109.12
5	D	758	GOL	O1-C1-C2	3.12	125.14	110.20

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	C	759	GOL	C1-C2-C3-O3
5	D	758	GOL	C1-C2-C3-O3

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms
5	D	758	GOL	O1-C1-C2-C3
5	C	759	GOL	O1-C1-C2-O2
5	C	759	GOL	O2-C2-C3-O3

There are no ring outliers.

5 monomers are involved in 8 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	C	759	GOL	1	0
4	B	802	AF3	1	0
4	D	804	AF3	1	0
4	A	801	AF3	1	0
5	D	758	GOL	4	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	268/271 (98%)	-0.13	4 (1%) 73 72	16, 31, 57, 75	0
1	B	268/271 (98%)	0.18	13 (4%) 29 28	20, 35, 79, 123	0
1	C	268/271 (98%)	0.48	30 (11%) 5 4	24, 41, 98, 144	0
1	D	268/271 (98%)	-0.01	2 (0%) 87 86	21, 35, 61, 87	0
All	All	1072/1084 (98%)	0.13	49 (4%) 32 31	16, 36, 77, 144	0

The worst 5 of 49 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	163	ASP	9.5
1	C	154	PRO	6.6
1	B	180	PHE	5.9
1	C	153	LEU	5.5
1	C	160	LEU	5.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
5	GOL	C	759	6/6	0.74	0.42	43,46,46,48	0
5	GOL	D	758	6/6	0.76	0.22	28,34,35,37	0
2	MG	C	807	1/1	0.81	0.09	39,39,39,39	0
4	AF3	D	804	4/4	0.84	0.18	54,55,55,56	0
4	AF3	C	803	4/4	0.91	0.21	58,58,58,58	0
4	AF3	B	802	4/4	0.91	0.20	51,52,52,52	0
4	AF3	A	801	4/4	0.93	0.26	41,43,44,46	0
3	ACT	A	800	4/4	0.93	0.27	28,30,31,31	0
2	MG	A	805	1/1	0.94	0.11	25,25,25,25	0
2	MG	D	808	1/1	0.95	0.07	32,32,32,32	0
2	MG	B	806	1/1	0.95	0.08	37,37,37,37	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.