



wwPDB X-ray Structure Validation Summary Report ⓘ

May 19, 2020 – 05:05 am BST

PDB ID : 4RS4
Title : Crystal structure and mutational analysis of the endoribonuclease from human coronavirus 229E
Authors : Huo, T.; Liu, X.; Yang, C.; Rao, Z.
Deposited on : 2014-11-06
Resolution : 2.96 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.96 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3104 (3.00-2.92)
Clashscore	141614	3462 (3.00-2.92)
Ramachandran outliers	138981	3340 (3.00-2.92)
Sidechain outliers	138945	3343 (3.00-2.92)
RSRZ outliers	127900	2986 (3.00-2.92)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	349	 4% 73% 23% ..
1	B	349	 % 75% 20% ..
1	C	349	 9% 62% 31% • 5%
1	D	349	 9% 62% 31% ..
1	E	349	 % 68% 28% ..
1	F	349	 % 73% 23% ..

2 Entry composition [i](#)

There is only 1 type of molecule in this entry. The entry contains 15989 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Uridylate-specific endoribonuclease.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	341	2674	1724	432	504	14	0	0	0
1	B	345	2695	1735	437	509	14	0	0	0
1	C	332	2601	1679	418	490	14	0	0	0
1	D	337	2636	1703	424	495	14	0	0	0
1	E	341	2669	1721	430	504	14	0	0	0
1	F	346	2714	1746	439	515	14	0	0	0

There are 30 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	SER	-	EXPRESSION TAG	UNP P0C6X1
A	17	SER	GLY	ENGINEERED MUTATION	UNP P0C6X1
A	142	ALA	THR	ENGINEERED MUTATION	UNP P0C6X1
A	219	MET	ILE	ENGINEERED MUTATION	UNP P0C6X1
A	252	SER	LEU	ENGINEERED MUTATION	UNP P0C6X1
B	0	SER	-	EXPRESSION TAG	UNP P0C6X1
B	17	SER	GLY	ENGINEERED MUTATION	UNP P0C6X1
B	142	ALA	THR	ENGINEERED MUTATION	UNP P0C6X1
B	219	MET	ILE	ENGINEERED MUTATION	UNP P0C6X1
B	252	SER	LEU	ENGINEERED MUTATION	UNP P0C6X1
C	0	SER	-	EXPRESSION TAG	UNP P0C6X1
C	17	SER	GLY	ENGINEERED MUTATION	UNP P0C6X1
C	142	ALA	THR	ENGINEERED MUTATION	UNP P0C6X1
C	219	MET	ILE	ENGINEERED MUTATION	UNP P0C6X1
C	252	SER	LEU	ENGINEERED MUTATION	UNP P0C6X1
D	0	SER	-	EXPRESSION TAG	UNP P0C6X1
D	17	SER	GLY	ENGINEERED MUTATION	UNP P0C6X1

Continued on next page...


Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
D	142	ALA	THR	ENGINEERED MUTATION	UNP P0C6X1
D	219	MET	ILE	ENGINEERED MUTATION	UNP P0C6X1
D	252	SER	LEU	ENGINEERED MUTATION	UNP P0C6X1
E	0	SER	-	EXPRESSION TAG	UNP P0C6X1
E	17	SER	GLY	ENGINEERED MUTATION	UNP P0C6X1
E	142	ALA	THR	ENGINEERED MUTATION	UNP P0C6X1
E	219	MET	ILE	ENGINEERED MUTATION	UNP P0C6X1
E	252	SER	LEU	ENGINEERED MUTATION	UNP P0C6X1
F	0	SER	-	EXPRESSION TAG	UNP P0C6X1
F	17	SER	GLY	ENGINEERED MUTATION	UNP P0C6X1
F	142	ALA	THR	ENGINEERED MUTATION	UNP P0C6X1
F	219	MET	ILE	ENGINEERED MUTATION	UNP P0C6X1
F	252	SER	LEU	ENGINEERED MUTATION	UNP P0C6X1


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Uridylate-specific endoribonuclease




- Molecule 1: Uridylate-specific endoribonuclease




- Molecule 1: Uridylate-specific endoribonuclease







• Molecule 1: Uridylate-specific endoribonuclease



• Molecule 1: Uridylate-specific endoribonuclease



• Molecule 1: Uridylate-specific endoribonuclease





4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	127.93Å 143.17Å 174.66Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	30.00 – 2.96 38.25 – 2.96	Depositor EDS
% Data completeness (in resolution range)	90.4 (30.00-2.96) 89.3 (38.25-2.96)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.59 (at 2.95Å)	Xtrriage
Refinement program	PHENIX (phenix.refine: 1.7_650)	Depositor
R, R_{free}	0.208 , 0.255 0.195 , 0.243	Depositor DCC
R_{free} test set	3387 reflections (5.07%)	wwPDB-VP
Wilson B-factor (Å ²)	56.9	Xtrriage
Anisotropy	0.595	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 45.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	15989	wwPDB-VP
Average B, all atoms (Å ²)	71.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.46% of the height of the origin peak. No significant pseudotranslation is detected.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.31	0/2732	0.49	1/3708 (0.0%)
1	B	0.30	0/2753	0.49	0/3737
1	C	0.28	0/2655	0.46	0/3604
1	D	0.28	0/2692	0.47	0/3653
1	E	0.30	0/2726	0.47	0/3700
1	F	0.29	0/2772	0.47	0/3761
All	All	0.29	0/16330	0.47	1/22163 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	D	0	1
1	F	0	1
All	All	0	2

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	A	109	GLU	N-CA-C	-6.04	94.69	111.00

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	D	142	ALA	Peptide
1	F	142	ALA	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2674	0	2657	58	0
1	B	2695	0	2668	58	0
1	C	2601	0	2568	68	0
1	D	2636	0	2621	74	0
1	E	2669	0	2652	65	0
1	F	2714	0	2691	49	0
All	All	15989	0	15857	358	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

The worst 5 of 358 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:315:VAL:HG11	1:B:318:LYS:HE2	1.42	1.01
1:F:157:LEU:HB2	1:F:162:ILE:HD13	1.58	0.86
1:A:288:PRO:HG3	1:D:162:ILE:HD11	1.57	0.85
1:D:334:TRP:HZ3	1:D:344:TYR:HE2	1.24	0.84
1:A:164:THR:HG23	1:A:174:ASN:HA	1.61	0.82

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	337/349 (97%)	325 (96%)	12 (4%)	0	100 100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	341/349 (98%)	328 (96%)	12 (4%)	1 (0%)	41	73
1	C	322/349 (92%)	307 (95%)	14 (4%)	1 (0%)	41	73
1	D	331/349 (95%)	311 (94%)	18 (5%)	2 (1%)	25	60
1	E	337/349 (97%)	321 (95%)	16 (5%)	0	100	100
1	F	342/349 (98%)	332 (97%)	10 (3%)	0	100	100
All	All	2010/2094 (96%)	1924 (96%)	82 (4%)	4 (0%)	47	79

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	236	VAL
1	D	237	VAL
1	D	190	HIS
1	C	246	LEU

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	298/306 (97%)	281 (94%)	17 (6%)	20	52
1	B	299/306 (98%)	277 (93%)	22 (7%)	13	41
1	C	290/306 (95%)	263 (91%)	27 (9%)	9	30
1	D	293/306 (96%)	266 (91%)	27 (9%)	9	30
1	E	297/306 (97%)	275 (93%)	22 (7%)	13	41
1	F	303/306 (99%)	283 (93%)	20 (7%)	16	46
All	All	1780/1836 (97%)	1645 (92%)	135 (8%)	13	39

5 of 135 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	315	VAL
1	D	162	ILE

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	F	235	HIS
1	C	324	ILE
1	D	84	VAL

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 12 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	326	ASN
1	D	117	ASN
1	F	171	ASN
1	C	232	ASN
1	E	232	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	341/349 (97%)	0.11	14 (4%) 37 24	32, 68, 115, 153	0
1	B	345/349 (98%)	-0.04	4 (1%) 79 63	33, 56, 100, 137	0
1	C	332/349 (95%)	0.44	32 (9%) 8 4	45, 80, 125, 144	0
1	D	337/349 (96%)	0.49	30 (8%) 9 5	37, 85, 145, 171	0
1	E	341/349 (97%)	0.02	4 (1%) 79 63	39, 62, 99, 136	0
1	F	346/349 (99%)	-0.02	2 (0%) 89 78	35, 52, 93, 143	0
All	All	2042/2094 (97%)	0.17	86 (4%) 36 23	32, 67, 118, 171	0

The worst 5 of 86 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	144	GLY	6.9
1	D	317	SER	6.7
1	D	316	VAL	6.7
1	D	314	THR	5.4
1	D	319	VAL	4.8

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers

There are no such residues in this entry.