



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 17, 2024 – 11:28 AM EST

PDB ID : 3SJA
Title : Crystal structure of *S. cerevisiae* Get3 in the open state in complex with Get1 cytosolic domain
Authors : Reitz, S.; Wild, K.; Sinning, I.
Deposited on : 2011-06-21
Resolution : 3.00 Å (reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2092 (3.00-3.00)
Clashscore	141614	2416 (3.00-3.00)
Ramachandran outliers	138981	2333 (3.00-3.00)
Sidechain outliers	138945	2336 (3.00-3.00)
RSRZ outliers	127900	1990 (3.00-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	362	
1	B	362	
1	E	362	
1	F	362	
1	I	362	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	C	65	 78% 14% 6%
2	D	65	 82% 11% 6%
2	G	65	 85% 9% 6%
2	H	65	 83% 11% 6%
2	J	65	 85% 9% 6%

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 15873 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ATPase GET3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	338	Total 2643	C 1667	N 441	O 516	S 19	0	0	0
1	B	341	Total 2663	C 1678	N 444	O 522	S 19	0	0	0
1	E	338	Total 2639	C 1665	N 441	O 514	S 19	0	0	0
1	F	341	Total 2663	C 1678	N 444	O 522	S 19	0	0	0
1	I	339	Total 2647	C 1669	N 442	O 517	S 19	0	0	0

There are 40 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	355	LEU	-	expression tag	UNP Q12154
A	356	GLU	-	expression tag	UNP Q12154
A	357	HIS	-	expression tag	UNP Q12154
A	358	HIS	-	expression tag	UNP Q12154
A	359	HIS	-	expression tag	UNP Q12154
A	360	HIS	-	expression tag	UNP Q12154
A	361	HIS	-	expression tag	UNP Q12154
A	362	HIS	-	expression tag	UNP Q12154
B	355	LEU	-	expression tag	UNP Q12154
B	356	GLU	-	expression tag	UNP Q12154
B	357	HIS	-	expression tag	UNP Q12154
B	358	HIS	-	expression tag	UNP Q12154
B	359	HIS	-	expression tag	UNP Q12154
B	360	HIS	-	expression tag	UNP Q12154
B	361	HIS	-	expression tag	UNP Q12154
B	362	HIS	-	expression tag	UNP Q12154
E	355	LEU	-	expression tag	UNP Q12154
E	356	GLU	-	expression tag	UNP Q12154
E	357	HIS	-	expression tag	UNP Q12154

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
E	358	HIS	-	expression tag	UNP Q12154
E	359	HIS	-	expression tag	UNP Q12154
E	360	HIS	-	expression tag	UNP Q12154
E	361	HIS	-	expression tag	UNP Q12154
E	362	HIS	-	expression tag	UNP Q12154
F	355	LEU	-	expression tag	UNP Q12154
F	356	GLU	-	expression tag	UNP Q12154
F	357	HIS	-	expression tag	UNP Q12154
F	358	HIS	-	expression tag	UNP Q12154
F	359	HIS	-	expression tag	UNP Q12154
F	360	HIS	-	expression tag	UNP Q12154
F	361	HIS	-	expression tag	UNP Q12154
F	362	HIS	-	expression tag	UNP Q12154
I	355	LEU	-	expression tag	UNP Q12154
I	356	GLU	-	expression tag	UNP Q12154
I	357	HIS	-	expression tag	UNP Q12154
I	358	HIS	-	expression tag	UNP Q12154
I	359	HIS	-	expression tag	UNP Q12154
I	360	HIS	-	expression tag	UNP Q12154
I	361	HIS	-	expression tag	UNP Q12154
I	362	HIS	-	expression tag	UNP Q12154

- Molecule 2 is a protein called Golgi to ER traffic protein 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	61	Total	C	N	O	S	0	0	0
			515	317	95	102	1			
2	D	61	Total	C	N	O	S	0	0	0
			515	317	95	102	1			
2	G	61	Total	C	N	O	S	0	0	0
			515	317	95	102	1			
2	H	61	Total	C	N	O	S	0	0	0
			515	317	95	102	1			
2	J	61	Total	C	N	O	S	0	0	0
			515	317	95	102	1			

There are 35 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	35	MET	-	expression tag	UNP P53192
C	94	HIS	-	expression tag	UNP P53192
C	95	HIS	-	expression tag	UNP P53192

Continued on next page...


Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	96	HIS	-	expression tag	UNP P53192
C	97	HIS	-	expression tag	UNP P53192
C	98	HIS	-	expression tag	UNP P53192
C	99	HIS	-	expression tag	UNP P53192
D	35	MET	-	expression tag	UNP P53192
D	94	HIS	-	expression tag	UNP P53192
D	95	HIS	-	expression tag	UNP P53192
D	96	HIS	-	expression tag	UNP P53192
D	97	HIS	-	expression tag	UNP P53192
D	98	HIS	-	expression tag	UNP P53192
D	99	HIS	-	expression tag	UNP P53192
G	35	MET	-	expression tag	UNP P53192
G	94	HIS	-	expression tag	UNP P53192
G	95	HIS	-	expression tag	UNP P53192
G	96	HIS	-	expression tag	UNP P53192
G	97	HIS	-	expression tag	UNP P53192
G	98	HIS	-	expression tag	UNP P53192
G	99	HIS	-	expression tag	UNP P53192
H	35	MET	-	expression tag	UNP P53192
H	94	HIS	-	expression tag	UNP P53192
H	95	HIS	-	expression tag	UNP P53192
H	96	HIS	-	expression tag	UNP P53192
H	97	HIS	-	expression tag	UNP P53192
H	98	HIS	-	expression tag	UNP P53192
H	99	HIS	-	expression tag	UNP P53192
J	35	MET	-	expression tag	UNP P53192
J	94	HIS	-	expression tag	UNP P53192
J	95	HIS	-	expression tag	UNP P53192
J	96	HIS	-	expression tag	UNP P53192
J	97	HIS	-	expression tag	UNP P53192
J	98	HIS	-	expression tag	UNP P53192
J	99	HIS	-	expression tag	UNP P53192

- Molecule 3 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total Zn 1 1	0	0
3	E	1	Total Zn 1 1	0	0
3	I	1	Total Zn 1 1	0	0


- Molecule 4 is PHOSPHATE ION (three-letter code: PO4) (formula: O₄P).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total O P 5 4 1	0	0
4	A	1	Total O P 5 4 1	0	0
4	B	1	Total O P 5 4 1	0	0
4	D	1	Total O P 5 4 1	0	0
4	E	1	Total O P 5 4 1	0	0
4	F	1	Total O P 5 4 1	0	0
4	I	1	Total O P 5 4 1	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total O 1 1	0	0
5	B	1	Total O 1 1	0	0
5	F	2	Total O 2 2	0	0
5	I	1	Total O 1 1	0	0




• Molecule 1: ATPase GET3

Chain F: 65% 25% 6%



• Molecule 1: ATPase GET3

Chain I: 65% 25% 6%




• Molecule 2: Golgi to ER traffic protein 1

Chain C: 78% 14% 6%




- Molecule 2: Golgi to ER traffic protein 1




- Molecule 2: Golgi to ER traffic protein 1



- Molecule 2: Golgi to ER traffic protein 1



- Molecule 2: Golgi to ER traffic protein 1



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	204.12Å 91.46Å 149.18Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	51.30 – 3.00 51.27 – 3.00	Depositor EDS
% Data completeness (in resolution range)	99.4 (51.30-3.00) 99.5 (51.27-3.00)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.12	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.52 (at 3.01Å)	Xtrriage
Refinement program	PHENIX (phenix.refine: 1.6_289)	Depositor
R, R_{free}	0.181 , 0.231 0.173 , 0.222	Depositor DCC
R_{free} test set	2857 reflections (5.06%)	wwPDB-VP
Wilson B-factor (Å ²)	76.6	Xtrriage
Anisotropy	0.087	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 73.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	15873	wwPDB-VP
Average B, all atoms (Å ²)	77.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.43% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, PO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.44	0/2686	0.61	0/3621
1	B	0.46	0/2706	0.60	0/3649
1	E	0.45	0/2682	0.62	0/3616
1	F	0.40	0/2706	0.56	0/3649
1	I	0.40	0/2690	0.57	0/3627
2	C	0.40	0/522	0.49	0/696
2	D	0.40	0/522	0.48	0/696
2	G	0.39	0/522	0.48	0/696
2	H	0.38	0/522	0.48	0/696
2	J	0.37	0/522	0.50	0/696
All	All	0.43	0/16080	0.58	0/21642

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	2
1	F	0	1
All	All	0	3

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	26	LYS	Peptide

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Group
1	B	284	ASN	Peptide
1	F	284	ASN	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2643	0	2628	81	0
1	B	2663	0	2645	79	0
1	E	2639	0	2627	80	0
1	F	2663	0	2644	67	0
1	I	2647	0	2631	74	0
2	C	515	0	505	4	0
2	D	515	0	505	3	0
2	G	515	0	505	1	0
2	H	515	0	505	2	0
2	J	515	0	505	1	0
3	A	1	0	0	0	0
3	E	1	0	0	0	0
3	I	1	0	0	0	0
4	A	10	0	0	0	0
4	B	5	0	0	0	0
4	D	5	0	0	0	0
4	E	5	0	0	0	0
4	F	5	0	0	1	0
4	I	5	0	0	0	0
5	A	1	0	0	0	0
5	B	1	0	0	0	0
5	F	2	0	0	0	0
5	I	1	0	0	0	0
All	All	15873	0	15700	378	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

The worst 5 of 378 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:352:ASP:HB3	1:I:69:LYS:HB3	1.36	1.03
1:A:9:LEU:O	1:A:13:ILE:HG13	1.68	0.94
1:A:230:PHE:HA	1:A:236:THR:HG21	1.49	0.93
1:A:239:VAL:HG13	1:A:268:SER:HB2	1.53	0.89
1:I:281:GLN:HA	1:I:283:HIS:ND1	1.87	0.89

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	334/362 (92%)	317 (95%)	16 (5%)	1 (0%)	41	76
1	B	337/362 (93%)	322 (96%)	14 (4%)	1 (0%)	41	76
1	E	334/362 (92%)	321 (96%)	10 (3%)	3 (1%)	17	55
1	F	337/362 (93%)	321 (95%)	12 (4%)	4 (1%)	13	48
1	I	335/362 (92%)	318 (95%)	14 (4%)	3 (1%)	17	55
2	C	59/65 (91%)	56 (95%)	3 (5%)	0	100	100
2	D	59/65 (91%)	57 (97%)	2 (3%)	0	100	100
2	G	59/65 (91%)	57 (97%)	2 (3%)	0	100	100
2	H	59/65 (91%)	58 (98%)	1 (2%)	0	100	100
2	J	59/65 (91%)	56 (95%)	3 (5%)	0	100	100
All	All	1972/2135 (92%)	1883 (96%)	77 (4%)	12 (1%)	25	64

5 of 12 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	285	CYS
1	F	173	THR
1	F	285	CYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	F	318	ALA
1	I	173	THR

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	297/317 (94%)	270 (91%)	27 (9%)	9 34
1	B	299/317 (94%)	274 (92%)	25 (8%)	11 38
1	E	296/317 (93%)	272 (92%)	24 (8%)	11 40
1	F	299/317 (94%)	266 (89%)	33 (11%)	6 25
1	I	297/317 (94%)	266 (90%)	31 (10%)	7 27
2	C	58/62 (94%)	54 (93%)	4 (7%)	15 48
2	D	58/62 (94%)	54 (93%)	4 (7%)	15 48
2	G	58/62 (94%)	54 (93%)	4 (7%)	15 48
2	H	58/62 (94%)	55 (95%)	3 (5%)	23 59
2	J	58/62 (94%)	54 (93%)	4 (7%)	15 48
All	All	1778/1895 (94%)	1619 (91%)	159 (9%)	9 35

5 of 159 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	F	341	ILE
1	I	170	THR
2	G	47	LYS
1	I	95	LYS
1	I	268	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 36 such sidechains are listed below:

Mol	Chain	Res	Type
1	I	122	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	J	62	GLN
1	I	178	GLN
1	I	267	ASN
1	E	18	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 10 ligands modelled in this entry, 3 are monoatomic - leaving 7 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	PO4	A	364	-	4,4,4	0.92	0	6,6,6	0.61	0
4	PO4	D	7	-	4,4,4	0.96	0	6,6,6	0.66	0
4	PO4	I	364	-	4,4,4	1.00	0	6,6,6	0.54	0
4	PO4	F	363	-	4,4,4	0.81	0	6,6,6	0.65	0
4	PO4	A	365	-	4,4,4	0.95	0	6,6,6	0.98	0
4	PO4	B	363	-	4,4,4	0.76	0	6,6,6	0.91	0
4	PO4	E	364	-	4,4,4	0.98	0	6,6,6	0.83	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	F	363	PO4	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	338/362 (93%)	-0.47	1 (0%) 94 84	36, 70, 124, 198	0
1	B	341/362 (94%)	-0.35	2 (0%) 89 72	35, 68, 118, 168	0
1	E	338/362 (93%)	-0.35	1 (0%) 94 84	41, 70, 120, 214	0
1	F	341/362 (94%)	-0.37	0 100 100	41, 75, 117, 154	0
1	I	339/362 (93%)	-0.25	5 (1%) 73 46	43, 76, 126, 249	0
2	C	61/65 (93%)	-0.55	0 100 100	42, 69, 106, 134	0
2	D	61/65 (93%)	-0.41	0 100 100	47, 70, 105, 135	0
2	G	61/65 (93%)	-0.49	0 100 100	46, 71, 108, 137	0
2	H	61/65 (93%)	-0.49	0 100 100	49, 73, 107, 133	0
2	J	61/65 (93%)	-0.37	0 100 100	47, 71, 105, 133	0
All	All	2002/2135 (93%)	-0.37	9 (0%) 92 79	35, 72, 119, 249	0

The worst 5 of 9 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	I	347	ILE	3.4
1	I	281	GLN	3.2
1	I	348	TYR	3.0
1	I	284	ASN	2.5
1	A	352	ASP	2.4

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	PO4	A	365	5/5	0.82	0.31	111,163,169,181	0
4	PO4	B	363	5/5	0.86	0.24	94,120,148,162	0
4	PO4	F	363	5/5	0.86	0.23	117,125,165,167	0
4	PO4	E	364	5/5	0.87	0.26	112,142,147,153	0
4	PO4	I	364	5/5	0.90	0.16	133,136,154,167	0
4	PO4	D	7	5/5	0.92	0.37	121,126,132,137	0
4	PO4	A	364	5/5	0.96	0.23	111,115,127,129	0
3	ZN	I	363	1/1	0.98	0.04	107,107,107,107	1
3	ZN	A	363	1/1	0.99	0.15	79,79,79,79	0
3	ZN	E	363	1/1	0.99	0.15	82,82,82,82	0

6.5 Other polymers [i](#)

There are no such residues in this entry.