



# wwPDB X-ray Structure Validation Summary Report ⓘ

Oct 9, 2023 – 02:43 PM EDT

PDB ID : 7TSI  
Title : Structure of human endothelial nitric oxide synthase heme domain in complex with 4-methyl-6-(3-((methylamino)methyl)phenyl)pyridin-2-amine  
Authors : Li, H.; Poulos, T.L.  
Deposited on : 2022-01-31  
Resolution : 2.10 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.35.1  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.35.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	5197 (2.10-2.10)
Clashscore	141614	5710 (2.10-2.10)
Ramachandran outliers	138981	5647 (2.10-2.10)
Sidechain outliers	138945	5648 (2.10-2.10)
RSRZ outliers	127900	5083 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	440	 6% 65% 23% 9%
1	B	440	 72% 18% 9%
1	C	440	 4% 65% 25% 9%
1	D	440	 72% 18% 9%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
6	GOL	C	508	-	X	-	-

## 2 Entry composition

There are 10 unique types of molecules in this entry. The entry contains 13853 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Nitric oxide synthase, endothelial.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	401	3207	2043	564	584	16	0	1	0
1	B	401	3208	2043	564	585	16	0	2	0
1	C	402	3212	2046	565	585	16	0	1	0
1	D	402	3214	2046	567	585	16	0	1	0

There are 4 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	298	GLU	ASP	variant	UNP P29474
B	298	GLU	ASP	variant	UNP P29474
C	298	GLU	ASP	variant	UNP P29474
D	298	GLU	ASP	variant	UNP P29474

- Molecule 2 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula:  $C_{34}H_{32}FeN_4O_4$ ).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			17	9	5	3		
3	B	1	Total	C	N	O	0	0
			17	9	5	3		
3	C	1	Total	C	N	O	0	0
			17	9	5	3		
3	D	1	Total	C	N	O	0	0
			17	9	5	3		

- Molecule 4 is 4-methyl-6-{3-[(methylamino)methyl]phenyl}pyridin-2-amine (three-letter code: KMI) (formula: C<sub>14</sub>H<sub>17</sub>N<sub>3</sub>) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	N	0	0
			17	14	3		
4	B	1	Total	C	N	0	0
			17	14	3		
4	C	1	Total	C	N	0	0
			17	14	3		
4	D	1	Total	C	N	0	0
			17	14	3		

- Molecule 5 is 2-[BIS-(2-HYDROXY-ETHYL)-AMINO]-2-HYDROXYMETHYL-PROPAN E-1,3-DIOL (three-letter code: BTB) (formula: C<sub>8</sub>H<sub>19</sub>NO<sub>5</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	A	1	Total	C	N	O	0	0
			14	8	1	5		
5	A	1	Total	C	N	O	0	0
			14	8	1	5		
5	A	1	Total	C	N	O	0	0
			14	8	1	5		
5	B	1	Total	C	N	O	0	0
			14	8	1	5		
5	B	1	Total	C	N	O	0	0
			14	8	1	5		
5	C	1	Total	C	N	O	0	0
			14	8	1	5		
5	C	1	Total	C	N	O	0	0
			14	8	1	5		
5	C	1	Total	C	N	O	0	0
			14	8	1	5		
5	D	1	Total	C	N	O	0	0
			14	8	1	5		
5	D	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 6 is GLYCEROL (three-letter code: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	1	Total C O 6 3 3	0	0
6	A	1	Total C O 6 3 3	0	0
6	A	1	Total C O 6 3 3	0	0
6	B	1	Total C O 6 3 3	0	0
6	B	1	Total C O 6 3 3	0	0
6	C	1	Total C O 6 3 3	0	0
6	C	1	Total C O 6 3 3	0	0
6	C	1	Total C O 6 3 3	0	0
6	C	1	Total C O 6 3 3	0	0
6	D	1	Total C O 6 3 3	0	0
6	D	1	Total C O 6 3 3	0	0

- Molecule 7 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	1	Total Cl 1 1	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	B	1	Total Cl 1 1	0	0
7	C	1	Total Cl 1 1	0	0
7	D	1	Total Cl 1 1	0	0

- Molecule 8 is GADOLINIUM ATOM (three-letter code: Gd) (formula: Gd).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
8	A	1	Total Gd 1 1	0	0
8	B	1	Total Gd 1 1	0	0
8	C	1	Total Gd 1 1	0	0
8	D	1	Total Gd 1 1	0	0

- Molecule 9 is ZINC ION (three-letter code: ZN) (formula: Zn).


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
9	A	1	Total Zn 1 1	0	0
9	C	1	Total Zn 1 1	0	0

- Molecule 10 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
10	A	81	Total O 81 81	0	0
10	B	154	Total O 154 154	0	0
10	C	92	Total O 92 92	0	0
10	D	161	Total O 161 161	0	0



- Molecule 1: Nitric oxide synthase, endothelial



- Molecule 1: Nitric oxide synthase, endothelial



## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	58.92Å 151.39Å 107.48Å 90.00° 90.62° 90.00°	Depositor
Resolution (Å)	49.11 – 2.10 49.10 – 2.10	Depositor EDS
% Data completeness (in resolution range)	98.1 (49.11-2.10) 95.7 (49.10-2.10)	Depositor EDS
$R_{merge}$	0.18	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.16 (at 2.10Å)	Xtrriage
Refinement program	PHENIX 1.11.1_2575	Depositor
R, $R_{free}$	0.217 , 0.277 0.212 , 0.273	Depositor DCC
$R_{free}$ test set	5403 reflections (4.99%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	34.0	Xtrriage
Anisotropy	0.872	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 37.1	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	0.239 for h,-k,-l	Xtrriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	13853	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	53.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.39% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: KMI, GD, ZN, GOL, CL, BTB, H4B, HEM

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.38	0/3302	0.54	0/4498
1	B	0.43	0/3306	0.57	0/4506
1	C	0.37	0/3307	0.53	0/4506
1	D	0.45	0/3309	0.59	0/4509
All	All	0.41	0/13224	0.56	0/18019

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3207	0	3112	73	1
1	B	3208	0	3109	56	0
1	C	3212	0	3116	76	0
1	D	3214	0	3116	52	0
2	A	43	0	30	3	0
2	B	43	0	30	3	0
2	C	43	0	30	2	0
2	D	43	0	30	1	0
3	A	17	0	15	2	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	B	17	0	15	1	0
3	C	17	0	15	1	0
3	D	17	0	15	1	0
4	A	17	0	0	1	0
4	B	17	0	0	0	0
4	C	17	0	0	1	0
4	D	17	0	0	1	0
5	A	42	0	57	5	0
5	B	28	0	37	7	0
5	C	42	0	55	6	0
5	D	28	0	36	6	1
6	A	18	0	24	0	0
6	B	12	0	16	0	0
6	C	24	0	32	1	0
6	D	12	0	16	0	0
7	A	1	0	0	0	0
7	B	1	0	0	0	0
7	C	1	0	0	0	0
7	D	1	0	0	1	0
8	A	1	0	0	0	0
8	B	1	0	0	0	0
8	C	1	0	0	0	0
8	D	1	0	0	0	0
9	A	1	0	0	0	0
9	C	1	0	0	0	0
10	A	81	0	0	3	0
10	B	154	0	0	5	0
10	C	92	0	0	6	0
10	D	161	0	0	5	0
All	All	13853	0	12906	261	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

The worst 5 of 261 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:239:GLY:HA3	1:A:297:ASP:H	1.46	0.79
1:B:247:GLN:HB2	1:B:250:ARG:HD3	1.64	0.79
1:C:382:CYS:HA	5:C:504:BTB:H12	1.63	0.79
1:D:124:LEU:HD11	1:D:154:GLU:HG3	1.67	0.75

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:200:ASP:OD1	1:C:200:ASP:N	2.20	0.74

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:152:GLU:OE2	5:D:505:BTB:O4[2_851]	1.94	0.26

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	398/440 (90%)	366 (92%)	29 (7%)	3 (1%)	19	15
1	B	399/440 (91%)	377 (94%)	22 (6%)	0	100	100
1	C	399/440 (91%)	378 (95%)	17 (4%)	4 (1%)	15	11
1	D	399/440 (91%)	389 (98%)	9 (2%)	1 (0%)	41	41
All	All	1595/1760 (91%)	1510 (95%)	77 (5%)	8 (0%)	29	26

5 of 8 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	277	HIS
1	C	89	GLN
1	A	203	SER
1	A	257	GLN
1	C	276	GLN

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	342/373 (92%)	322 (94%)	20 (6%)	20	17
1	B	342/373 (92%)	327 (96%)	15 (4%)	28	28
1	C	342/373 (92%)	326 (95%)	16 (5%)	26	25
1	D	342/373 (92%)	327 (96%)	15 (4%)	28	28
All	All	1368/1492 (92%)	1302 (95%)	66 (5%)	25	24

5 of 66 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	136	SER
1	D	200	ASP
1	D	436	LYS
1	B	136	SER
1	B	128	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	233	GLN
1	C	132	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 43 ligands modelled in this entry, 10 are monoatomic - leaving 33 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
6	GOL	A	507	-	5,5,5	0.36	0	5,5,5	0.44	0
6	GOL	C	510	-	5,5,5	0.34	0	5,5,5	0.24	0
3	H4B	B	502	-	16,18,18	0.91	0	11,26,26	2.71	6 (54%)
6	GOL	B	506	-	5,5,5	0.34	0	5,5,5	0.37	0
3	H4B	C	502	-	16,18,18	0.93	0	11,26,26	2.63	4 (36%)
2	HEM	D	501	1	41,50,50	1.53	6 (14%)	45,82,82	1.84	13 (28%)
5	BTB	A	506	-	13,13,13	0.42	0	7,16,16	0.63	0
6	GOL	A	509	-	5,5,5	0.40	0	5,5,5	0.33	0
5	BTB	C	504	8	13,13,13	0.48	0	7,16,16	1.17	1 (14%)
5	BTB	C	505	-	13,13,13	0.62	0	7,16,16	1.02	1 (14%)
6	GOL	D	506	-	5,5,5	0.46	0	5,5,5	0.44	0
3	H4B	D	502	-	16,18,18	0.95	1 (6%)	11,26,26	2.64	4 (36%)
5	BTB	C	506	-	13,13,13	0.42	0	7,16,16	0.44	0
4	KMI	A	503	-	18,18,18	0.43	0	23,24,24	1.63	4 (17%)
5	BTB	B	504	8	13,13,13	0.43	0	7,16,16	0.55	0
6	GOL	B	507	-	5,5,5	0.36	0	5,5,5	0.32	0
5	BTB	A	504	8	13,13,13	0.33	0	7,16,16	0.69	0
5	BTB	D	505	-	13,13,13	0.65	0	7,16,16	0.99	0
2	HEM	B	501	1	41,50,50	1.43	5 (12%)	45,82,82	1.95	15 (33%)
6	GOL	C	509	-	5,5,5	0.36	0	5,5,5	0.27	0
2	HEM	C	501	1	41,50,50	1.53	7 (17%)	45,82,82	1.74	8 (17%)
3	H4B	A	502	-	16,18,18	1.02	0	11,26,26	2.62	5 (45%)
5	BTB	A	505	-	13,13,13	0.73	1 (7%)	7,16,16	1.19	1 (14%)
6	GOL	C	507	-	5,5,5	0.41	0	5,5,5	0.17	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	HEM	A	501	1	41,50,50	1.57	4 (9%)	45,82,82	1.55	9 (20%)
6	GOL	D	507	-	5,5,5	0.36	0	5,5,5	0.32	0
6	GOL	C	508	-	5,5,5	0.77	0	5,5,5	1.80	2 (40%)
5	BTB	B	505	-	13,13,13	0.56	0	7,16,16	1.11	0
5	BTB	D	504	8	13,13,13	0.54	0	7,16,16	0.65	0
4	KMI	D	503	-	18,18,18	0.59	0	23,24,24	1.41	3 (13%)
4	KMI	B	503	-	18,18,18	0.68	1 (5%)	23,24,24	1.17	3 (13%)
4	KMI	C	503	-	18,18,18	0.51	0	23,24,24	1.85	4 (17%)
6	GOL	A	508	-	5,5,5	0.60	0	5,5,5	1.18	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	GOL	A	507	-	-	2/4/4/4	-
6	GOL	C	510	-	-	1/4/4/4	-
3	H4B	B	502	-	-	0/8/17/17	0/2/2/2
6	GOL	B	506	-	-	2/4/4/4	-
3	H4B	C	502	-	-	0/8/17/17	0/2/2/2
2	HEM	D	501	1	-	0/12/54/54	-
5	BTB	A	506	-	-	2/21/21/21	-
6	GOL	A	509	-	-	2/4/4/4	-
5	BTB	C	504	8	-	6/21/21/21	-
5	BTB	C	505	-	-	11/21/21/21	-
6	GOL	D	506	-	-	3/4/4/4	-
3	H4B	D	502	-	-	0/8/17/17	0/2/2/2
5	BTB	C	506	-	-	14/21/21/21	-
4	KMI	A	503	-	-	0/6/7/7	0/2/2/2
5	BTB	B	504	8	-	3/21/21/21	-
6	GOL	B	507	-	-	2/4/4/4	-
5	BTB	A	504	8	-	8/21/21/21	-
5	BTB	D	505	-	-	8/21/21/21	-
2	HEM	B	501	1	-	2/12/54/54	-
6	GOL	C	509	-	-	2/4/4/4	-
2	HEM	C	501	1	-	2/12/54/54	-
3	H4B	A	502	-	-	3/8/17/17	0/2/2/2

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	BTB	A	505	-	-	12/21/21/21	-
6	GOL	C	507	-	-	4/4/4/4	-
2	HEM	A	501	1	-	2/12/54/54	-
6	GOL	D	507	-	-	2/4/4/4	-
6	GOL	C	508	-	-	4/4/4/4	-
5	BTB	B	505	-	-	11/21/21/21	-
5	BTB	D	504	8	-	6/21/21/21	-
4	KMI	D	503	-	-	4/6/7/7	0/2/2/2
4	KMI	B	503	-	-	0/6/7/7	0/2/2/2
4	KMI	C	503	-	-	0/6/7/7	0/2/2/2
6	GOL	A	508	-	-	4/4/4/4	-

The worst 5 of 25 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	501	HEM	C3C-C2C	-4.40	1.34	1.40
2	A	501	HEM	C3C-C2C	-4.20	1.34	1.40
2	C	501	HEM	C3C-CAC	3.71	1.55	1.47
2	A	501	HEM	C3C-CAC	3.62	1.55	1.47
2	B	501	HEM	C3C-CAC	3.42	1.54	1.47

The worst 5 of 83 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	C	503	KMI	C06-N01-C02	6.68	122.76	118.43
4	A	503	KMI	C06-N01-C02	6.03	122.33	118.43
3	C	502	H4B	C8A-C4A-C4	6.00	119.90	114.57
3	D	502	H4B	C8A-C4A-C4	5.89	119.81	114.57
3	A	502	H4B	C8A-C4A-C4	5.86	119.78	114.57

There are no chirality outliers.

5 of 122 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
5	A	504	BTB	C1-C2-C3-O3
5	A	504	BTB	C4-C2-C3-O3
5	A	504	BTB	N-C2-C3-O3
5	A	504	BTB	C1-C2-C4-O4
5	A	504	BTB	C3-C2-C4-O4


There are no ring outliers.


20 monomers are involved in 43 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	502	H4B	1	0
3	C	502	H4B	1	0
2	D	501	HEM	1	0
5	A	506	BTB	3	0
5	C	504	BTB	5	0
3	D	502	H4B	1	0
5	C	506	BTB	1	0
4	A	503	KMI	1	0
5	B	504	BTB	2	0
5	D	505	BTB	3	1
2	B	501	HEM	3	0
2	C	501	HEM	2	0
3	A	502	H4B	2	0
5	A	505	BTB	2	0
6	C	507	GOL	1	0
2	A	501	HEM	3	0
5	B	505	BTB	5	0
5	D	504	BTB	3	0
4	D	503	KMI	1	0
4	C	503	KMI	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.


















## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	401/440 (91%)	0.51	28 (6%) 16 20	28, 59, 107, 156	0
1	B	401/440 (91%)	0.09	2 (0%) 91 92	26, 41, 81, 115	0
1	C	402/440 (91%)	0.31	18 (4%) 33 38	29, 56, 99, 129	0
1	D	402/440 (91%)	0.01	5 (1%) 79 82	25, 40, 66, 118	0
All	All	1606/1760 (91%)	0.23	53 (3%) 46 53	25, 48, 96, 156	0

The worst 5 of 53 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	238	ARG	6.3
1	A	153	VAL	5.8
1	A	254	TYR	4.4
1	A	204	ALA	4.2
1	A	275	ILE	4.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)


There are no monosaccharides in this entry.


### 6.4 Ligands [i](#)


In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
6	GOL	D	507	6/6	0.73	0.15	65,71,78,79	0
6	GOL	C	509	6/6	0.78	0.16	51,53,58,61	0
8	GD	A	511	1/1	0.82	0.12	138,138,138,138	0
6	GOL	D	506	6/6	0.83	0.17	54,67,76,88	0
5	BTB	C	506	14/14	0.83	0.12	71,83,87,88	0
5	BTB	A	506	14/14	0.83	0.12	73,88,94,95	0
6	GOL	B	507	6/6	0.84	0.11	64,69,73,73	0
6	GOL	C	510	6/6	0.85	0.11	60,68,69,72	0
6	GOL	A	509	6/6	0.85	0.12	56,62,63,66	0
5	BTB	A	505	14/14	0.87	0.19	45,64,73,77	0
3	H4B	C	502	17/17	0.89	0.21	59,66,70,70	0
3	H4B	A	502	17/17	0.90	0.17	49,62,76,76	0
5	BTB	B	505	14/14	0.90	0.18	40,69,72,77	0
5	BTB	C	504	14/14	0.91	0.15	48,62,73,74	0
6	GOL	B	506	6/6	0.91	0.17	53,61,67,69	0
5	BTB	D	504	14/14	0.91	0.16	38,64,73,78	0
5	BTB	D	505	14/14	0.91	0.20	47,65,79,82	0
5	BTB	B	504	14/14	0.92	0.14	20,51,65,66	0
3	H4B	D	502	17/17	0.92	0.16	41,56,66,69	0
5	BTB	A	504	14/14	0.92	0.16	55,77,92,96	0
6	GOL	A	507	6/6	0.92	0.20	57,59,69,70	0
4	KMI	C	503	17/17	0.93	0.22	35,50,76,77	0
5	BTB	C	505	14/14	0.94	0.20	25,58,68,72	0
4	KMI	A	503	17/17	0.94	0.16	35,47,68,72	0
6	GOL	C	507	6/6	0.94	0.21	47,50,56,67	0
3	H4B	B	502	17/17	0.94	0.12	41,48,59,61	0
6	GOL	C	508	6/6	0.95	0.21	25,29,51,56	0
4	KMI	D	503	17/17	0.95	0.12	17,32,61,64	0
4	KMI	B	503	17/17	0.95	0.14	22,33,57,57	0
2	HEM	A	501	43/43	0.96	0.18	38,57,74,78	0
2	HEM	C	501	43/43	0.97	0.12	26,45,61,73	0
6	GOL	A	508	6/6	0.97	0.16	31,51,68,70	0
2	HEM	D	501	43/43	0.97	0.13	19,35,49,59	0
7	CL	A	510	1/1	0.98	0.07	46,46,46,46	0
2	HEM	B	501	43/43	0.98	0.13	16,33,50,61	0
8	GD	C	512	1/1	0.98	0.10	80,80,80,80	1
7	CL	C	511	1/1	0.99	0.09	44,44,44,44	0
7	CL	D	508	1/1	0.99	0.09	40,40,40,40	0
8	GD	D	509	1/1	0.99	0.18	48,48,48,48	0
9	ZN	A	512	1/1	0.99	0.16	52,52,52,52	0
8	GD	B	509	1/1	1.00	0.17	43,43,43,43	0
7	CL	B	508	1/1	1.00	0.11	35,35,35,35	0
9	ZN	C	513	1/1	1.00	0.16	51,51,51,51	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.















**Electron density around HEM C 501:**


$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



**Electron density around HEM D 501:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





## 6.5 Other polymers [i](#)

There are no such residues in this entry.