



# wwPDB X-ray Structure Validation Summary Report ⓘ

Apr 28, 2024 – 12:21 pm BST

PDB ID : 1UWK  
Title : The High Resolution Structure of Urocanate Hydratase from *Pseudomonas putida* in complex with urocanate  
Authors : Kessler, D.; Retey, J.; Schulz, G.E.  
Deposited on : 2004-02-05  
Resolution : 1.19 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.4, CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.36.2
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.36.2


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.19 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1223 (1.22-1.18)
Clashscore	141614	1286 (1.22-1.18)
Ramachandran outliers	138981	1240 (1.22-1.18)
Sidechain outliers	138945	1239 (1.22-1.18)
RSRZ outliers	127900	1200 (1.22-1.18)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	557	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red, orange, yellow, green, grey);"></div> <div style="display: flex; justify-content: space-between; margin-top: 2px;"> <span>%</span> <span>87%</span> <span>9%</span> <span>..</span> </div> </div>
1	B	557	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red, orange, yellow, green, grey);"></div> <div style="display: flex; justify-content: space-between; margin-top: 2px;"> <span>2%</span> <span>88%</span> <span>9%</span> <span>..</span> </div> </div>

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 9437 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called UROCANATE HYDRATASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	554	Total	C	N	O	S	0	4	0
			4250	2667	757	804	22			
1	B	553	Total	C	N	O	S	0	1	0
			4239	2660	754	803	22			

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	198	SER	CYS	engineered mutation	UNP P25080
A	455	ALA	ARG	engineered mutation	UNP P25080
B	198	SER	CYS	engineered mutation	UNP P25080
B	455	ALA	ARG	engineered mutation	UNP P25080
A	164	SER	THR	conflict	UNP P25080
A	165	LEU	VAL	conflict	UNP P25080
A	167	GLY	ALA	conflict	UNP P25080
A	338	ASP	ASN	conflict	UNP P25080
B	164	SER	THR	conflict	UNP P25080
B	165	LEU	VAL	conflict	UNP P25080
B	167	GLY	ALA	conflict	UNP P25080
B	338	ASP	ASN	conflict	UNP P25080

- Molecule 2 is NICOTINAMIDE-ADENINE-DINUCLEOTIDE (three-letter code: NAD) (formula: C<sub>21</sub>H<sub>27</sub>N<sub>7</sub>O<sub>14</sub>P<sub>2</sub>).



- [illegible]

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			10	6	2	2		



*Continued from previous page...*

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	B	1	Total	C	N	O	0	0
			10	6	2	2		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	434	Total	O	0	0
			434	434		
4	B	406	Total	O	0	0
			406	406		



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	111.43Å 71.62Å 129.16Å 90.00° 98.75° 90.00°	Depositor
Resolution (Å)	50.00 – 1.19 19.57 – 1.19	Depositor EDS
% Data completeness (in resolution range)	94.7 (50.00-1.19) 98.8 (19.57-1.19)	Depositor EDS
$R_{merge}$	0.08	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.63 (at 1.19Å)	Xtriage
Refinement program	SHELXL-97	Depositor
R, $R_{free}$	0.141 , 0.179 0.135 , 0.172	Depositor DCC
$R_{free}$ test set	15983 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	8.2	Xtriage
Anisotropy	0.248	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.41 , 54.8	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.45$ , $\langle L^2 \rangle = 0.28$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.98	EDS
Total number of atoms	9437	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	12.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.24% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NAD, URO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	2.06	20/4367 (0.5%)	1.60	58/5924 (1.0%)
1	B	0.83	5/4336 (0.1%)	1.31	51/5884 (0.9%)
All	All	1.57	25/8703 (0.3%)	1.46	109/11808 (0.9%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	3

The worst 5 of 25 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	168[A]	LYS	CE-NZ	57.87	2.93	1.49
1	A	168[B]	LYS	CE-NZ	57.87	2.93	1.49
1	A	47[A]	LYS	CD-CE	54.25	2.86	1.51
1	A	47[B]	LYS	CD-CE	54.25	2.86	1.51
1	A	236	ILE	CB-CG1	30.99	2.40	1.54

The worst 5 of 109 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	47[A]	LYS	CD-CE-NZ	-32.34	37.32	111.70
1	A	47[B]	LYS	CD-CE-NZ	-32.34	37.32	111.70
1	A	168[A]	LYS	CG-CD-CE	-25.30	35.98	111.90
1	A	168[B]	LYS	CG-CD-CE	-25.30	35.98	111.90
1	A	285	ARG	CD-NE-CZ	22.20	154.67	123.60



There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	18[A]	ASN	Sidechain
1	A	18[B]	ASN	Sidechain
1	A	4	ASN	Peptide

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4250	0	4148	84	0
1	B	4239	0	4143	36	0
2	A	44	0	26	3	0
2	B	44	0	26	4	0
3	A	10	0	4	1	0
3	B	10	0	5	1	0
4	A	434	0	0	11	0
4	B	406	0	0	9	0
All	All	9437	0	8352	117	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

The worst 5 of 117 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:198[A]:SER:OG	1:A:198[A]:SER:CB	1.69	1.41
1:A:18[B]:ASN:CB	1:A:18[B]:ASN:CG	2.02	1.26
1:A:168[B]:LYS:NZ	1:A:168[B]:LYS:HE2	1.47	1.09
1:A:168[B]:LYS:NZ	1:A:168[B]:LYS:HE3	1.47	1.07
1:A:47[B]:LYS:CE	1:A:47[B]:LYS:HD2	1.52	1.03

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	556/557 (100%)	537 (97%)	18 (3%)	1 (0%)	47	19
1	B	552/557 (99%)	531 (96%)	21 (4%)	0	100	100
All	All	1108/1114 (100%)	1068 (96%)	39 (4%)	1 (0%)	51	19

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	5	ASN

### 5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	438/438 (100%)	427 (98%)	11 (2%)	47	10
1	B	435/438 (99%)	427 (98%)	8 (2%)	59	21
All	All	873/876 (100%)	854 (98%)	19 (2%)	52	14

5 of 19 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	282	GLU
1	B	402	SER
1	B	532	ARG
1	B	297	LYS
1	A	316	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 10 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	268	HIS
1	B	391	HIS
1	B	500	GLN
1	A	391	HIS
1	A	490	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	NAD	B	1558	-	42,48,48	1.86	6 (14%)	50,73,73	2.56	16 (32%)
2	NAD	A	1558	-	42,48,48	1.63	4 (9%)	50,73,73	2.08	12 (24%)
3	URO	A	1559	-	8,10,10	2.62	3 (37%)	7,12,12	5.60	4 (57%)
3	URO	B	1559	-	8,10,10	2.45	3 (37%)	7,12,12	3.58	2 (28%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral

centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAD	B	1558	-	-	5/26/62/62	0/5/5/5
2	NAD	A	1558	-	-	4/26/62/62	0/5/5/5
3	URO	A	1559	-	-	1/3/5/5	0/1/1/1
3	URO	B	1559	-	-	3/3/5/5	0/1/1/1

The worst 5 of 16 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	1558	NAD	C3N-C7N	-7.19	1.39	1.50
2	A	1558	NAD	C3N-C7N	-7.02	1.40	1.50
2	B	1558	NAD	O4D-C1D	6.42	1.50	1.41
3	A	1559	URO	CAJ-CAD	-4.62	1.35	1.47
3	B	1559	URO	CAJ-CAD	-4.13	1.36	1.47

The worst 5 of 34 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	1559	URO	CAJ-CAD-CAC	12.65	156.46	126.14
3	B	1559	URO	CAJ-CAD-CAC	8.31	146.05	126.14
2	A	1558	NAD	O7N-C7N-N7N	-7.57	111.83	122.58
2	B	1558	NAD	O7N-C7N-C3N	7.14	128.17	119.63
2	A	1558	NAD	C3N-C7N-N7N	6.60	125.67	117.75

There are no chirality outliers.

5 of 13 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	1558	NAD	O4D-C1D-N1N-C2N
2	A	1558	NAD	O4D-C1D-N1N-C6N
2	A	1558	NAD	C2D-C1D-N1N-C2N
2	B	1558	NAD	O4D-C1D-N1N-C2N
2	B	1558	NAD	O4D-C1D-N1N-C6N

There are no ring outliers.

4 monomers are involved in 9 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	1558	NAD	4	0


*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1558	NAD	3	0
3	A	1559	URO	1	0
3	B	1559	URO	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	554/557 (99%)	-0.30	6 (1%) 80 80	5, 9, 21, 34	0
1	B	553/557 (99%)	-0.25	9 (1%) 72 72	6, 10, 23, 44	0
All	All	1107/1114 (99%)	-0.28	15 (1%) 75 75	5, 10, 22, 44	0

The worst 5 of 15 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	4	ASN	4.3
1	B	164	SER	3.6
1	B	290	THR	3.2
1	A	557	GLY	3.0
1	B	282	GLU	3.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)


There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)


In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	URO	A	1559	10/10	0.85	0.26	16,42,48,52	0
3	URO	B	1559	10/10	0.90	0.28	12,51,61,61	0
2	NAD	A	1558	44/44	0.99	0.06	5,6,47,73	0
2	NAD	B	1558	44/44	0.99	0.06	7,8,39,74	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.







## 6.5 Other polymers [i](#)

There are no such residues in this entry.