



# wwPDB X-ray Structure Validation Summary Report ⓘ

May 22, 2020 – 12:26 am BST

PDB ID : 3V89  
Title : The crystal structure of transferrin binding protein A (TbpA) from *Neisseria meningitidis* serogroup B in complex with the C-lobe of human transferrin  
Authors : Noinaj, N.; Oke, M.; Easley, N.; Zak, O.; Aisen, P.; Buchanan, S.K.  
Deposited on : 2011-12-22  
Resolution : 3.10 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 3.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1094 (3.10-3.10)
Clashscore	141614	1184 (3.10-3.10)
Ramachandran outliers	138981	1141 (3.10-3.10)
Sidechain outliers	138945	1141 (3.10-3.10)
RSRZ outliers	127900	1067 (3.10-3.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	904	
2	B	343	

## 2 Entry composition [i](#)

There are 2 unique types of molecules in this entry. The entry contains 9110 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Transferrin-binding protein A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	853	6558	4091	1182	1274	11	0	0	0

There are 13 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	12	MET	-	EXPRESSION TAG	UNP Q9JP J0
A	13	ASP	-	EXPRESSION TAG	UNP Q9JP J0
A	14	ILE	-	EXPRESSION TAG	UNP Q9JP J0
A	15	HIS	-	EXPRESSION TAG	UNP Q9JP J0
A	16	HIS	-	EXPRESSION TAG	UNP Q9JP J0
A	17	HIS	-	EXPRESSION TAG	UNP Q9JP J0
A	18	HIS	-	EXPRESSION TAG	UNP Q9JP J0
A	19	HIS	-	EXPRESSION TAG	UNP Q9JP J0
A	20	HIS	-	EXPRESSION TAG	UNP Q9JP J0
A	21	HIS	-	EXPRESSION TAG	UNP Q9JP J0
A	22	HIS	-	EXPRESSION TAG	UNP Q9JP J0
A	23	HIS	-	EXPRESSION TAG	UNP Q9JP J0
A	24	HIS	-	EXPRESSION TAG	UNP Q9JP J0

- Molecule 2 is a protein called Serotransferrin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	339	2552	1587	436	503	26	0	0	0

There is a discrepancy between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
B	429	VAL	ILE	VARIANT	UNP P02787





• Molecule 2: Serotransferrin

Chain B: 47% 45% 6%



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	58.05Å 107.59Å 130.72Å 90.00° 94.48° 90.00°	Depositor
Resolution (Å)	29.96 – 3.10 41.69 – 3.09	Depositor EDS
% Data completeness (in resolution range)	96.8 (29.96-3.10) 91.6 (41.69-3.09)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.06	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.77 (at 3.06Å)	Xtrriage
Refinement program	PHENIX (phenix.refine: 1.7.2_865)	Depositor
R, $R_{free}$	0.224 , 0.287 0.225 , 0.283	Depositor DCC
$R_{free}$ test set	1407 reflections (4.95%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	57.6	Xtrriage
Anisotropy	0.388	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.29 , 52.8	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.90	EDS
Total number of atoms	9110	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	73.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.06% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.31	0/6701	0.57	0/9081
2	B	0.28	0/2605	0.50	0/3535
All	All	0.30	0/9306	0.55	0/12616

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	289	VAL	Peptide
1	A	743	ASP	Peptide

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	6558	0	6177	605	1
2	B	2552	0	2359	148	1

*Continued on next page...*

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	9110	0	8536	740	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 42.

The worst 5 of 740 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:854:TYR:CE1	1:A:856:VAL:HG12	1.47	1.46
1:A:855:THR:CG2	1:A:861:THR:HB	1.43	1.45
1:A:52:ASP:HB3	1:A:53:ASN:CB	1.51	1.40
1:A:854:TYR:CD1	1:A:856:VAL:HG12	1.60	1.37
1:A:75:ASN:ND2	1:A:77:ARG:H	1.19	1.34

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:290:SER:OG	2:B:438:ASP:O[2_454]	2.02	0.18

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	847/904 (94%)	757 (89%)	72 (8%)	18 (2%)	7	30
2	B	337/343 (98%)	311 (92%)	26 (8%)	0	100	100
All	All	1184/1247 (95%)	1068 (90%)	98 (8%)	18 (2%)	10	39

5 of 18 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	52	ASP
1	A	55	VAL
1	A	56	THR
1	A	58	LEU
1	A	800	PRO

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	662/741 (89%)	530 (80%)	132 (20%)	<b>1</b> <b>5</b>
2	B	271/292 (93%)	233 (86%)	38 (14%)	<b>3</b> <b>15</b>
All	All	933/1033 (90%)	763 (82%)	170 (18%)	<b>1</b> <b>7</b>

5 of 170 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	508	SER
1	A	625	SER
2	B	611	ASN
1	A	524	TYR
1	A	566	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 8 such sidechains are listed below:

Mol	Chain	Res	Type
1	A	439	ASN
2	B	535	HIS
1	A	829	ASN
1	A	233	HIS
1	A	698	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

#### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

#### 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

#### 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

#### 5.7 Other polymers [i](#)

There are no such residues in this entry.

#### 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	853/904 (94%)	-0.21	8 (0%) 84 69	24, 73, 122, 160	0
2	B	339/343 (98%)	-0.34	0 100 100	31, 69, 112, 126	0
All	All	1192/1247 (95%)	-0.25	8 (0%) 87 75	24, 72, 118, 160	0

The worst 5 of 8 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	802	GLY	3.2
1	A	621	HIS	2.7
1	A	320	PHE	2.5
1	A	414	TYR	2.4
1	A	408	TYR	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

There are no ligands in this entry.

### 6.5 Other polymers [i](#)

There are no such residues in this entry.