



Full wwPDB X-ray Structure Validation Report ⓘ

Jun 17, 2024 – 04:28 AM EDT

PDB ID : 2WSH
Title : Structure of bacteriophage T4 EndoII E118A mutant
Authors : Andersson, C.E.; Lagerback, P.; Carlson, K.
Deposited on : 2009-09-07
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 1.20.1
EDS : 2.37.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.37.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	6207 (1.90-1.90)
Clashscore	141614	6847 (1.90-1.90)
Ramachandran outliers	138981	6760 (1.90-1.90)
Sidechain outliers	138945	6760 (1.90-1.90)
RSRZ outliers	127900	6082 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	143	<div> <div>%</div> <div> <div></div> <div>76%</div> <div>16%</div> <div>• 6%</div> </div> </div>
1	B	143	<div> <div>%</div> <div> <div></div> <div>73%</div> <div>18%</div> <div>• 8%</div> </div> </div>
1	C	143	<div> <div>%</div> <div> <div></div> <div>73%</div> <div>16%</div> <div>• 10%</div> </div> </div>
1	D	143	<div> <div></div> <div> <div></div> <div>74%</div> <div>19%</div> <div>7%</div> </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	PO4	A	1133	-	-	X	-

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 4775 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ENDONUCLEASE II.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	134	Total	C	N	O	S	Se	0	6	0
			1138	724	193	217	1	3			
1	B	131	Total	C	N	O	S	Se	0	1	0
			1080	693	183	200	1	3			
1	C	129	Total	C	N	O	S	Se	0	1	0
			1064	683	181	196	1	3			
1	D	133	Total	C	N	O	S	Se	0	2	0
			1101	704	187	206	1	3			

There are 4 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	118	ALA	GLU	engineered mutation	UNP P07059
B	118	ALA	GLU	engineered mutation	UNP P07059
C	118	ALA	GLU	engineered mutation	UNP P07059
D	118	ALA	GLU	engineered mutation	UNP P07059

- Molecule 2 is PHOSPHATE ION (three-letter code: PO4) (formula: O₄P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	P	0	0
			5	4	1		

- Molecule 3 is DI(HYDROXYETHYL)ETHER (three-letter code: PEG) (formula: $C_4H_{10}O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	B	1	Total	C	O	0	0
			7	4	3		
3	C	1	Total	C	O	0	0
			7	4	3		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	70	Total 70	O 70	0	0
4	B	104	Total 104	O 104	0	0
4	C	98	Total 98	O 98	0	0
4	D	101	Total 101	O 101	0	0

- Molecule 1: ENDONUCLEASE II



LYS	LYS	LYS	LYS	LYS	H-2	A5	T6	E7	F10	I11	M21	M42	D43	V46	V47	I48	M59	R62	T63	A64	R67	A68	D69	K70	THR	SER	D73	S77	L100	S101	M102	M109	T110	I111	L116	E117	A118	P119	M126	P127	P128	M129	M130	I131	GLN	LYS
-----	-----	-----	-----	-----	-----	----	----	----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	-----	-----

MSE	HIS	HIS	HIS	HIS	H-3	H-3	H-2	H-1	M1	K2	E3	I4			I11	K12	Y13	L27	S28	L29		V46	A64	I65	N66	R67	LYS	ASP	LYS	THR	SER	D73		Y93		C97		T113	A118	P119		M126	P127	P128		I131	G1N	HIS	LYS	LYS	LYS	LVS
-----	-----	-----	-----	-----	-----	-----	-----	-----	----	----	----	----	--	--	-----	-----	-----	-----	-----	-----	--	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	--	-----	--	-----	--	------	------	------	--	------	------	------	--	------	-----	-----	-----	-----	-----	-----

MSE	HIS	A5	F10	E17	K25	G26	L27	N34	I41	I48	R57	Y60	S77	S82	A83	E86	F98	N99	M109	A112	P119	F125	M126	P127	P128	M129	M130	I131	GLN	HIS	LVS	LVS	TYS
-----	-----	----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	------	------	------	------	------	------	------	------	------	------	-----	-----	-----	-----	-----

4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	57.42Å 104.97Å 57.87Å 90.00° 113.56° 90.00°	Depositor
Resolution (Å)	53.04 – 1.90 53.04 – 1.90	Depositor EDS
% Data completeness (in resolution range)	92.5 (53.04-1.90) 98.5 (53.04-1.90)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.76 (at 1.90Å)	Xtriage
Refinement program	PHENIX (PHENIX.REFINE)	Depositor
R, R_{free}	0.200 , 0.247 0.193 , 0.238	Depositor DCC
R_{free} test set	2466 reflections (5.06%)	wwPDB-VP
Wilson B-factor (Å ²)	25.9	Xtriage
Anisotropy	0.131	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 50.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.035 for l,-k,h	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	4775	wwPDB-VP
Average B, all atoms (Å ²)	32.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.77% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PEG, PO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.24	0/1156	0.46	0/1556
1	B	0.24	0/1097	0.42	0/1473
1	C	0.24	0/1082	0.42	0/1455
1	D	0.25	0/1119	0.43	0/1505
All	All	0.24	0/4454	0.44	0/5989

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1138	0	1144	21	0
1	B	1080	0	1099	29	0
1	C	1064	0	1076	22	0
1	D	1101	0	1117	15	0
2	A	5	0	0	2	0
3	B	7	0	10	2	0
3	C	7	0	10	2	0
4	A	70	0	0	0	0
4	B	104	0	0	4	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	C	98	0	0	1	0
4	D	101	0	0	0	0
All	All	4775	0	4456	83	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (83) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:48:ILE:H	1:B:130:ASN:HD21	1.10	0.97
1:D:48:ILE:H	1:D:130:ASN:HD21	1.22	0.87
1:C:-2:HIS:HB3	1:C:4:ILE:HD11	1.58	0.82
1:D:127:PRO:HB2	1:D:130:ASN:HD22	1.48	0.79
1:A:98:PHE:HZ	1:B:102:MSE:HE1	1.50	0.76
1:B:127:PRO:HB2	1:B:130:ASN:HD22	1.51	0.73
1:D:34[A]:ASN:HD21	1:D:112:ALA:HB1	1.53	0.73
1:A:31[A]:ASN:HD22	1:A:32:LYS:N	1.86	0.72
1:C:131:ILE:HD11	3:C:1132:PEG:H11	1.71	0.72
1:D:-2:HIS:HE1	1:D:125:PHE:O	1.76	0.68
1:A:46:VAL:HA	1:A:128:PRO:HD2	1.78	0.66
1:B:118:ALA:HB3	1:B:119:PRO:HD3	1.77	0.65
1:B:111:ILE:HD11	1:B:116:LEU:HD21	1.78	0.64
1:C:-1:HIS:CE1	1:C:2:LYS:HG2	2.34	0.62
1:A:98:PHE:CZ	1:B:102:MSE:HE1	2.32	0.61
1:C:64:ALA:O	1:C:67:ARG:HG2	2.00	0.61
1:D:5:ALA:HA	1:D:10:PHE:HB2	1.82	0.61
1:B:111:ILE:HG13	4:B:2093:HOH:O	2.01	0.61
1:D:83:ALA:O	1:D:86:GLU:HG2	2.00	0.60
1:B:62:ARG:HD3	4:B:2060:HOH:O	2.01	0.60
1:A:5:ALA:HA	1:A:10:PHE:HB2	1.83	0.60
1:C:-1:HIS:HE1	1:C:2:LYS:HG2	1.66	0.59
1:A:72:SER:O	1:A:73:ASP:HB3	2.01	0.59
1:D:99:ASN:HD22	1:D:112:ALA:HA	1.68	0.58
1:B:67:ARG:HD2	1:B:69:ASP:OD2	2.03	0.58
1:A:57[B]:ARG:NH1	1:A:60:TYR:CD2	2.73	0.57
1:A:57[A]:ARG:HD2	2:A:1133:PO4:O4	2.05	0.56
1:A:118:ALA:HB3	1:A:119:PRO:HD3	1.88	0.56
1:B:46:VAL:HA	1:B:128:PRO:HD2	1.89	0.54
1:D:118:ALA:HB3	1:D:119:PRO:HD3	1.88	0.54
1:B:102:MSE:HE3	1:D:98:PHE:CE1	2.44	0.53

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:111:ILE:HD11	1:B:116:LEU:HD11	1.92	0.52
1:A:109:MSE:N	1:A:109:MSE:HE3	2.25	0.52
1:C:126:ASN:N	1:C:127:PRO:HD3	2.25	0.52
1:B:129:TRP:HA	3:B:1132:PEG:H31	1.91	0.52
1:C:13:TYR:CZ	1:C:29:ILE:HD13	2.46	0.51
1:B:100:LEU:HB2	1:B:111:ILE:CG2	2.41	0.51
1:D:34[A]:ASN:ND2	1:D:112:ALA:HB1	2.25	0.50
1:D:77:SER:O	1:D:81:HIS:HD2	1.95	0.50
1:A:57[B]:ARG:O	1:A:60:TYR:HB3	2.12	0.49
1:B:7:GLU:HG2	4:B:2005:HOH:O	2.13	0.49
1:C:-1:HIS:HB3	1:C:3:GLU:CG	2.43	0.49
1:C:-1:HIS:HB3	1:C:3:GLU:HG2	1.93	0.49
1:C:46:VAL:HA	1:C:128:PRO:HD2	1.95	0.48
1:C:-2:HIS:HB3	1:C:4:ILE:CD1	2.38	0.48
1:B:109:MSE:N	1:B:109:MSE:HE3	2.28	0.48
1:C:97:CYS:HB3	1:C:113:THR:OG1	2.14	0.47
1:A:35:VAL:HG12	1:A:96:GLN:HA	1.95	0.47
1:C:131:ILE:CD1	3:C:1132:PEG:H11	2.43	0.47
1:B:131:ILE:HD12	3:B:1132:PEG:H42	1.97	0.47
1:A:57[B]:ARG:NE	2:A:1133:PO4:O4	2.48	0.47
1:B:48:ILE:H	1:B:130:ASN:ND2	1.94	0.46
1:B:126:ASN:N	1:B:127:PRO:HD3	2.30	0.46
1:C:65:ILE:HD12	1:C:66:ASN:N	2.31	0.46
1:C:13:TYR:OH	1:C:29:ILE:HD13	2.15	0.46
1:B:5:ALA:HA	1:B:10:PHE:HB2	1.98	0.46
1:B:59:ASN:ND2	4:B:2060:HOH:O	2.48	0.45
1:C:118:ALA:HB3	1:C:119:PRO:HD3	1.98	0.45
1:C:1:MSE:HG2	1:C:93:TYR:CE2	2.51	0.45
1:A:1:MSE:HB3	1:A:43[B]:ASP:OD1	2.16	0.45
1:C:127:PRO:HA	1:C:128:PRO:HD3	1.88	0.44
1:B:64:ALA:HB3	1:B:77:SER:CB	2.47	0.44
1:D:126:ASN:N	1:D:127:PRO:HD3	2.33	0.44
1:A:-2:HIS:HB3	1:A:4:ILE:HD11	2.00	0.43
1:D:17:GLU:HG3	1:D:25:LYS:HD2	2.01	0.43
1:C:4:ILE:HD12	1:C:4:ILE:H	1.83	0.43
1:A:-1:HIS:CE1	1:A:43[B]:ASP:OD2	2.71	0.43
1:B:109:MSE:HE3	1:B:109:MSE:H	1.82	0.43
1:B:111:ILE:HG23	1:B:111:ILE:O	2.19	0.43
1:D:41:ILE:HD11	1:D:129:TRP:CH2	2.53	0.43
1:C:-3:HIS:HA	4:C:2038:HOH:O	2.19	0.42
1:D:57[A]:ARG:O	1:D:60:TYR:HB3	2.19	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:58:ILE:HD12	1:A:92:PHE:CE1	2.54	0.42
1:B:100:LEU:HB2	1:B:111:ILE:HG22	2.01	0.42
1:B:42:ASN:O	1:B:43:ASP:HB2	2.19	0.41
1:B:111:ILE:CD1	1:B:116:LEU:HD11	2.51	0.41
1:B:59:ASN:ND2	1:B:62:ARG:HH11	2.18	0.41
1:C:118:ALA:N	1:C:119:PRO:CD	2.83	0.41
1:B:11:ILE:HG23	1:C:11:ILE:HD11	2.03	0.41
1:A:71[A]:THR:O	1:A:71[A]:THR:HG22	2.20	0.41
1:A:54:LEU:O	1:A:58:ILE:HG12	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	138/143 (96%)	132 (96%)	6 (4%)	0	100	100
1	B	128/143 (90%)	127 (99%)	1 (1%)	0	100	100
1	C	126/143 (88%)	123 (98%)	3 (2%)	0	100	100
1	D	133/143 (93%)	131 (98%)	2 (2%)	0	100	100
All	All	525/572 (92%)	513 (98%)	12 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	127/126 (101%)	122 (96%)	5 (4%)	32	23
1	B	119/126 (94%)	118 (99%)	1 (1%)	81	82
1	C	117/126 (93%)	114 (97%)	3 (3%)	46	39
1	D	122/126 (97%)	119 (98%)	3 (2%)	47	41
All	All	485/504 (96%)	473 (98%)	12 (2%)	50	41

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	31[A]	ASN
1	A	31[B]	ASN
1	A	34	ASN
1	A	72	SER
1	A	109	MSE
1	B	109	MSE
1	C	-3	HIS
1	C	3	GLU
1	C	27	LEU
1	D	27	LEU
1	D	109	MSE
1	D	131	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	34	ASN
1	A	59	ASN
1	B	59	ASN
1	B	130	ASN
1	D	-2	HIS
1	D	66	ASN
1	D	81	HIS
1	D	99	ASN
1	D	130	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no monosaccharides in this entry.

5.6 Ligand geometry ⓘ

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	PO4	A	1133	-	4,4,4	0.95	0	6,6,6	0.60	0
3	PEG	C	1132	-	6,6,6	0.49	0	5,5,5	1.46	0
3	PEG	B	1132	-	6,6,6	0.50	0	5,5,5	1.40	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	PEG	C	1132	-	-	2/4/4/4	-
3	PEG	B	1132	-	-	2/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	B	1132	PEG	O2-C3-C4-O4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
3	B	1132	PEG	C1-C2-O2-C3
3	C	1132	PEG	O2-C3-C4-O4
3	C	1132	PEG	C4-C3-O2-C2

There are no ring outliers.

3 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1133	PO4	2	0
3	C	1132	PEG	2	0
3	B	1132	PEG	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	131/143 (91%)	-0.33	1 (0%) 86 87	20, 32, 54, 69	0
1	B	128/143 (89%)	-0.36	1 (0%) 86 87	16, 29, 52, 78	0
1	C	126/143 (88%)	-0.36	2 (1%) 72 74	18, 29, 51, 75	0
1	D	130/143 (90%)	-0.39	0 100 100	17, 27, 49, 64	0
All	All	515/572 (90%)	-0.36	4 (0%) 86 87	16, 29, 53, 78	0

All (4) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	67	ARG	3.1
1	A	-1	HIS	2.3
1	B	21	ASN	2.1
1	C	66	ASN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	PEG	C	1132	7/7	0.89	0.13	36,38,54,57	0
2	PO4	A	1133	5/5	0.93	0.18	39,44,62,68	0
3	PEG	B	1132	7/7	0.95	0.09	30,36,44,49	0

6.5 Other polymers [i](#)

There are no such residues in this entry.