



# Full wwPDB X-ray Structure Validation Report ⓘ

Jun 16, 2024 – 11:34 AM EDT

PDB ID : 4WSH  
Title : Crystal structure of Probable Uroporphyrinogen decarboxylase (UPD) (URO-D) from *Pseudomonas aeruginosa*  
Authors : Seattle Structural Genomics Center for Infectious Disease (SSGCID)  
Deposited on : 2014-10-27  
Resolution : 1.95 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.37.1
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.37.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	2580 (1.96-1.96)
Clashscore	141614	2705 (1.96-1.96)
Ramachandran outliers	138981	2678 (1.96-1.96)
Sidechain outliers	138945	2678 (1.96-1.96)
RSRZ outliers	127900	2539 (1.96-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	363	<div> <div>3%</div> <div> <div></div> <div>94%</div> <div></div> </div> <div></div> </div>
1	B	363	<div> <div>6%</div> <div> <div></div> <div>91%</div> <div></div> </div> <div></div> </div>

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 5849 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Uroporphyrinogen decarboxylase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	355	Total	C	N	O	S	0	13	0
			2771	1761	491	502	17			
1	B	347	Total	C	N	O	S	0	8	0
			2655	1693	465	480	17			

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-7	MET	-	initiating methionine	UNP P95458
A	-6	ALA	-	expression tag	UNP P95458
A	-5	HIS	-	expression tag	UNP P95458
A	-4	HIS	-	expression tag	UNP P95458
A	-3	HIS	-	expression tag	UNP P95458
A	-2	HIS	-	expression tag	UNP P95458
A	-1	HIS	-	expression tag	UNP P95458
A	0	HIS	-	expression tag	UNP P95458
B	-7	MET	-	initiating methionine	UNP P95458
B	-6	ALA	-	expression tag	UNP P95458
B	-5	HIS	-	expression tag	UNP P95458
B	-4	HIS	-	expression tag	UNP P95458
B	-3	HIS	-	expression tag	UNP P95458
B	-2	HIS	-	expression tag	UNP P95458
B	-1	HIS	-	expression tag	UNP P95458
B	0	HIS	-	expression tag	UNP P95458

- Molecule 2 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Cl	0	0
			1	1		
2	B	1	Total	Cl	0	0
			1	1		

- Molecule 3 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	S	0	0
			5	4	1		
3	A	1	Total	O	S	0	0
			5	4	1		
3	A	1	Total	O	S	0	0
			5	4	1		
3	A	1	Total	O	S	0	0
			5	4	1		
3	A	1	Total	O	S	0	0
			5	4	1		
3	A	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	211	Total	O	0	0
			211	211		
4	B	165	Total	O	0	0
			165	165		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Uroporphyrinogen decarboxylase



- Molecule 1: Uroporphyrinogen decarboxylase



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	71.76Å 73.88Å 141.88Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	39.49 – 1.95 39.49 – 1.95	Depositor EDS
% Data completeness (in resolution range)	99.6 (39.49-1.95) 99.7 (39.49-1.95)	Depositor EDS
$R_{merge}$	0.04	Depositor
$R_{sym}$	0.04	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.46 (at 1.95Å)	Xtriage
Refinement program	PHENIX	Depositor
R, $R_{free}$	0.158 , 0.201 0.158 , 0.201	Depositor DCC
$R_{free}$ test set	2854 reflections (5.14%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	32.2	Xtriage
Anisotropy	0.338	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 51.3	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.017 for k,h,-l	Xtriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	5849	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	36.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.92% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.36	0/2864	0.51	0/3885
1	B	0.33	0/2740	0.47	0/3721
All	All	0.35	0/5604	0.49	0/7606

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2771	0	2740	10	0
1	B	2655	0	2611	11	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	35	0	0	1	0
3	B	10	0	0	0	0
4	A	211	0	0	2	0
4	B	165	0	0	0	0
All	All	5849	0	5351	20	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (20) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:79:ILE:O	4:A:685:HOH:O	2.17	0.59
1:A:85:ALA:HB2	1:A:122:LEU:HD21	1.85	0.59
1:A:19:ASP:OD1	4:A:681:HOH:O	2.17	0.58
1:B:64:LEU:HD11	1:B:73:ALA:HB2	1.87	0.57
1:A:150:TRP:O	1:A:154:THR:HG23	2.12	0.50
1:B:150:TRP:O	1:B:154:THR:HG23	2.12	0.50
1:B:79:ILE:HD12	1:B:145:PHE:HE2	1.78	0.48
1:A:27[B]:ARG:HD3	3:A:402:SO4:O4	2.16	0.46
1:B:251:LYS:HE3	1:B:328:HIS:ND1	2.31	0.46
1:B:150:TRP:CZ2	1:B:215:LEU:HD21	2.51	0.45
1:B:27:ARG:HD2	1:B:329:GLY:HA3	1.98	0.45
1:B:251:LYS:HE3	1:B:328:HIS:CG	2.53	0.44
1:A:47:MET:SD	1:A:92:PHE:HE2	2.41	0.44
1:A:51[B]:MET:HG2	1:A:124:TYR:CE2	2.54	0.43
1:A:295:PRO:HG2	1:A:328:HIS:HB3	2.01	0.43
1:B:79:ILE:HD12	1:B:145:PHE:CE2	2.54	0.43
1:A:174:ASP:OD1	1:B:301:ASN:ND2	2.53	0.41
1:A:79:ILE:HD12	1:A:79:ILE:HA	1.89	0.40
1:B:229:ARG:HA	1:B:263:VAL:HG13	2.03	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	366/363 (101%)	358 (98%)	8 (2%)	0	100	100
1	B	351/363 (97%)	343 (98%)	8 (2%)	0	100	100
All	All	717/726 (99%)	701 (98%)	16 (2%)	0	100	100

There are no Ramachandran outliers to report.



### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	280/286 (98%)	279 (100%)	1 (0%)	91	90
1	B	266/286 (93%)	265 (100%)	1 (0%)	91	90
All	All	546/572 (96%)	544 (100%)	2 (0%)	91	90

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	237	ARG
1	B	134	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 11 ligands modelled in this entry, 2 are monoatomic - leaving 9 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The

Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	SO4	B	402	-	4,4,4	0.13	0	6,6,6	0.12	0
3	SO4	A	401	-	4,4,4	0.14	0	6,6,6	0.23	0
3	SO4	A	403	-	4,4,4	0.13	0	6,6,6	0.14	0
3	SO4	A	405	-	4,4,4	0.14	0	6,6,6	0.05	0
3	SO4	A	404	-	4,4,4	0.15	0	6,6,6	0.15	0
3	SO4	A	402	-	4,4,4	0.15	0	6,6,6	0.13	0
3	SO4	B	401	-	4,4,4	0.23	0	6,6,6	0.22	0
3	SO4	A	407	-	4,4,4	0.13	0	6,6,6	0.09	0
3	SO4	A	406	-	4,4,4	0.14	0	6,6,6	0.05	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	402	SO4	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	355/363 (97%)	0.05	12 (3%) 45 55	20, 29, 54, 86	0
1	B	347/363 (95%)	0.11	21 (6%) 21 29	24, 34, 71, 99	0
All	All	702/726 (96%)	0.08	33 (4%) 31 41	20, 31, 66, 99	0

All (33) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	91	TYR	7.2
1	A	92	PHE	6.1
1	A	94	THR	6.0
1	B	98	PRO	4.8
1	A	355	GLY	4.8
1	B	104	VAL	4.7
1	A	1	MET	4.6
1	A	163[A]	LYS	4.5
1	B	108	ALA	4.5
1	B	3	ALA	4.4
1	A	95	GLY	4.3
1	B	100	PHE	4.2
1	B	110	ILE	4.1
1	B	90	LEU	3.7
1	B	115	VAL	3.6
1	B	112	ALA	3.6
1	A	2	THR	3.3
1	A	96	GLU	3.3
1	B	103	VAL	3.3
1	B	106	SER	3.0
1	B	107	LEU	3.0
1	B	99	ARG	3.0
1	B	162	SER	2.7
1	B	102	LYS	2.6

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	97	GLY	2.6
1	B	89	GLY	2.4
1	B	118	PRO	2.4
1	A	45	ASP	2.4
1	A	207	ILE	2.3
1	B	113	LEU	2.3
1	B	87	GLY	2.1
1	A	43	ALA	2.0
1	B	105	SER	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	SO4	A	407	5/5	0.80	0.40	127,129,131,132	0
3	SO4	A	403	5/5	0.84	0.17	40,51,62,68	5
3	SO4	A	406	5/5	0.89	0.16	120,120,123,124	0
3	SO4	A	405	5/5	0.91	0.21	98,102,103,108	0
3	SO4	B	402	5/5	0.94	0.14	87,92,96,97	0
3	SO4	A	404	5/5	0.97	0.11	70,71,72,74	0
2	CL	B	400	1/1	0.98	0.05	37,37,37,37	0
3	SO4	A	401	5/5	0.98	0.14	36,37,48,51	0
3	SO4	A	402	5/5	0.98	0.13	48,51,56,58	0
3	SO4	B	401	5/5	0.99	0.08	31,34,39,39	0
2	CL	A	400	1/1	0.99	0.07	42,42,42,42	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.