



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 27, 2024 – 03:03 pm BST

PDB ID : 2WYH
Title : Structure of the Streptococcus pyogenes family GH38 alpha-mannosidase
Authors : Suits, M.D.L.; Zhu, Y.; Taylor, E.J.; Zechel, D.L.; Gilbert, H.J.; Davies, G.J.
Deposited on : 2009-11-16
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.4, CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36.2
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	6207 (1.90-1.90)
Clashscore	141614	6847 (1.90-1.90)
Ramachandran outliers	138981	6760 (1.90-1.90)
Sidechain outliers	138945	6760 (1.90-1.90)
RSRZ outliers	127900	6082 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	923	<div> <div>%</div> <div> <div></div> <div>92%</div> <div>6%</div> <div>.</div> </div> </div>
1	B	923	<div> <div>3%</div> <div> <div></div> <div>91%</div> <div>5%</div> <div>.</div> </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	GOL	A	903	-	-	X	-
3	GOL	A	906	-	-	X	-
4	TRS	B	903	-	-	X	-

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 15430 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called ALPHA-MANNOSIDASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	905	Total	C	N	O	S	0	9	0
			7272	4600	1247	1395	30			
1	B	891	Total	C	N	O	S	0	6	0
			7161	4525	1231	1376	29			

- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Zn	0	0
			1	1		
2	B	1	Total	Zn	0	0
			1	1		

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O 6 3 3	0	0
3	A	1	Total C O 6 3 3	0	0
3	A	1	Total C O 6 3 3	0	0
3	A	1	Total C O 6 3 3	0	0
3	A	1	Total C O 6 3 3	0	0
3	A	1	Total C O 6 3 3	0	0
3	A	1	Total C O 6 3 3	0	0
3	B	1	Total C O 6 3 3	0	0
3	B	1	Total C O 6 3 3	0	0
3	B	1	Total C O 6 3 3	0	0
3	B	1	Total C O 6 3 3	0	0
3	B	1	Total C O 6 3 3	0	0

- Molecule 4 is 2-AMINO-2-HYDROXYMETHYL-PROPANE-1,3-DIOL (three-letter code: TRS) (formula: $C_4H_{12}NO_3$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	B	1	Total	C	N	O	0	0
			8	4	1	3		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	470	Total	O	0	0
			470	470		
5	B	445	Total	O	0	0
			445	445		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: ALPHA-MANNOSIDASE



• Molecule 1: ALPHA-MANNOSIDASE



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	92.57Å 88.50Å 134.69Å 90.00° 108.97° 90.00°	Depositor
Resolution (Å)	127.00 – 1.90 46.28 – 1.90	Depositor EDS
% Data completeness (in resolution range)	99.1 (127.00-1.90) 99.1 (46.28-1.90)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.66 (at 1.90Å)	Xtriage
Refinement program	REFMAC 5.5.0109	Depositor
R, R_{free}	0.177 , 0.205 0.179 , 0.205	Depositor DCC
R_{free} test set	8041 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å ²)	18.9	Xtriage
Anisotropy	0.118	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 47.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.014 for h,-k,-h-l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	15430	wwPDB-VP
Average B, all atoms (Å ²)	16.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.61% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: TRS, ZN, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.46	0/7469	0.58	1/10140 (0.0%)
1	B	0.47	0/7342	0.57	0/9965
All	All	0.46	0/14811	0.57	1/20105 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	3
1	B	0	2
All	All	0	5

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	-5	LEU	CA-CB-CG	5.51	127.98	115.30

There are no chirality outliers.

All (5) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	84	GLY	Mainchain,Peptide
1	A	90	GLN	Peptide
1	B	84	GLY	Mainchain,Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	7272	0	7044	40	0
1	B	7161	0	6918	35	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	42	0	56	20	0
3	B	30	0	40	2	0
4	B	8	0	12	12	0
5	A	470	0	0	3	0
5	B	445	0	0	1	0
All	All	15430	0	14070	76	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (76) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:479:PRO:HB3	3:A:906:GOL:H31	1.52	0.88
1:A:202:GLU:H	3:A:905:GOL:H31	1.36	0.86
3:A:904:GOL:H31	5:A:2208:HOH:O	1.75	0.86
1:B:156:PHE:HA	1:B:169:GLN:HG3	1.55	0.85
1:B:480:LYS:H	3:B:906:GOL:H12	1.40	0.85
1:A:24:GLN:HG2	3:A:903:GOL:O3	1.78	0.84
1:B:28:ARG:HH22	4:B:903:TRS:H21	1.42	0.82
1:B:28:ARG:HH12	4:B:903:TRS:H32	1.42	0.82
1:A:480:LYS:H	3:A:906:GOL:H12	1.41	0.82
1:A:202:GLU:H	3:A:905:GOL:C3	1.98	0.76
1:A:379:TYR:HA	4:B:903:TRS:H12	1.74	0.68
1:A:687:ASP:H	3:A:908:GOL:C1	2.08	0.67
1:A:-9:GLY:O	1:A:-6:VAL:HG22	1.96	0.66
1:B:28:ARG:HH22	4:B:903:TRS:C2	2.12	0.63
3:A:903:GOL:H11	1:B:382:LYS:NZ	2.13	0.62
1:A:379:TYR:HA	4:B:903:TRS:C1	2.30	0.62
1:A:521:THR:HG23	1:A:522:VAL:H	1.65	0.61
1:B:299:SER:OG	1:B:756[A]:ARG:HD3	2.00	0.61

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:156:PHE:HA	1:B:169:GLN:CG	2.30	0.61
1:A:688[B]:HIS:ND1	3:A:908:GOL:H32	2.16	0.60
1:B:155:GLY:C	1:B:157:ASN:H	2.04	0.60
1:B:28:ARG:HH12	4:B:903:TRS:C3	2.14	0.60
1:B:28:ARG:NH1	4:B:903:TRS:H32	2.13	0.59
1:A:462:ARG:NH2	5:A:2261:HOH:O	2.34	0.59
1:A:154:THR:O	1:A:169:GLN:O	2.21	0.58
1:B:479:PRO:HB3	3:B:906:GOL:H31	1.87	0.56
3:A:906:GOL:H32	5:A:2277:HOH:O	2.05	0.55
1:B:28:ARG:NH2	4:B:903:TRS:H21	2.17	0.55
1:B:24:GLN:HG2	4:B:903:TRS:O3	2.08	0.54
1:A:24:GLN:CG	3:A:903:GOL:O3	2.52	0.54
1:A:735:ASN:HD22	1:A:738:LEU:H	1.54	0.53
1:B:735:ASN:HD22	1:B:738:LEU:H	1.55	0.53
1:B:675:ARG:HG3	1:B:756[A]:ARG:HG3	1.89	0.52
3:A:903:GOL:H11	1:B:382:LYS:HZ2	1.75	0.52
1:B:28:ARG:HH22	4:B:903:TRS:H32	1.74	0.51
1:B:160:VAL:O	1:B:162:THR:N	2.44	0.51
1:A:28:ARG:HH12	3:A:903:GOL:C3	2.25	0.50
1:A:735:ASN:ND2	1:A:738:LEU:H	2.09	0.50
1:B:356:GLY:HA2	1:B:363:HIS:CE1	2.46	0.50
1:A:-6:VAL:HG23	1:A:-5:LEU:HD13	1.93	0.50
1:A:546:ARG:HH22	3:A:907:GOL:H11	1.77	0.49
1:A:356:GLY:HA2	1:A:363:HIS:CE1	2.47	0.49
1:A:521:THR:HG23	1:A:522:VAL:N	2.27	0.49
1:A:466:GLU:OE1	1:A:495:ARG:HD2	2.13	0.49
1:B:21:ALA:H	1:B:24:GLN:NE2	2.10	0.49
1:A:224:HIS:CE1	1:A:267:ASP:HB3	2.49	0.48
1:A:673:ASN:HA	1:A:757:SER:O	2.13	0.48
1:A:479:PRO:CB	3:A:906:GOL:H31	2.35	0.48
1:A:382:LYS:NZ	4:B:903:TRS:H22	2.29	0.48
1:A:349:HIS:N	1:A:350:PRO:CD	2.78	0.47
1:B:523:SER:HB3	1:B:599:ASN:OD1	2.15	0.46
1:A:860:ASP:OD2	1:A:862:SER:OG	2.32	0.46
1:B:155:GLY:C	1:B:157:ASN:N	2.69	0.46
1:A:541:ILE:HD13	1:A:594:LEU:HD11	1.99	0.45
1:A:425[A]:LYS:HA	1:A:425[A]:LYS:HD3	1.85	0.44
1:A:378:HIS:ND1	4:B:903:TRS:O1	2.50	0.44
1:B:531[A]:CYS:SG	1:B:538:LYS:HG3	2.58	0.44
1:A:28:ARG:HH12	3:A:903:GOL:H31	1.82	0.44
1:A:333:TYR:CD1	1:A:839:ARG:HG3	2.52	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:620:LYS:HB3	1:B:620:LYS:HE3	1.71	0.44
1:A:382:LYS:HE2	1:A:382:LYS:HB3	1.70	0.43
1:A:688[B]:HIS:CE1	3:A:908:GOL:H32	2.54	0.43
1:B:673:ASN:HA	1:B:757:SER:O	2.19	0.43
1:B:3:THR:O	1:B:3:THR:HG23	2.18	0.42
1:B:150:GLY:HA3	1:B:283:ILE:HG22	2.01	0.42
1:A:67:GLU:OE2	1:B:336:ASP:OD2	2.37	0.42
1:B:349:HIS:N	1:B:350:PRO:CD	2.82	0.42
1:B:632:ILE:HG23	5:B:2300:HOH:O	2.18	0.42
1:B:812[B]:ASN:OD1	1:B:814:HIS:HD2	2.03	0.42
1:A:202:GLU:N	3:A:905:GOL:H31	2.18	0.41
1:A:24:GLN:HE21	3:A:903:GOL:C3	2.34	0.41
3:A:903:GOL:H32	1:B:379:TYR:HD1	1.85	0.41
1:B:532:LEU:HD21	1:B:606:LEU:HD23	2.02	0.41
1:B:33:ILE:HB	1:B:61:TYR:CE1	2.55	0.40
1:A:153:PRO:HB2	1:A:219:PHE:CD1	2.57	0.40
1:A:435:LYS:HE2	1:A:435:LYS:HB3	1.82	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	910/923 (99%)	885 (97%)	23 (2%)	2 (0%)	47	38
1	B	893/923 (97%)	866 (97%)	23 (3%)	4 (0%)	34	24
All	All	1803/1846 (98%)	1751 (97%)	46 (3%)	6 (0%)	41	31

All (6) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	161	ASP

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	18	TRP
1	B	18	TRP
1	A	85	PRO
1	B	85	PRO
1	B	84	GLY

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	790/810 (98%)	781 (99%)	9 (1%)	73	73
1	B	775/810 (96%)	764 (99%)	11 (1%)	67	65
All	All	1565/1620 (97%)	1545 (99%)	20 (1%)	71	68

All (20) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	-5	LEU
1	A	180	ASN
1	A	186	LEU
1	A	448	LEU
1	A	464	GLU
1	A	494	LEU
1	A	495	ARG
1	A	612	PHE
1	A	824	TYR
1	B	151	ILE
1	B	448	LEU
1	B	494	LEU
1	B	553	GLN
1	B	612	PHE
1	B	620	LYS
1	B	756[A]	ARG
1	B	756[B]	ARG
1	B	824	TYR

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	862	SER
1	B	871	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	402	GLN
1	A	735	ASN
1	B	735	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 15 ligands modelled in this entry, 2 are monoatomic - leaving 13 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	GOL	A	906	-	5,5,5	0.48	0	5,5,5	0.62	0
3	GOL	A	908	-	5,5,5	0.42	0	5,5,5	0.57	0
3	GOL	A	909	-	5,5,5	0.34	0	5,5,5	0.36	0
3	GOL	B	907	-	5,5,5	0.38	0	5,5,5	0.30	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	GOL	B	906	-	5,5,5	0.40	0	5,5,5	0.40	0
3	GOL	A	903	-	5,5,5	0.36	0	5,5,5	0.81	0
3	GOL	B	908	-	5,5,5	0.39	0	5,5,5	0.44	0
4	TRS	B	903	-	7,7,7	0.49	0	9,9,9	0.26	0
3	GOL	A	905	-	5,5,5	0.43	0	5,5,5	0.54	0
3	GOL	B	905	-	5,5,5	0.40	0	5,5,5	0.44	0
3	GOL	B	904	-	5,5,5	0.46	0	5,5,5	0.56	0
3	GOL	A	907	-	5,5,5	0.37	0	5,5,5	0.36	0
3	GOL	A	904	-	5,5,5	0.40	0	5,5,5	0.64	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	A	906	-	-	4/4/4/4	-
3	GOL	A	908	-	-	4/4/4/4	-
3	GOL	A	909	-	-	0/4/4/4	-
3	GOL	B	907	-	-	0/4/4/4	-
3	GOL	B	906	-	-	2/4/4/4	-
3	GOL	A	903	-	-	0/4/4/4	-
3	GOL	B	908	-	-	0/4/4/4	-
4	TRS	B	903	-	-	9/9/9/9	-
3	GOL	A	905	-	-	1/4/4/4	-
3	GOL	B	905	-	-	4/4/4/4	-
3	GOL	B	904	-	-	0/4/4/4	-
3	GOL	A	907	-	-	4/4/4/4	-
3	GOL	A	904	-	-	1/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (29) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	906	GOL	O1-C1-C2-C3
3	A	907	GOL	O1-C1-C2-C3
3	A	908	GOL	O1-C1-C2-C3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
3	A	908	GOL	C1-C2-C3-O3
3	B	905	GOL	C1-C2-C3-O3
3	B	905	GOL	O2-C2-C3-O3
3	B	906	GOL	O1-C1-C2-O2
4	B	903	TRS	C2-C-C1-O1
4	B	903	TRS	C3-C-C1-O1
4	B	903	TRS	N-C-C1-O1
4	B	903	TRS	C1-C-C3-O3
4	B	903	TRS	C2-C-C3-O3
4	B	903	TRS	N-C-C3-O3
3	B	905	GOL	O1-C1-C2-O2
3	A	904	GOL	O1-C1-C2-C3
3	A	906	GOL	C1-C2-C3-O3
3	A	907	GOL	C1-C2-C3-O3
3	B	905	GOL	O1-C1-C2-C3
3	B	906	GOL	O1-C1-C2-C3
3	A	906	GOL	O1-C1-C2-O2
3	A	908	GOL	O1-C1-C2-O2
3	A	908	GOL	O2-C2-C3-O3
4	B	903	TRS	C1-C-C2-O2
3	A	906	GOL	O2-C2-C3-O3
3	A	907	GOL	O1-C1-C2-O2
3	A	905	GOL	O1-C1-C2-C3
3	A	907	GOL	O2-C2-C3-O3
4	B	903	TRS	N-C-C2-O2
4	B	903	TRS	C3-C-C2-O2

There are no ring outliers.

8 monomers are involved in 34 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	906	GOL	4	0
3	A	908	GOL	3	0
3	B	906	GOL	2	0
3	A	903	GOL	8	0
4	B	903	TRS	12	0
3	A	905	GOL	3	0
3	A	907	GOL	1	0
3	A	904	GOL	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	905/923 (98%)	-0.24	9 (0%) 82 84	10, 15, 24, 33	0
1	B	891/923 (96%)	-0.14	24 (2%) 54 57	9, 14, 25, 56	0
All	All	1796/1846 (97%)	-0.19	33 (1%) 68 71	9, 14, 24, 56	0

All (33) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	160	VAL	14.7
1	B	162	THR	11.3
1	B	161	ASP	10.0
1	B	163	SER	8.5
1	B	156	PHE	7.8
1	A	-6	VAL	7.3
1	B	159	GLN	6.5
1	B	157	ASN	5.3
1	B	158	ASN	4.8
1	B	164	GLU	4.5
1	B	275	ASN	3.9
1	B	527	ASP	3.8
1	A	160	VAL	3.6
1	B	167	SER	3.5
1	B	154	THR	3.5
1	B	522	VAL	3.4
1	B	274	GLU	3.3
1	A	275	ASN	3.2
1	B	155	GLY	3.2
1	B	706	ALA	3.2
1	A	529	ASN	3.1
1	A	-5	LEU	3.1
1	B	529	ASN	3.0
1	A	448	LEU	2.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	449	SER	2.8
1	A	399	MET	2.8
1	B	447	ARG	2.6
1	B	523	SER	2.6
1	B	3	THR	2.3
1	B	449	SER	2.2
1	B	528	SER	2.2
1	B	867	ASN	2.2
1	A	887	GLN	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	GOL	A	908	6/6	0.58	0.34	39,39,41,43	0
3	GOL	B	904	6/6	0.73	0.27	32,33,33,33	0
3	GOL	A	906	6/6	0.74	0.22	24,26,26,29	0
3	GOL	B	906	6/6	0.75	0.23	27,29,30,33	0
3	GOL	A	905	6/6	0.77	0.22	31,33,34,37	0
3	GOL	B	905	6/6	0.79	0.27	39,40,40,40	0
3	GOL	A	907	6/6	0.79	0.17	37,38,39,40	0
3	GOL	A	909	6/6	0.80	0.18	49,49,50,50	0
4	TRS	B	903	8/8	0.83	0.24	29,29,31,31	0
3	GOL	A	903	6/6	0.85	0.18	24,26,27,29	0
3	GOL	A	904	6/6	0.86	0.18	16,24,25,28	0
3	GOL	B	907	6/6	0.88	0.15	38,38,39,39	0
3	GOL	B	908	6/6	0.89	0.16	16,26,27,27	0
2	ZN	A	902	1/1	0.99	0.03	29,29,29,29	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	ZN	B	902	1/1	1.00	0.03	28,28,28,28	0

6.5 Other polymers [i](#)

There are no such residues in this entry.