



wwPDB X-ray Structure Validation Summary Report ⓘ

Jan 6, 2021 – 10:04 PM GMT

PDB ID : 6XVI
Title : Crystal structure of Megabody Mb-Nb207-c7HopQ_A12
Authors : Steyaert, J.; Uchanski, T.; Fischer, B.
Deposited on : 2020-01-22
Resolution : 2.60 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.16
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.16


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3163 (2.60-2.60)
Clashscore	141614	3518 (2.60-2.60)
Ramachandran outliers	138981	3455 (2.60-2.60)
Sidechain outliers	138945	3455 (2.60-2.60)
RSRZ outliers	127900	3104 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	521	 70% 13% 16%
1	B	521	 70% 11% 19%
1	C	521	 69% 13% 17%
1	D	521	 68% 12% 20%

2 Entry composition i

There are 2 unique types of molecules in this entry. The entry contains 13015 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Outer membrane protein,Outer membrane protein,Mb-Nb207-c7HopQ_A12.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	437	3310	2050	585	662	13	0	0	0
1	B	424	3207	1988	564	642	13	0	0	0
1	C	430	3255	2012	576	654	13	0	0	0
1	D	418	3167	1960	557	637	13	0	0	0

There are 52 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	GLN	-	expression tag	UNP B5Z8H1
A	2	VAL	-	expression tag	UNP B5Z8H1
A	3	GLN	-	expression tag	UNP B5Z8H1
A	4	LEU	-	expression tag	UNP B5Z8H1
A	5	VAL	-	expression tag	UNP B5Z8H1
A	6	GLU	-	expression tag	UNP B5Z8H1
A	7	SER	-	expression tag	UNP B5Z8H1
A	8	GLY	-	expression tag	UNP B5Z8H1
A	9	GLY	-	expression tag	UNP B5Z8H1
A	10	GLY	-	expression tag	UNP B5Z8H1
A	11	LEU	-	expression tag	UNP B5Z8H1
A	12	VAL	-	expression tag	UNP B5Z8H1
A	13	ARG	-	expression tag	UNP B5Z8H1
B	1	GLN	-	expression tag	UNP B5Z8H1
B	2	VAL	-	expression tag	UNP B5Z8H1
B	3	GLN	-	expression tag	UNP B5Z8H1
B	4	LEU	-	expression tag	UNP B5Z8H1
B	5	VAL	-	expression tag	UNP B5Z8H1
B	6	GLU	-	expression tag	UNP B5Z8H1
B	7	SER	-	expression tag	UNP B5Z8H1

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	8	GLY	-	expression tag	UNP B5Z8H1
B	9	GLY	-	expression tag	UNP B5Z8H1
B	10	GLY	-	expression tag	UNP B5Z8H1
B	11	LEU	-	expression tag	UNP B5Z8H1
B	12	VAL	-	expression tag	UNP B5Z8H1
B	13	ARG	-	expression tag	UNP B5Z8H1
C	1	GLN	-	expression tag	UNP B5Z8H1
C	2	VAL	-	expression tag	UNP B5Z8H1
C	3	GLN	-	expression tag	UNP B5Z8H1
C	4	LEU	-	expression tag	UNP B5Z8H1
C	5	VAL	-	expression tag	UNP B5Z8H1
C	6	GLU	-	expression tag	UNP B5Z8H1
C	7	SER	-	expression tag	UNP B5Z8H1
C	8	GLY	-	expression tag	UNP B5Z8H1
C	9	GLY	-	expression tag	UNP B5Z8H1
C	10	GLY	-	expression tag	UNP B5Z8H1
C	11	LEU	-	expression tag	UNP B5Z8H1
C	12	VAL	-	expression tag	UNP B5Z8H1
C	13	ARG	-	expression tag	UNP B5Z8H1
D	1	GLN	-	expression tag	UNP B5Z8H1
D	2	VAL	-	expression tag	UNP B5Z8H1
D	3	GLN	-	expression tag	UNP B5Z8H1
D	4	LEU	-	expression tag	UNP B5Z8H1
D	5	VAL	-	expression tag	UNP B5Z8H1
D	6	GLU	-	expression tag	UNP B5Z8H1
D	7	SER	-	expression tag	UNP B5Z8H1
D	8	GLY	-	expression tag	UNP B5Z8H1
D	9	GLY	-	expression tag	UNP B5Z8H1
D	10	GLY	-	expression tag	UNP B5Z8H1
D	11	LEU	-	expression tag	UNP B5Z8H1
D	12	VAL	-	expression tag	UNP B5Z8H1
D	13	ARG	-	expression tag	UNP B5Z8H1

- Molecule 2 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	36	Total O 36 36	0	0
2	B	21	Total O 21 21	0	0
2	C	9	Total O 9 9	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	D	10	Total	O	0	0
			10	10		

4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	96.05Å 117.38Å 96.84Å 90.00° 90.72° 90.00°	Depositor
Resolution (Å)	29.62 – 2.60 29.62 – 2.60	Depositor EDS
% Data completeness (in resolution range)	94.6 (29.62-2.60) 94.6 (29.62-2.60)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.10 (at 2.61Å)	Xtriage
Refinement program	PHENIX 1.12_2829	Depositor
R, R_{free}	0.224 , 0.250 0.224 , 0.251	Depositor DCC
R_{free} test set	3141 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	66.2	Xtriage
Anisotropy	0.039	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 33.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.000 for l,k,-h 0.074 for h,-k,-l 0.019 for l,-k,h	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	13015	wwPDB-VP
Average B, all atoms (Å ²)	76.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.60% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.64	0/3358	0.72	0/4547
1	B	0.62	0/3253	0.68	0/4402
1	C	0.62	0/3300	0.71	0/4466
1	D	0.57	0/3212	0.71	0/4348
All	All	0.61	0/13123	0.71	0/17763

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3310	0	3246	45	0
1	B	3207	0	3136	36	0
1	C	3255	0	3184	53	0
1	D	3167	0	3091	59	0
2	A	36	0	0	1	0
2	B	21	0	0	0	0
2	C	9	0	0	1	0
2	D	10	0	0	0	0
All	All	13015	0	12657	189	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

The worst 5 of 189 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:306:TYR:CD2	1:D:351:LYS:HA	1.57	1.37
1:D:215:ILE:HD11	1:D:266:ARG:NH1	1.72	1.03
1:D:306:TYR:CE2	1:D:351:LYS:HA	1.95	1.01
1:D:306:TYR:CD2	1:D:351:LYS:CA	2.49	0.95
1:D:215:ILE:HD11	1:D:266:ARG:HH12	1.25	0.94

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	423/521 (81%)	412 (97%)	11 (3%)	0	100	100
1	B	408/521 (78%)	397 (97%)	11 (3%)	0	100	100
1	C	416/521 (80%)	401 (96%)	15 (4%)	0	100	100
1	D	404/521 (78%)	397 (98%)	6 (2%)	1 (0%)	47	71
All	All	1651/2084 (79%)	1607 (97%)	43 (3%)	1 (0%)	51	75

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	D	71	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	355/427 (83%)	350 (99%)	5 (1%)	67	85
1	B	342/427 (80%)	337 (98%)	5 (2%)	65	83
1	C	347/427 (81%)	338 (97%)	9 (3%)	46	72
1	D	338/427 (79%)	332 (98%)	6 (2%)	59	80
All	All	1382/1708 (81%)	1357 (98%)	25 (2%)	59	80

5 of 25 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	258	GLN
1	C	349	THR
1	D	479	TYR
1	C	307	THR
1	C	350	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	107	GLN
1	C	254	GLN
1	C	326	ASN
1	C	462	ASN
1	D	262	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2	OWAB(Å ²)	Q < 0.9
1	A	437/521 (83%)	0.36	4 (0%) 84 82	37, 68, 105, 155	0
1	B	424/521 (81%)	0.42	8 (1%) 66 62	43, 69, 110, 139	0
1	C	430/521 (82%)	0.55	23 (5%) 26 20	42, 72, 126, 147	0
1	D	418/521 (80%)	0.58	25 (5%) 21 16	46, 78, 131, 182	0
All	All	1709/2084 (82%)	0.47	60 (3%) 44 36	37, 72, 121, 182	0

The worst 5 of 60 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	217	GLN	7.2
1	C	132	MET	6.1
1	D	310	PRO	5.7
1	A	311	GLY	4.9
1	C	312	GLU	4.8

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers

There are no such residues in this entry.