



wwPDB X-ray Structure Validation Summary Report ⓘ

Nov 14, 2023 – 12:50 AM JST

PDB ID : 5YMT
Title : Functional and structural characterization of P[19] rotavirus VP8* interaction with histo-blood group antigens
Authors : Sun, X.; Duan, Z.
Deposited on : 2017-10-22
Resolution : 2.20 Å (reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4898 (2.20-2.20)
Clashscore	141614	5594 (2.20-2.20)
Ramachandran outliers	138981	5503 (2.20-2.20)
Sidechain outliers	138945	5504 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	160	84% 16% .
1	B	160	82% 17% .
1	C	160	81% 18% .
1	D	160	83% 16% .
1	E	160	85% 14% .
1	F	160	81% 19% .
1	G	160	70% 29% .

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	H	160	 82% 18% .
1	I	160	 72% 28% .
1	J	160	 82% 18% .
1	K	160	 86% 14%
1	L	160	 76% 23% .
1	M	160	 81% 19% .
1	N	160	 69% 30% .
2	O	4	 25% 75%
2	P	4	 100%
2	Q	4	 75% 25%

2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 18722 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Outer capsid protein VP4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	160	1279	808	216	252	3	0	0	0
1	B	160	1279	808	216	252	3	0	0	0
1	C	160	1287	812	218	254	3	0	1	0
1	D	160	1279	808	216	252	3	0	0	0
1	E	160	1279	808	216	252	3	0	0	0
1	F	160	1279	808	216	252	3	0	0	0
1	G	160	1279	808	216	252	3	0	0	0
1	H	160	1279	808	216	252	3	0	0	0
1	I	160	1279	808	216	252	3	0	0	0
1	J	160	1279	808	216	252	3	0	0	0
1	K	160	1279	808	216	252	3	0	0	0
1	L	160	1279	808	216	252	3	0	0	0
1	M	160	1279	808	216	252	3	0	0	0
1	N	160	1279	808	216	252	3	0	0	0

- Molecule 2 is an oligosaccharide called beta-D-galactopyranose-(1-3)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-3)-beta-D-galactopyranose-(1-4)-alpha-D-glucopyranose.



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace	
2	O	4	Total	C	N	O	0	0	0
			48	26	1	21			
2	P	4	Total	C	N	O	0	0	0
			48	26	1	21			
2	Q	4	Total	C	N	O	0	0	0
			48	26	1	21			

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	94	Total	O	0	0
			94	94		
3	B	79	Total	O	0	0
			79	79		
3	C	66	Total	O	0	0
			66	66		
3	D	52	Total	O	0	0
			52	52		
3	E	61	Total	O	0	0
			61	61		
3	F	34	Total	O	0	0
			34	34		
3	G	30	Total	O	0	0
			30	30		
3	H	39	Total	O	0	0
			39	39		
3	I	26	Total	O	0	0
			26	26		
3	J	42	Total	O	0	0
			42	42		
3	K	28	Total	O	0	0
			28	28		
3	L	49	Total	O	0	0
			49	49		
3	M	34	Total	O	0	0
			34	34		
3	N	30	Total	O	0	0
			30	30		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Outer capsid protein VP4

Chain A:  84% 16%



- Molecule 1: Outer capsid protein VP4

Chain B:  82% 17%



- Molecule 1: Outer capsid protein VP4

Chain C:  81% 18%



- Molecule 1: Outer capsid protein VP4

Chain D:  83% 16%



- Molecule 1: Outer capsid protein VP4

Chain E:  85% 14%



- Molecule 1: Outer capsid protein VP4

Chain F:  81% 19%



- Molecule 1: Outer capsid protein VP4



- Molecule 1: Outer capsid protein VP4



- Molecule 1: Outer capsid protein VP4



- Molecule 1: Outer capsid protein VP4




- Molecule 1: Outer capsid protein VP4



- Molecule 1: Outer capsid protein VP4





- Molecule 1: Outer capsid protein VP4

Chain M: 81% 19%



- Molecule 1: Outer capsid protein VP4

Chain N: 69% 30%



- Molecule 2: beta-D-galactopyranose-(1-3)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-3)-beta-D-galactopyranose-(1-4)-alpha-D-glucopyranose

Chain O: 25% 75%



- Molecule 2: beta-D-galactopyranose-(1-3)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-3)-beta-D-galactopyranose-(1-4)-alpha-D-glucopyranose

Chain P: 100%



- Molecule 2: beta-D-galactopyranose-(1-3)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-3)-beta-D-galactopyranose-(1-4)-alpha-D-glucopyranose

Chain Q: 75% 25%



4 Data and refinement statistics

Property	Value	Source
Space group	P 32	Depositor
Cell constants a, b, c, α , β , γ	131.67Å 131.67Å 150.53Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	39.91 – 2.20 39.91 – 2.20	Depositor EDS
% Data completeness (in resolution range)	89.5 (39.91-2.20) 89.5 (39.91-2.20)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.96 (at 2.20Å)	Xtriage
Refinement program	PHENIX 1.11.1_2575	Depositor
R, R_{free}	0.226 , 0.267 0.226 , 0.267	Depositor DCC
R_{free} test set	6659 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å ²)	29.3	Xtriage
Anisotropy	0.170	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 19.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.35$, $\langle L^2 \rangle = 0.18$	Xtriage
Estimated twinning fraction	0.267 for -h,-k,l 0.206 for h,-h-k,-l 0.205 for -k,-h,-l	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	18722	wwPDB-VP
Average B, all atoms (Å ²)	35.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 1.91% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NAG, GLC, GAL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.37	0/1310	0.59	0/1786
1	B	0.37	0/1310	0.57	0/1786
1	C	0.36	0/1318	0.58	1/1797 (0.1%)
1	D	0.36	0/1310	0.56	1/1786 (0.1%)
1	E	0.36	0/1310	0.57	0/1786
1	F	0.33	0/1310	0.57	1/1786 (0.1%)
1	G	0.33	0/1310	0.53	0/1786
1	H	0.33	0/1310	0.55	1/1786 (0.1%)
1	I	0.38	1/1310 (0.1%)	0.56	1/1786 (0.1%)
1	J	0.34	0/1310	0.61	2/1786 (0.1%)
1	K	0.33	0/1310	0.53	1/1786 (0.1%)
1	L	0.34	0/1310	0.55	0/1786
1	M	0.33	0/1310	0.55	1/1786 (0.1%)
1	N	0.52	1/1310 (0.1%)	0.58	0/1786
All	All	0.36	2/18348 (0.0%)	0.57	9/25015 (0.0%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	N	87	GLU	CG-CD	14.58	1.73	1.51
1	I	101	VAL	CB-CG1	6.26	1.66	1.52

The worst 5 of 9 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	F	30	LEU	CA-CB-CG	8.54	134.95	115.30
1	J	2	LEU	CA-CB-CG	8.12	133.97	115.30
1	C	30	LEU	CA-CB-CG	7.43	132.40	115.30
1	D	30	LEU	CA-CB-CG	7.08	131.57	115.30
1	H	30	LEU	CA-CB-CG	6.74	130.80	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1279	0	1237	22	0
1	B	1279	0	1237	21	0
1	C	1287	0	1242	24	0
1	D	1279	0	1237	23	0
1	E	1279	0	1237	14	2
1	F	1279	0	1237	20	1
1	G	1279	0	1237	34	1
1	H	1279	0	1237	23	0
1	I	1279	0	1237	36	1
1	J	1279	0	1237	20	0
1	K	1279	0	1237	17	0
1	L	1279	0	1237	36	0
1	M	1279	0	1237	24	1
1	N	1279	0	1237	34	1
2	O	48	0	42	0	0
2	P	48	0	42	1	0
2	Q	48	0	42	0	0
3	A	94	0	0	6	0
3	B	79	0	0	2	0
3	C	66	0	0	6	1
3	D	52	0	0	2	0
3	E	61	0	0	5	0
3	F	34	0	0	4	0
3	G	30	0	0	8	0
3	H	39	0	0	5	0
3	I	26	0	0	5	0
3	J	42	0	0	7	1
3	K	28	0	0	7	0
3	L	49	0	0	8	1
3	M	34	0	0	5	0
3	N	30	0	0	7	0
All	All	18722	0	17449	332	5

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

The worst 5 of 332 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:N:77:MET:SD	3:N:222:HOH:O	1.96	1.24
1:A:119:ASN:OD1	3:A:201:HOH:O	1.80	1.00
1:L:85:ASN:ND2	3:L:201:HOH:O	1.97	0.97
1:D:149:GLU:OE2	3:D:701:HOH:O	1.84	0.95
1:G:2:LEU:O	3:G:701:HOH:O	1.83	0.95

All (5) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:F:128:SER:OG	1:N:82:ASN:O[3_555]	2.11	0.09
3:C:730:HOH:O	3:J:219:HOH:O[2_554]	2.13	0.07
1:E:82:ASN:O	1:G:128:SER:OG[2_454]	2.15	0.05
1:E:69:ASN:OD1	3:L:202:HOH:O[2_454]	2.17	0.03
1:I:82:ASN:O	1:M:128:SER:OG[3_445]	2.18	0.02

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	158/160 (99%)	152 (96%)	6 (4%)	0	100	100
1	B	158/160 (99%)	149 (94%)	9 (6%)	0	100	100
1	C	159/160 (99%)	154 (97%)	5 (3%)	0	100	100
1	D	158/160 (99%)	154 (98%)	4 (2%)	0	100	100
1	E	158/160 (99%)	152 (96%)	6 (4%)	0	100	100
1	F	158/160 (99%)	155 (98%)	3 (2%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	G	158/160 (99%)	154 (98%)	4 (2%)	0	100	100
1	H	158/160 (99%)	152 (96%)	6 (4%)	0	100	100
1	I	158/160 (99%)	151 (96%)	7 (4%)	0	100	100
1	J	158/160 (99%)	154 (98%)	4 (2%)	0	100	100
1	K	158/160 (99%)	151 (96%)	7 (4%)	0	100	100
1	L	158/160 (99%)	151 (96%)	7 (4%)	0	100	100
1	M	158/160 (99%)	152 (96%)	6 (4%)	0	100	100
1	N	158/160 (99%)	154 (98%)	4 (2%)	0	100	100
All	All	2213/2240 (99%)	2135 (96%)	78 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	146/146 (100%)	145 (99%)	1 (1%)	84	91
1	B	146/146 (100%)	144 (99%)	2 (1%)	67	80
1	C	147/146 (101%)	143 (97%)	4 (3%)	44	57
1	D	146/146 (100%)	144 (99%)	2 (1%)	67	80
1	E	146/146 (100%)	145 (99%)	1 (1%)	84	91
1	F	146/146 (100%)	145 (99%)	1 (1%)	84	91
1	G	146/146 (100%)	144 (99%)	2 (1%)	67	80
1	H	146/146 (100%)	145 (99%)	1 (1%)	84	91
1	I	146/146 (100%)	146 (100%)	0	100	100
1	J	146/146 (100%)	146 (100%)	0	100	100
1	K	146/146 (100%)	145 (99%)	1 (1%)	84	91
1	L	146/146 (100%)	145 (99%)	1 (1%)	84	91
1	M	146/146 (100%)	144 (99%)	2 (1%)	67	80

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	N	146/146 (100%)	145 (99%)	1 (1%)	84	91
All	All	2045/2044 (100%)	2026 (99%)	19 (1%)	78	88

5 of 19 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	K	114	HIS
1	M	114	HIS
1	N	146	ARG
1	M	30	LEU
1	D	116	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	119	ASN
1	F	114	HIS
1	I	64	GLN
1	J	50	ASN
1	L	55	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

12 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	GLC	O	1	2	12,12,12	0.69	0	17,17,17	0.82	0
2	GAL	O	2	2	11,11,12	0.81	1 (9%)	15,15,17	0.89	0
2	NAG	O	3	2	14,14,15	0.41	0	17,19,21	0.76	1 (5%)
2	GAL	O	4	2	11,11,12	0.86	1 (9%)	15,15,17	1.05	1 (6%)
2	GLC	P	1	2	12,12,12	0.62	0	17,17,17	0.91	1 (5%)
2	GAL	P	2	2	11,11,12	0.80	1 (9%)	15,15,17	1.01	2 (13%)
2	NAG	P	3	2	14,14,15	0.29	0	17,19,21	0.62	0
2	GAL	P	4	2	11,11,12	0.94	1 (9%)	15,15,17	1.11	0
2	GLC	Q	1	2	12,12,12	0.66	0	17,17,17	0.88	0
2	GAL	Q	2	2	11,11,12	0.46	0	15,15,17	1.07	0
2	NAG	Q	3	2	14,14,15	0.16	0	17,19,21	0.57	0
2	GAL	Q	4	2	11,11,12	0.92	1 (9%)	15,15,17	1.22	3 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GLC	O	1	2	-	0/2/22/22	0/1/1/1
2	GAL	O	2	2	-	0/2/19/22	0/1/1/1
2	NAG	O	3	2	-	0/6/23/26	0/1/1/1
2	GAL	O	4	2	-	2/2/19/22	0/1/1/1
2	GLC	P	1	2	-	0/2/22/22	0/1/1/1
2	GAL	P	2	2	-	1/2/19/22	0/1/1/1
2	NAG	P	3	2	-	0/6/23/26	0/1/1/1
2	GAL	P	4	2	-	2/2/19/22	0/1/1/1
2	GLC	Q	1	2	-	0/2/22/22	0/1/1/1
2	GAL	Q	2	2	-	0/2/19/22	0/1/1/1
2	NAG	Q	3	2	-	0/6/23/26	0/1/1/1
2	GAL	Q	4	2	-	2/2/19/22	0/1/1/1

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	Q	4	GAL	O5-C5	2.63	1.48	1.43
2	P	2	GAL	O5-C5	2.08	1.47	1.43
2	P	4	GAL	O5-C5	2.06	1.47	1.43
2	O	2	GAL	C2-C3	-2.06	1.49	1.52
2	O	4	GAL	O5-C5	2.02	1.47	1.43

The worst 5 of 8 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	Q	4	GAL	O5-C1-C2	-2.57	106.81	110.77
2	P	2	GAL	C1-O5-C5	2.49	115.57	112.19
2	P	2	GAL	C2-C3-C4	-2.27	106.97	110.89
2	Q	4	GAL	C1-O5-C5	2.24	115.22	112.19
2	O	4	GAL	C1-O5-C5	2.16	115.12	112.19

There are no chirality outliers.

5 of 7 torsion outliers are listed below:



Mol	Chain	Res	Type	Atoms
2	P	4	GAL	O5-C5-C6-O6
2	Q	4	GAL	O5-C5-C6-O6
2	P	4	GAL	C4-C5-C6-O6
2	O	4	GAL	O5-C5-C6-O6
2	Q	4	GAL	C4-C5-C6-O6


There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	P	3	NAG	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

Unable to reproduce the depositors R factor - this section is therefore empty.


6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

Unable to reproduce the depositors R factor - this section is therefore empty.

6.3 Carbohydrates [i](#)




Unable to reproduce the depositors R factor - this section is therefore empty.

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around Chain P:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around Chain Q:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.4 Ligands

Unable to reproduce the depositors R factor - this section is therefore empty.

6.5 Other polymers

Unable to reproduce the depositors R factor - this section is therefore empty.